

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
24 March 2005 (24.03.2005)

PCT

(10) International Publication Number
WO 2005/026362 A2

(51) International Patent Classification⁷: **C12N 15/12**,
C07K 14/47, 14/705, 16/18, 16/28, 14/62, G01N 33/68,
33/50, C12Q 1/68, A61K 38/17, 48/00, C12N 15/11

Molecular/Cancer Biology Laboratory Biomedicum
Helsinki, P.O. Box 63 (Haartmaninkatu 8), FIN-00014
Helsinki (FI).

(21) International Application Number:
PCT/US2004/029245

(74) Agent: **NEVILLE, Katherine, L.**; Marshall, Gerstein
& Borun LLP, 233 S. Wacker Drive, 6300 Sears Tower,
Chicago, IL 60606-6357 (US).

(22) International Filing Date:
8 September 2004 (08.09.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/500,977 8 September 2003 (08.09.2003) US

(71) Applicants (for all designated States except US): **LUD-
WIG INSTITUTE FOR CANCER RESERACH**
[US/US]; 605 THIRD AVENUE, New York, NY 10158
(US). **LICENTIA LTD.** [FI/FI]; Erottajankatu 19 B, 6th
Floor, FIN-00130 Helsinki (FI).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **ALITALO, Kari**
[FI/FI]; Nyrikintie 4A, 02100 Espoo (FI). **MAKINEN,
Taija** [FI/DE]; Department of Molecule Neurobiology,
Max-Planck Institute of Neurobiology, Am Klopfer-
spitz 18A, 82152 Martinsried (DE). **PETROVA, Ta-
tiana** [RU/FI]; Molecular/Cancer Biology Laboratory
Biomedicum Helsinki, P.O. Box 63 (Haartmaninkatu 8),
FIN-00014 University of Helsinki (FI). **SAHARINEN,
Pipsa** [FI/FI]; Molecular/Cancer Biology Laboratory
Biomedicum Helsinki, P.O. Box 63 (Haartmaninkatu 8),
FIN-00014 Helsinki (FI). **SAHARINEN, Juha** [FI/FI];

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI,
SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: LYMPHATIC AND BLOOD ENDOTHELIAL CELL GENES

(57) Abstract: The invention provides polynucleotides and genes that are differentially expressed in lymphatic versus blood vascular endothelial cells. These genes are useful targets for treating diseases involving lymphatic vessels, such as lymphedema, various inflammatory diseases, and cancer metastasis via the lymphatic system.



WO 2005/026362 A2

LYMPHATIC AND BLOOD ENDOTHELIAL CELL GENES

BACKGROUND OF THE INVENTION

Field of the Invention

5 The invention relates to polynucleotides and proteins specifically expressed in lymphatic endothelial cells.

Description of the Related Art

Recent evidence on the association of lymphangiogenic growth factors
10 with intralymphatic growth and metastasis of cancers (Mandriota, et al., *EMBO J.* 20:672-682 (2001); Skobe, et al., *Nat. Med.* 7:192-198 (2001); Stacker, et al., *Nat. Med.* 7:186-191 (2001); Karpanen, et al., *Cancer Res.* 61:1786-1790 (2001)) has raised hopes that lymphatic vessels could be used as an additional target for tumor therapy. Cancer cells spread within the body by direct invasion to surrounding
15 tissues, spreading to body cavities, invasion into the blood vascular system (hematogenous metastasis), as well as spread via the lymphatic system (lymphatic metastasis). Regional lymph node dissemination is the first step in the metastasis of several common cancers and correlates highly with the prognosis of the disease. The lymph nodes that are involved in draining tissue fluid from the tumor area are called
20 sentinel nodes, and diagnostic measures are in place to find these nodes and to remove them in cases of suspected metastasis. However, in spite of its clinical relevance, little is known about the mechanisms leading to metastasis via the bloodstream or via the lymphatics.

Until recently, the lymphatic vessels have received much less attention
25 than blood vessels, despite their importance in medicine. Lymphatic vessels collect protein-rich fluid and white blood cells from the interstitial space of most tissues and transport them as a whitish opaque fluid, the lymph, into the blood circulation. Small lymphatic vessels coalesce into larger vessels, which drain the lymph through the thoracic duct into large veins in the neck region. Lymph nodes serve as filtering
30 stations along the lymphatic vessels and lymph movement is propelled by the contraction of smooth muscles surrounding collecting lymphatic vessels and by bodily

movements, the direction of flow being secured by valves as it is in veins. The lymphatic capillaries are lined by endothelial cells, which have distinct junctions with frequent large interendothelial gaps. The lymphatic capillaries also lack a continuous basement membrane, and are devoid of pericytes. Anchoring filaments connect the abluminal surfaces of lymphatic endothelial cells to the perivascular extracellular matrix and pull to maintain vessel patency in the presence of tissue edema. The absence or obstruction of lymphatic vessels, which is usually the result of an infection, surgery, or radiotherapy and in rare cases, a genetic defect, causes accumulation of a protein-rich fluid in tissues, lymphedema. The lymphatic system is also critical in fat absorption from the gut and in immune responses. Bacteria, viruses, and other foreign materials are taken up by the lymphatic vessels and transported to the lymph nodes, where the foreign material is presented to immune cells and where dendritic cells traverse via the lymphatics. There has been slow progress in the understanding of and ability to manipulate the lymphatic vessels.

Abnormal development or function of the lymphatic ECs can result in tumors or malformations of the lymphatic vessels, such as lymphangiomas or lymphangiectasis. Witte, et al., *Regulation of Angiogenesis* (eds. Goldber, I.D. & Rosen, E.M.) 65-112 (Birkäuser, Basel, Switzerland, 1997). The VEGFR-3 tyrosine kinase receptor is expressed in the normal lymphatic endothelium and is upregulated in many types of vascular tumors, including Kaposi's sarcomas. Jussila, et al., *Cancer Res* 58, 1955-1604 (1998); Partanen, et al., *Cancer* 86:2406-2412 (1999). Absence or dysfunction of lymphatic vessels which can result from an infection, surgery, radiotherapy or from a genetic defect, causes lymphedema, which is characterized by a chronic accumulation of protein-rich fluid in the tissues that leads to swelling. The importance of VEGFR-3 signaling for lymphangiogenesis was revealed in the genetics of familial lymphedema, a disease characterized by a hypoplasia of cutaneous lymphatic vessels, which leads to a disfiguring and disabling swelling of the extremities. Witte, et al., *Regulation of Angiogenesis* (eds. Goldber, I.D. & Rosen, E.M.) 65-112 (Birkäuser, Basel, Switzerland, 1997); Rockson, S.G., *Am. J. Med.* 110, 288-295 (2001). Some members of families with lymphedema are heterozygous for missense mutations of the *VEGFR3* exons encoding the tyrosine

kinase domain, which results in an inactive receptor protein. Karkkainen, et al., *Nature Genet.* 25:153-159 (2000); Irrthum, et al., *Am. J. Hum. Genet.* 67:295-301 (2000).

5 There is a need in the art for information on the transcriptional program which controls the diversity of endothelial cells, and into the mechanisms of angiogenesis and lymphangiogenesis. There is also a need in the art for new vascular markers, which may be used as valuable targets in the study of a number of diseases involving the lymphatic vessels, including tumor metastasis.

SUMMARY OF THE INVENTION

10 The compositions of the present invention include isolated polynucleotides and polypeptides, in particular, lymphatic endothelial genes, isolated polypeptides encoded by these polynucleotides, recombinant DNA molecules, cloned genes or degenerate variants thereof, especially naturally occurring variants such as allelic variants, and antibodies that specifically recognize one or more epitopes
15 present on such polypeptides.

 The compositions of the present invention additionally include vectors, including expression vectors, containing the polynucleotides of the invention, cells genetically engineered to contain such polynucleotides and cells genetically engineered to express such polynucleotides.

20 In selected embodiments, such isolated polynucleotides of the invention represent a polynucleotide comprising a nucleotide sequence selected from the group consisting of SEQ ID NOS: 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 and 41, or a fragment thereof that encodes a corresponding polypeptide.

25 The polynucleotides of the present invention also include, but are not limited to, a polynucleotide that hybridizes to the complement of the nucleotide sequence of SEQ ID NOS: 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 and 41 under highly stringent hybridization conditions; a polynucleotide that hybridizes to the complement of the nucleotide sequence of SEQ ID NOS: 5, 7, 9, 11,
30 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 and 41 under moderately stringent

hybridization conditions; a polynucleotide which is an allelic variant of any polynucleotide recited above; a polynucleotide which encodes a species homologue of any of the proteins recited above; or a polynucleotide that encodes a polypeptide comprising a specific domain or truncation of the polypeptide encoded by any one of
5 SEQ ID NOS: 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 and 41. Exemplary high stringency hybridization conditions are hybridization at 42°C for 20 hours in a solution containing 50% formamide, 5xSSPE, 5x Denhardt's solution, 0.1% SDS and 0.1 mg/ml denatured salmon sperm DNA, with a wash in 1xSSC, 0.1% SDS for 30 minutes at 65°C.

10 Another aspect of the invention is drawn to LEC and BEC polypeptides, including polypeptides encoded by the polynucleotides described above. In some embodiments, the polypeptides are the mature forms of the polypeptides of the invention. Expressly contemplated is a purified and isolated polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID
15 NOS: 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 43, 44, 45, and 46; and a purified and isolated polypeptide comprising an amino acid sequence selected from the group consisting of: (a) SEQ ID NOS: 6, 8, 10, 12, 14, 16, 20, 22, 24, 42 and 43; and (b) an extracellular domain fragment of at least 10 amino acids of an amino acid sequence of (a). Further, this aspect of the invention includes a
20 purified and isolated, soluble polypeptide as described immediately above, comprising an extracellular domain fragment of an amino acid sequence selected from the group consisting of: SEQ ID NOS: 6, 8, 10, 12, 14, 16, 20, 22, 24, 42 and 43 wherein the polypeptide lacks any transmembrane domain. Such a polypeptide may further lack any intracellular domain. Also, the invention contemplates a fusion
25 protein comprising a polypeptide as described above fused, e.g., to an immunoglobulin fragment comprising an immunoglobulin constant region.

Another aspect of the invention is an isolated polypeptide comprising an amino acid sequence at least 95% identical to amino acids 1-1005 of SEQ ID NO: 43. In a related aspect, a composition is contemplated which comprises a soluble
30 polypeptide comprising a fragment of one of the amino acid sequences disclosed herein that comprises a functional protein domain. One exemplary embodiment

comprises a fragment of the amino acid sequence set forth in SEQ ID NO: 43, wherein said fragment lacks the transmembrane and intracellular amino acids from approximately residues 1006-1028 of SEQ ID NO: 43. Further contemplated is an isolated polypeptide comprising an adhesion domain having an amino acid sequence at least 95% identical to a fragment of a polypeptide comprising the amino acid sequence set forth in SEQ ID NO: 20 or SEQ ID NO: 43, wherein said fragment includes at least one vonWillebrand factor domain or at least one Cache domain.

In an additional aspect, the invention contemplates polypeptide fragments of any one of SEQ ID NOS: 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 43, 44, 45, and 46, wherein the fragment comprises at least one protein domain of the given polypeptide sequence, as set out herein. The invention further contemplates polypeptide fragments comprising an amino acid sequence at least 95% identical to a fragment of any one of SEQ ID NOS: 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 43, 44, 45, and 46 comprising a protein domain of the polypeptide set forth herein. Exemplary embodiments include, e.g., polypeptide fragments comprising a protein domain that is a protein kinase domain selected from the group consisting of a protein kinase domain set out in SEQ ID NO: 8, a protein kinase domain set out in SEQ ID NO: 26, a protein kinase domain at approximately amino acids 88-342 of SEQ ID NO: 42, and a protein kinase domain at approximately amino acids 51-317 of SEQ ID NO: 46; a Zinc binding domain selected from the group consisting of a Zinc-binding domain set out in SEQ ID NO: 36, a zinc-binding domain at approximately amino acids 627-561 of SEQ ID NO: 6 and a zinc-binding domain at approximately amino acids 1030-1094 of SEQ ID NO: 45.

Another aspect of the invention is drawn to an isolated polypeptide at least 95% identical to a polypeptide set forth herein, comprising a particular biological activity associated with a protein domain as set forth herein. Exemplary biological activities include ubiquitin ligase activity of a Hect domain in SEQ ID NO: 28; guanine-nucleotide-dissociation-stimulating activity of at least one RCC1-like domain in SEQ ID NO: 28; cation transport activity of a cation transport domain set forth in SEQ ID NO: 12; GTP-binding activity of at least one GTP-binding

domain set forth in SEQ ID NO: 32, SRPY domain activity of at least one SPRY domain of SEQ ID NO: 6, and Kelch domain activity of at least one Kelch domain of SEQ ID NO: 30 or SEQ ID NO: 43. In another aspect, the invention contemplates an isolated polypeptide comprising an amino acid sequence at least 95% identical to a transcription factor selected from the group consisting of SEQ ID NO: 30, SEQ ID NO: 38, SEQ ID NO: 40, and SEQ ID NO: 44, or a protein involved in cytoskeletal regulation selected from the group consisting of SEQ ID NO: 18, SEQ ID NO: 36 and SEQ ID NO: 45.

In a related aspect, the invention provides a composition comprising a polynucleotide or polypeptide or protein as described above and a pharmaceutically acceptable diluent, carrier or adjuvant. Polypeptide compositions of the invention may comprise an acceptable carrier, such as a hydrophilic, *e.g.*, pharmaceutically acceptable, carrier. Further provided is a kit comprising such a composition and a protocol for administering the pharmaceutical composition to a mammalian subject to modulate the lymphatic system in the subject. The invention also provides an antibody that specifically binds to a polypeptide as described above, and that antibody is humanized in some embodiments. Antibodies contemplated by the invention include, for example, monoclonal and polyclonal antibodies, single chain antibodies, chimeric antibodies, bifunctional/bispecific antibodies, humanized antibodies, human antibodies, single and multiple domain antibodies and complementarity-determining region (CDR)-grafted antibodies, including compounds which include CDR sequences that specifically recognize a polypeptide of the invention, provided that such antibodies each specifically bind at least one polypeptide according to the invention. Still further, the invention provides a protein comprising an antigen binding domain of an antibody that specifically binds to a polypeptide as described hereinabove, wherein the protein specifically binds to the polypeptide.

The invention also relates to methods for producing a polypeptide comprising growing a culture of the cells of the invention in a suitable culture medium, and purifying the protein from the culture or from an extract of the cells. In particular, the invention contemplates a method for producing a LEC polypeptide comprising steps of growing a host cell transformed or transfected with an expression

vector as described herein under conditions in which the cell expresses the polypeptide encoded by the polynucleotide.

Methods of identifying the products and compositions described herein are also provided by the invention. In particular, the invention provides a method of identifying a LEC nucleic acid comprising: (a) contacting a biological sample containing a candidate LEC nucleic acid with a polynucleotide comprising a fragment of at least 14 contiguous nucleotides of a sequence selected from the group consisting of SEQ ID NOS: 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 and 41, or a complement thereof, under the following stringent hybridization conditions: (i) hybridization at 42°C for 20 hours in a solution containing 50% formamide, 5xSSPE, 5x Denhardt's solution, 0.1% SDS and 0.1 mg/ml denatured salmon sperm DNA, and (ii) washing for 30 minutes at 65°C in 1xSSC, 0.1% SDS; and (b) detecting hybridization of the candidate LEC nucleic acid and the polynucleotide, thereby identifying a LEC nucleic acid.

The invention also provides a method of identifying a LEC protein comprising: (a) contacting a biological sample containing a candidate LEC protein with a LEC protein binding partner selected from the group consisting of an antibody as described herein or a protein or polypeptide as described herein, under conditions suitable for binding therebetween; and (b) detecting binding between the candidate LEC protein and the LEC binding partner, thereby identifying a LEC protein.

Another related aspect of the invention is a method of identifying a LEC comprising: (a) contacting a biological sample comprising cells with a LEC binding partner under conditions suitable for binding therebetween, wherein the LEC binding partner comprises an antibody that binds to a polypeptide comprising a sequence selected from the group consisting of SEQ ID NOS: 6, 8, 10, 12, 14, 16, 20, 22, 24, 42 and 43, or comprises an antigen binding fragment of the antibody; and (b) identifying a LEC by detecting binding between a cell and the LEC binding partner, where binding of the LEC binding partner to the cell identifies a LEC.

Polynucleotides according to the invention have numerous applications in a variety of techniques known to those skilled in the art of molecular biology. These techniques include use as hybridization probes, use as primers for PCR, use for

chromosome and gene mapping, use in the recombinant production of protein, and use in generation of anti-sense DNA or RNA, their chemical analogs and the like. For example, when the expression of an mRNA is largely restricted to a particular cell or tissue type, such as a lymphatic endothelial cell, polynucleotides of the invention can be used as hybridization probes to detect the presence of the particular cell or tissue mRNA in a sample using, *e.g.*, *in situ* hybridization.

In another aspect, the invention provides a composition comprising an isolated polynucleotide that comprises a nucleotide sequence that encodes a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 43, 44, 45, and 46 and a pharmaceutically acceptable diluent, carrier or adjuvant. In some embodiments, the composition comprises a polynucleotide that comprises a nucleotide sequence selected from the group consisting of SEQ ID NOS: 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 and 41, or a fragment thereof that encodes the polypeptide.

Still another aspect of the invention is an expression vector comprising an expression control sequence operably linked to a polynucleotide that comprises a nucleotide sequence that encodes a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 43, 44, 45, and 46. In some embodiments, the expression vector is a replication-deficient adenoviral or adeno-associated viral vector containing the polynucleotide. A related aspect of the invention is a composition comprising an expression vector as described above and a pharmaceutically acceptable diluent, carrier, or adjuvant. Further, the invention provides a kit comprising the composition containing either the above-described polynucleotide or vector and a pharmaceutically acceptable diluent, carrier or adjuvant, packaged with a protocol for administering the composition to a mammalian subject to modulate the lymphatic system in the subject.

The invention further provides a host cell transformed or transfected with an expression vector as described above.

The polypeptides according to the invention can be used in a variety of conventional procedures and methods that are currently applied to other proteins. In addition, a polypeptide of the invention can be used to generate an antibody that specifically binds the polypeptide.

5 In one aspect of the invention, a method is provided for differentially modulating the growth or differentiation of blood endothelial cells (BEC) or lymphatic endothelial cells (LEC), comprising contacting endothelial cells with a composition comprising an agent that differentially modulates blood or lymphatic endothelial cells, said agent selected from the group consisting of: (a) a polypeptide
10 that comprises an amino acid sequence of a BEC polypeptide or a LEC polypeptide, or an active fragment of the polypeptide ; (b) a polynucleotide that comprises a nucleotide sequence that encodes a polypeptide according to (a); (c) an antibody that specifically binds to a polypeptide according to (a); (d) a polypeptide comprising a fragment of the antibody of (c), wherein the fragment and the antibody bind to the
15 polypeptide; (e) an antisense nucleic acid to a human gene or mRNA encoding the polypeptide of (a); (f) an interfering RNA (RNAi) to a human gene or mRNA encoding the polypeptide of (a). In a related aspect, the method of the invention provides an antibody, a peptide or a small molecular weight compound for differentially modulating the growth or differentiation of blood endothelial cells
20 (BEC) or lymphatic endothelial cells (LEC). The method may involve endothelial cell contact with the composition *ex vivo* or *in vivo*. The composition may comprise a pharmaceutically acceptable diluent, adjuvant, or carrier, and the contacting step may comprise administering the composition to a mammalian subject to differentially modulate BECs or LECs in the mammalian subject.

25 Further, the method may comprise identifying a human subject with a disorder characterized by hyperproliferation of LECs; and administering to the human subject the composition, wherein the agent differentially inhibits LEC growth compared to BEC growth; alternatively the method may comprise identifying a human subject with a disorder characterized by hyperproliferation of LECs; screening
30 LECs of the subject to identify overexpression of a polypeptide set forth in any of Tables 1 and 2 herein; and administering to the human subject the composition,

wherein the agent differentially inhibits LEC growth compared to BEC growth by inhibiting expression of the polypeptide identified by the screening step.

This aspect of the invention also contemplates a method of modulating the growth of lymphatic endothelial cells in a human subject, comprising steps of

5 identifying a human subject with a hypoproliferative lymphatic disorder; screening the subject to identify underexpression or underactivity of a LEC polypeptide set forth in any of Table 1 or Table 2 herein; administering to the human subject the composition, wherein the agent comprises the LEC polypeptide (a) identified by the screening step or an active fragment of the polypeptide, or comprises the

10 polynucleotide (b) that comprises a nucleotide sequence that encodes the polypeptide.

A related aspect of the invention is drawn to a use of an agent for the manufacture of a medicament for the differential modulation of blood vessel endothelial cell (BEC) or lymphatic vessel endothelial cell (LEC) growth or differentiation, the agent selected from the group consisting of: (a) a polypeptide that

15 comprises an amino acid sequence of a BEC polypeptide or a LEC polypeptide, or an active fragment of the polypeptide ; (b) a polynucleotide that comprises a nucleotide sequence that encodes a polypeptide according to (a); (c) an antibody that specifically binds to a polypeptide according to (a); (d) a polypeptide comprising a fragment of the antibody of (c), wherein the fragment and the antibody bind to the polypeptide; (e)

20 an antisense nucleic acid to a human gene or mRNA encoding the polypeptide of (a); (f) an interfering RNA (RNAi) to a human gene or mRNA encoding the polypeptide of (a). It is further contemplated that an antibody, a peptide or a small molecular weight compound may be useful for the manufacture of a medicament for the differential modulation of blood endothelial cell (BEC) or lymphatic endothelial cell

25 (LEC) growth or differentiation. In some embodiments, the medicament may promote growth and/or differentiation of a BEC or LEC; in other embodiments, the medicament inhibits growth and/or differentiation of such a cell.

In another aspect, the invention provides a method of identifying compounds that modulate growth of endothelial cells, comprising culturing

30 endothelial cells in the presence and absence of a compound; and measuring expression of at least one BEC or LEC gene in the cells, wherein the gene is a gene

set forth in any of Tables 1 and 2 herein, wherein a change in expression of at least one BEC gene in the presence compared to the absence of the compound identifies the compound as a modulator of BEC growth, and wherein a change in expression of at least one LEC gene in the presence compared to the absence of the compound
5 identifies the compound as a modulator of LEC growth. The method may be used to screen for a compound that selectively modulates BEC or LEC growth or differentiation, wherein the measuring step comprises measuring expression of at least one BEC gene and at least one LEC gene in the cells, and wherein the method comprises screening for a compound that selectively modulates BEC or LEC growth
10 or differentiation by selecting a compound that differentially modulates expression of the at least one BEC gene compared to expression of the at least one LEC gene. In a related aspect, the method further comprises a second BEC or LEC gene that encodes a polypeptide set forth in any of Tables 3 and 4 of PCT/US03/06900.

Further, the invention comprehends a method or use according to the
15 aspects of the invention described above, wherein the polypeptide is a LEC polypeptide selected from a gene found in any of Tables 1 and 2 herein, and the agent differentially modulates LEC growth or differentiation over BEC growth or differentiation. In some embodiments, the LEC polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 6, 8, 10, 12, 14, 16, 20,
20 22, 24, 42 and 43; in other embodiments, the LEC polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 20 and SEQ ID NO: 32. In these embodiments, an agent may be an antibody that specifically binds to a LEC polypeptide as described above, or a polypeptide fragment of such an antibody. Further, the agent may be an extracellular domain of a polypeptide described above, a
25 polynucleotide encoding an extracellular domain, or an antisense molecule or nucleic acid. Alternatively, the polypeptide is a BEC polypeptide selected from the BEC polypeptides set forth in any of Tables 1 and 2 herein, and the agent differentially modulates BEC growth or differentiation over LEC growth or differentiation. It is further contemplated that the above methods further comprise screening LECs of the
30 subject to identify overexpression of a second polypeptide set forth in any of Tables 3 and 4 of PCT/US03/06900 (specifically incorporated herein by reference).

The methods of the present invention further relate to methods for detecting the presence of the polynucleotides or polypeptides of the invention in a sample. Such methods can, for example, be utilized as part of a prognostic and/or diagnostic evaluation of disorders as recited above and for the identification of
5 subjects exhibiting a predisposition to such conditions. Furthermore, the invention provides methods for evaluating the efficacy of drugs, and monitoring the progress of patients, involved in clinical trials for the treatment of disorders related to lymphatic endothelial cells.

The invention also provides methods for the identification of
10 compounds that modulate the expression of the polynucleotides and/or polypeptides of the invention. Such methods can be utilized, for example, for the identification of compounds that can ameliorate symptoms of disorders related to expression of proteins encoded by any one of SEQ ID NOS: 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 and 41 as recited above. Such methods can include, but are
15 not limited to, assays for identifying compounds and other substances that interact with (*e.g.*, bind to) the polypeptides of the invention.

Further, the invention provides a method of assaying for risk of developing hereditary lymphedema, comprising (a) assaying nucleic acid of a human subject for a mutation that correlates with a hereditary lymphedema phenotype and
20 alters the encoded amino acid sequence of at least one gene allele of the human subject when compared to the amino acid sequence of the polypeptide encoded by a corresponding wild-type gene allele, wherein the wild-type polypeptide is a polypeptide identified in any of Tables 1 and 2 herein. Alternatively, a method of assaying for risk of developing hereditary lymphedema, comprises (a) assaying
25 nucleic acid of a human subject for a mutation that correlates with a hereditary lymphedema phenotype and alters the encoded amino acid sequence of at least one gene allele of the human subject when compared to the amino acid sequence of the polypeptide encoded by a corresponding wild-type gene allele, wherein the wild-type polypeptide comprises an amino acid sequence selected from the group consisting of
30 SEQ ID NOS: 6, 8, 10, 12, 14, 16, 20, 22, 24, 42 and 43; (b) correlating the presence or absence of the mutation in the nucleic acid to a risk of developing hereditary

lymphedema, wherein the presence of the mutation in the nucleic acid correlates with an increased risk of developing hereditary lymphedema, and wherein the absence of the mutation in the nucleic acid correlates with no increased risk of developing hereditary lymphedema.

5 In another method of assaying for risk of developing hereditary lymphedema, the steps comprise (a) assaying nucleic acid of a human subject for a mutation that alters the encoded amino acid sequence of at least one intracellular protein or transcription factor allele of the human subject and alters transcription modulation activity of the transcription factor polypeptide encoded by the allele, when
10 compared to the transcription modulation activity of a transcription factor polypeptide encoded by a wild-type allele, wherein the wild-type intracellular protein or transcription factor polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 18, 26, 28, 30, 32, 34, 36, 38, 40, 44, 45, and 46; and (b) correlating the presence or absence of the mutation in the nucleic acid to a risk of
15 developing hereditary lymphedema, wherein the presence of the mutation in the nucleic acid correlates with an increased risk of developing hereditary lymphedema, and wherein the absence of the mutation in the nucleic acid correlates with no increased risk of developing hereditary lymphedema. As one example, the assaying identifies a mutation altering activity of a transactivating or DNA binding domain of a
20 Sox18 allele. In some embodiments of the method, the mutation reduces transcriptional activation of a SOX18-responsive gene compared to transcriptional activation of the gene in the absence of said mutation.

 In a related aspect, the invention provides a method of assaying for risk of developing hereditary lymphedema, comprising (a) assaying nucleic acid of a
25 human subject for a mutation that alters the encoded amino acid sequence of at least one LEC gene allele of the human subject and alters the binding affinity of the adhesion polypeptide encoded by the LEC gene allele, when compared to the binding affinity of an adhesion polypeptide encoded by a wild-type allele, wherein the wild-type adhesion polypeptide comprises an amino acid sequence of SEQ ID NO: 20; and
30 (b) correlating the presence or absence of the mutation in the nucleic acid to a risk of developing hereditary lymphedema, wherein the presence of the mutation in the

nucleic acid correlates with an increased risk of developing hereditary lymphedema, and wherein the absence of the mutation in the nucleic acid correlates with no increased risk of developing hereditary lymphedema.

In the methods of assaying for risk of developing hereditary lymphedema according to the invention, the assaying may identify the presence of the mutation, and the correlating step may identify the increased risk of the patient developing hereditary lymphedema.

A related method according to the invention is a method of screening a human subject for an increased risk of developing hereditary lymphedema comprising assaying nucleic acid of a human subject for a mutation that alters the encoded amino acid sequence of at least one polypeptide comprising an amino acid sequence found in any of Tables 1 and 2. In some embodiments of this method, the polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NOS: 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 43, 44, 45, and 46 in a manner that correlates with the risk of developing hereditary lymphedema, and it is expressly contemplated that the polypeptide may comprise the amino acid sequences set forth in SEQ ID NOS: 6, 8, 10, 12, 14, 16, 20, 22, 24, 42 and 43.

A related aspect of the invention is drawn to methods of assaying or screening for risk of developing hereditary lymphedema as described above, wherein the method comprises at least one procedure selected from the group consisting of: (a) determining a nucleotide sequence of at least one codon of at least one polynucleotide of the human subject; (b) performing a hybridization assay to determine whether nucleic acid from the human subject has a nucleotide sequence identical to or different from one or more reference sequences; (c) performing a polynucleotide migration assay to determine whether nucleic acid from the human subject has a nucleotide sequence identical to or different from one or more reference sequences; and (d) performing a restriction endonuclease digestion to determine whether nucleic acid from the human subject has a nucleotide sequence identical to or different from one or more reference sequences.

A related aspect of the invention provides methods of assaying or screening for risk of developing hereditary lymphedema as described above, wherein

the method comprises: performing a polymerase chain reaction (PCR) to amplify nucleic acid comprising the coding sequence of the LEC polynucleotide, and determining nucleotide sequence of the amplified nucleic acid.

Further provided by the invention is a method of screening for a hereditary lymphedema genotype in a human subject, comprising: (a) providing a biological sample comprising nucleic acid from said subject, and (b) analyzing the nucleic acid for the presence of a mutation altering the encoded amino acid sequence of the at least one allele of at least one gene in the human subject relative to a human gene encoding an amino acid sequence selected from the group consisting of SEQ ID NOS: 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 43, 44, 45, and 46, wherein the presence of a mutation altering the encoded amino acid sequence in the human subject in a manner that correlates with lymphedema in human subjects identifies a hereditary lymphedema genotype. In some embodiments of this method, the biological sample is a cell sample. In other embodiments of this method, the analyzing comprises sequencing a portion of the nucleic acid. In still further embodiments of this method, the human subject has a hereditary lymphedema genotype identified by the method of screening.

In is contemplated that the above methods further comprise assaying the nucleic acid for a second mutation that correlates with a hereditary lymphedema phenotype and alters the encoded amino acid sequence of a second gene allele of the human subject when compared to the amino acid sequence of the polypeptide encoded by a corresponding wild-type allele of said second gene, wherein the wild-type polypeptide encoded by said second gene is a polypeptide identified in any of Tables 3 and 4 of PCT/US03/06900.

Another aspect of the invention provides a method of inhibiting lymphangiogenesis comprising administering to a subject an inhibitor of a LEC transmembrane polypeptide, wherein the LEC transmembrane polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NOS: 6, 8, 10, 12, 14, 16, 20, 22, 24, 42 and 43, and wherein the inhibitor is selected from the group consisting of (a) a soluble extracellular domain fragment of the LEC transmembrane polypeptide; (b) an antibody that binds to the extracellular domain of the LEC

transmembrane polypeptide; (c) a polypeptide comprising an antigen binding domain of the antibody according to (b); and (d) an antisense nucleic acid complementary to the nucleic acid encoding the LEC transmembrane polypeptide or its complement. In some embodiments of the method, the inhibitor is a polypeptide comprising an
5 extracellular domain fragment of an LEC polypeptide. It is further contemplated that the sequence of said extracellular domain is selected from the group consisting of amino acids 1-1005 of SEQ ID NO: 43. In some embodiments of the method, the subject is a human containing a tumor.

In a related aspect, the invention provides a method for modulating
10 lymphangiogenesis in a mammalian subject comprising: administering to a mammalian subject in need of modulation of lymphangiogenesis an antisense molecule to a LEC polynucleotide, in an amount effective to inhibit transcription or translation of the polypeptide encoded by the LEC polynucleotide, wherein the LEC polynucleotide comprises a nucleotide sequence selected from the group consisting of
15 SEQ ID NOS: 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 and 41. Antisense molecules of suitable sequence and length are known in the art, and include sequences perfectly complementary to a coding region of a coding strand of a nucleic acid, as well as lengths capable of relatively specific and stable hybridization to a target *in vivo* (e.g., lengths of each incremental integer in the range of 8-50, with a
20 preferred length of at least 17 nucleotides).

The methods of the invention also include methods for the treatment of disorders related to lymphatic endothelial cells as recited above which may involve the administration of such compounds to individuals exhibiting symptoms or tendencies related to such disorders.

25 In another aspect, the invention provides a method of treating hereditary lymphedema, comprising: (a) identifying a human subject with hereditary lymphedema and with a mutation that alters the encoded amino acid sequence of at least one polypeptide of the human subject, relative to the amino acid sequence of a polypeptide comprising an amino acid sequence selected from the group consisting of
30 SEQ ID NOS: 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 43, 44, 45, and 46; and (b) administering to the subject a lymphatic growth factor

selected from the group consisting of a VEGF-C polynucleotide, a VEGF-C polypeptide, a VEGF-D polynucleotide, and a VEGF-D polypeptide.

The invention also provides a method of treating hereditary lymphedema comprising: identifying a human subject with lymphedema and with a mutation in at least one allele of a gene encoding a LEC protein identified in any of Tables 1 and 2, wherein the mutation correlates with lymphedema in human subjects, and with the proviso that the LEC-specific protein is not VEGFR-3; and administering to the subject a composition comprising a lymphatic growth agent selected from the group consisting of VEGF-C polypeptides, VEGF-D polypeptides, VEGF-C polynucleotides, and VEGF-D polynucleotides. The invention also comprehends use of a lymphatic growth agent selected from the group consisting of VEGF-C polypeptides, VEGF-D polypeptides, VEGF-C polynucleotides, and VEGF-D polynucleotides in the manufacture of a medicament for the treatment of hereditary lymphedema resulting from a mutation in a LEC gene identified in any of Tables 1 and 2. It is contemplated that the method further comprises a mutation in at least one allele of a second gene encoding a LEC protein identified in any of Tables 3 and 4 of PCT/US03/06900, wherein the mutation in said second gene correlates with lymphedema in human subjects.

In addition, the invention encompasses methods for treating such diseases or disorders by administering compounds and other substances that modulate the overall activity of the target gene products. Compounds and other substances can effect such modulation either at the level of target gene expression or target protein activity. These treatment methods include the administration of a polypeptide or a polynucleotide according to the invention to an endothelial cell, e.g., a LEC and/or a BEC, or to an organism such as a human patient. An exemplary method according to this aspect of the invention is the administration of a therapeutic selected from the group consisting of an antisense polynucleotide capable of modulating the expression of at least one polynucleotide according to the invention, a polypeptide according to the invention, a polynucleotide according to the invention, an antibody or antibody fragment specifically recognizing a polypeptide according to the invention, a VEGF-

C polynucleotide, a VEGF-C polypeptide, a VEGF-D polynucleotide, a VEGF-D polypeptide and a soluble VEGFR-3 polypeptide.

In another aspect, the invention provides a method of screening for an endothelial cell disorder or predisposition to the disorder, comprising obtaining a biological sample containing endothelial cell mRNA from a human subject; and measuring expression of a BEC or LEC gene from the amount of mRNA in the sample transcribed from the gene, wherein the BEC or LEC gene encodes a polypeptide identified in any of Tables 1 and 2. In a related aspect, the method further comprises a second BEC or LEC gene that encodes a polypeptide set forth in any of Tables 3 and 4 of PCT/US03/06900.

The invention also relates to a method of inhibiting the growth of a lymphatic endothelial cell, the method comprising contacting the cell with a composition comprising at least one antibody conjugated to an agent capable of inhibiting the growth, wherein the agent is selected from the group consisting of a cytotoxic agent and a cytostatic agent, and wherein the antibody specifically binds to a polypeptide encoded by a polynucleotide comprising a sequence selected from the group consisting of SEQ ID NOS: 5, 7, 9, 11, 13, 15, 19, 21 and 23. In specific embodiments of this method, the polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NOS: 6, 8, 10, 12, 14, 16, 20, 22, 24, 42 and 43.

The invention further relates to methods of detecting a lymphatic endothelial cell, the method comprising contacting the cell with a composition comprising at least one antibody conjugated to a detectable agent, such as a fluorescent molecule or a radiolabeled molecule. In specific embodiments, the antibody specifically binds to a polypeptide encoded by a polynucleotide comprising a sequence selected from the group consisting of SEQ ID NOS: 5, 7, 9, 11, 13, 15, 19, 21 and 23. In further specific embodiments of this method, the polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NOS: 6, 8, 10, 12, 14, 16, 20, 22, 24, 42 and 43.

The invention still further relates to methods of isolating a lymphatic endothelial cell, comprising contacting the cell with a solid matrix comprising at least

one antibody capable of binding to a transmembrane protein in the cell membrane of the cell, and isolating cells specifically bound to the antibody matrix. In specific embodiments, the antibody specifically binds to a polypeptide encoded by a polynucleotide comprising a sequence selected from the group consisting of SEQ ID NOS: 5, 7, 9, 11, 13, 15, 19, 21 and 23. In further specific embodiments of this method, the polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NOS: 6, 8, 10, 12, 14, 16, 20, 22, 24, 42 and 43.

The invention also relates to the administration of an agonist or antagonist to a lymphatic endothelial cell surface receptor, comprising selecting an antibody, a peptide or a small molecular weight compound that is capable of specifically binding to a lymphatic endothelial cell-specific protein, wherein the antibody, peptide or small molecular weight compound is an agonist or antagonist for a growth factor receptor, a cytokine receptor, a chemokine receptor, or a hemopoietic receptor, and contacting the antibody, peptide or small molecular weight compound with the lymphatic endothelial cell in need of growth stimulation or inhibition. In specific embodiments, such lymphatic endothelial cells are associated with lymphedema, lymphangioma, lymphangiomyeloma, lymphangiomatosis, lymphangiectasis, lymphosarcoma, and lymphangiosclerosis.

The invention also relates to the administration of a cytotoxic or cytostatic drug to a cancerous lymphatic endothelial cell, comprising selecting an antibody, a peptide or a small molecular weight compound that is capable of specifically binding to a lymphatic endothelial cell-specific protein, wherein the antibody, peptide or small molecular weight compound is complexed to the cytotoxic or cytostatic drug. In specific embodiments, administration of such complexes is useful in the treatment of malignant tumor diseases prone to metastatic spread through the lymphatic system.

The invention also provides a method of monitoring the efficacy or toxicity of a drug on endothelial cells, comprising the steps of measuring expression of at least one BEC or LEC gene in endothelial cells of a mammalian subject before and after administering a drug to the subject, wherein the at least one LEC or BEC gene encodes a polypeptide found in any of Tables 1 and 2 herein, and wherein

changes in expression of the LEC or BEC gene correlates with efficacy or toxicity of the drug on endothelial cells. It is contemplated that the method further comprises a second BEC or LEC gene that encodes a polypeptide set forth in any of Tables 3 and 4 of PCT/US03/06900.

5 The invention further relates to a method of detecting a lymphatic endothelial cell, comprising contacting said cell with an antibody as described herein, wherein the antibody is detectably labeled.

 The invention still further relates to a method of inhibiting at least one biological activity of a lymphatic endothelial cell, comprising contacting the cell with
10 an agent capable of binding to at least one polypeptide encoded by any one of SEQ ID NOS: 6, 8, 10, 12, 14, 16, 20, 22, 24, 42 and 43, wherein the activity of the polypeptide is reduced relative to the activity of a polypeptide that is not contacted with the agent.

 The invention also relates to a method of inhibiting the growth of a
15 lymphatic endothelial cell, the method comprising contacting the cell with an antisense oligonucleotide capable of specifically binding to at least one polynucleotide selected from the group consisting of SEQ ID NOS: 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 and 41. In a specific embodiment, the antisense oligonucleotide consists essentially of about 12 to about 25 contiguous
20 nucleotides of any one of SEQ ID NOS: 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 and 41.

 It is further contemplated that the methods of the invention, such as identifying a LEC nucleic acid; modulating the growth of lymphatic endothelial cells in a human subject; identifying compounds that modulate growth of endothelial cells;
25 identifying a human subject with a disorder characterized by hyperproliferation of LECs; assaying for risk of developing hereditary lymphedema; screening for an increased risk of developing hereditary lymphedema; screening for a hereditary lymphedema genotype in a human subject; and inhibiting lymphangiogenesis, may be carried out using compositions comprising polypeptide fragments as described above.
30 In some embodiments, the polypeptide is at least 95% identical to a polypeptide set forth herein, comprising a particular biological activity associated with a protein

domain as set forth herein. The polypeptide may have transcription factor activity or cytoskeletal regulation activity. Additional exemplary biological activities include the ubiquitin ligase activity of a Hect domain, guanine-nucleotide-dissociation-stimulating activity of at least one RCC1-like domain, cation transport activity of a cation transport domain, GTP-binding activity of at least one GTP-binding domain, SPRY domain activity of at least one SPRY domain, and Kelch domain activity of at least one Kelch domain. Variations on the methods above may also be carried out using the above-described polypeptide fragment compositions.

Additional features and variations of the invention will be apparent to those skilled in the art from the entirety of this application, and all such features are intended as aspects of the invention.

DETAILED DESCRIPTION OF THE INVENTION

A major role of the lymphatic vasculature is to remove an excess of the protein-rich interstitial fluid that continuously escapes from the blood capillaries, and to return it to the blood circulation (Witte, M.H., et al., *Microsc. Res. Tech.* 55:122-145. 2001; Karpanen, T., et al., *J. Exp. Med.* 194:F37-F42. 2001; Karkkainen, M.J., et al., *Trends Mol. Med.* 7:18-22, 2001). In addition, the lymphatic system provides constant immune surveillance by filtering lymph and its antigens through the chain of lymph nodes, and also serves as one of the major routes for absorption of lipids from the gut. It has been known for a long time that in many types of cancer the lymphatic vessels provide a major pathway for tumor metastasis, and regional lymph node dissemination correlates with the progression of the disease. Hereditary lymphedema, post-surgical secondary lymphedema and lymphatic obstruction in filariasis, are all characterized by disabling and disfiguring swelling of the affected areas, linked to the insufficiency or obstruction of the lymphatics. Witte, M.J., et al., *Microsc. Res. Tech* 55:122-145 (2001).

In spite of the importance of lymphatic vessels in medicine, the cell biology of this part of the vascular system has received little attention until recently. Studies during the past four years have uncovered the existence of the lymphatic

specific vascular endothelial growth factors VEGF-C and VEGF-D, which serve as ligands for the receptor tyrosine kinase VEGFR-3, and demonstrated their importance for the normal development of the lymphatic vessels (*See, Jeltsch et al., Science* 276:1423-1425 (1997); Veikkola, T., *et al., EMBO J.* 20:1223-1231 (2001); Mäkinen, T., *et al., Nat. Med.* 7:199-205 (2001)). These molecules also appear to be involved in the development of lymphedema and lymphatic metastasis (Karpanen *et al., J. Exp. Med.* 194:F37-F42 (2001); Karkkainen *et al., Trends Mol. Med.* 7:18-22, 2001).

The growth factor Vascular Endothelial Growth Factor C (VEGF-C), as well as native human, non-human mammalian, and avian polynucleotide sequences encoding VEGF-C, and VEGF-C variants and analogs, have been described in detail in International Patent Application Number PCT/US98/01973, filed February 2, 1998 and published on August 6, 1998 as International Publication Number WO 98/33917; in Joukov *et al., J. Biol. Chem.*, 273(12): 6599-6602 (1998); and in Joukov *et al., EMBO J.*, 16(13): 3898-3911 (1997), all of which are incorporated herein by reference in their entirety. As explained therein in detail, human VEGF-C (SEQ ID NOS: 1 and 2) is initially produced in human cells as a prepro-VEGF-C polypeptide of 419 amino acids. A cDNA encoding human VEGF-C (SEQ ID NO: 1) has been deposited with the American Type Culture Collection (ATCC), 10801 University Blvd., Manassas, VA 20110-2209 (USA), pursuant to the provisions of the Budapest Treaty (Deposit date of 24 July 1995 and ATCC Accession Number 97231). VEGF-C sequences from other species also have been reported. See Genbank Accession Nos. MMU73620 (*Mus musculus*); and CCY15837 (*Coturnix coturnix*) for example, incorporated herein by reference.

The prepro-VEGF-C polypeptide is processed in multiple stages to produce a mature and most active VEGF-C polypeptide of about 21-23 kD, as assessed by SDS-PAGE under reducing conditions (SEQ ID NO: 2). Such processing includes cleavage of a signal peptide (residues 1-31); cleavage of a carboxyl-terminal peptide (corresponding approximately to amino acids 228-419 and having a pattern of spaced Cysteine residues reminiscent of a Balbiani ring 3 protein (BR3P) sequence [Dignam *et al., Gene*, 88:133-40 (1990); Paulsson *et al., J. Mol. Biol.*, 211:331-49 (1990)]) to produce a partially-processed form of about 29 kD; and cleavage

(apparently extracellularly) of an amino-terminal peptide (corresponding approximately to amino acids 32-103) to produce a fully-processed mature form of about 21-23 kD. Experimental evidence demonstrates that partially-processed forms of VEGF-C (e.g., the 29 kD form) are able to bind VEGFR-3 (Flt4 receptor), whereas high affinity binding to VEGFR-2 occurs only with the fully processed forms of VEGF-C. It appears that VEGF-C polypeptides naturally associate as non-disulfide linked dimers.

It has been demonstrated that amino acids 103-227 of VEGF-C are not all critical for maintaining VEGF-C functions. A polypeptide consisting of amino acids 113-213 (and lacking residues 103-112 and 214-227) retains the ability to bind and stimulate VEGF-C receptors, and it is expected that a polypeptide spanning from about residue 131 to about residue 211 will retain VEGF-C biological activity. The Cysteine residue at position 156 has been shown to be important for VEGFR-2 binding ability. However, VEGF-C ΔC_{156} polypeptides (i.e., analogs that lack this Cysteine due to deletion or substitution) remain potent activators of VEGFR-3. The Cysteine at position 165 of VEGF-C polypeptide is essential for binding either receptor, whereas analogs lacking the Cysteine at positions 83 or 137 compete with native VEGF-C for binding with both receptors and stimulate both receptors.

VEGF-D is structurally and functionally most closely related to VEGF-C (*see* U.S. Patent 6,235,713 and International Patent Publ. No. WO 98/07832, incorporated herein by reference). See SEQ ID NO: 3 for the polynucleotide sequence of VEGF-D; the encoded amino acid sequence is set forth in SEQ ID NO: 4. Like VEGF-C, VEGF-D is initially expressed as a prepro-peptide that undergoes N-terminal and C-terminal proteolytic processing, and forms non-covalently linked dimers. VEGF-D stimulates mitogenic responses in endothelial cells *in vitro*. During embryogenesis, VEGF-D is expressed in a complex temporal and spatial pattern, and its expression persists in the heart, lung, and skeletal muscles in adults. Isolation of a biologically active fragment of VEGF-D designated VEGF-D $\Delta N\Delta C$, is described in International Patent Publication No. WO 98/07832, incorporated herein by reference. VEGF-D $\Delta N\Delta C$ consists of amino acid residues 93 to 201 of VEGF-D (SEQ ID NO: 4) optionally linked to the affinity tag peptide FLAG[®], or other sequences.

The prepro-VEGF-D polypeptide has a putative signal peptide of 21 amino acids and is apparently proteolytically processed in a manner analogous to the processing of prepro-VEGF-C. A “recombinantly matured” VEGF-D lacking residues 1-92 and 202-354 retains the ability to activate receptors VEGFR-2 and VEGFR-3, and appears to associate as non-covalently linked dimers. Thus, preferred VEGF-D polynucleotides include those polynucleotides that comprise a nucleotide sequence encoding amino acids 93-201. The guidance provided above for introducing function-preserving modifications into VEGF-C polypeptides is also suitable for introducing function-preserving modifications into VEGF-D polypeptides. As another aspect of the invention, practice of the invention methods is contemplated wherein VEGF-D polypeptides are employed in lieu of VEGF-C polypeptides.

When compared with the blood vascular endothelium, the lymphatic endothelium exhibits specific morphological and molecular characteristics. For example, the lymphatic capillaries are larger than blood capillaries, they have an irregular or collapsed lumen with no red blood cells, a discontinuous basal lamina, overlapping intercellular junctional complexes and anchoring filaments that connect the lymphatic endothelial cells to the extracellular matrix (Witte, M.H., et al., *Microsc. Res. Tech.* 55:122-145 (2001)). Unlike the blood capillaries, the lymphatic capillaries lack pericyte coverage. At the molecular level several lymphatic specific markers have been identified, including VEGFR-3, the Prox-1 transcription factor, the hyaluronan receptor LYVE-1, the membrane mucoprotein podoplanin, the beta-chemokine receptor D6, the cytoskeletal proteins desmoplakin I and II and the macrophage mannose receptor I (Wigle, J.T. & Oliver, G., *Cell* 98:769-778 (1999); Banerji, S., et al., *J. Cell Biol.* 144:789-801 (1999); Breiteneder-Geleff, S., et al., *Am. J. Pathol.* 154:385-394 (1999); Nibbs, R.J., et al., *Am. J. Pathol.* 158:867-877 (2001); Ebata, N., et al., *Microvasc. Res.* 61:40-48. (2001); Irjala, H., et al., *J. Exp. Med.* 194:1033-1041 (2001)). The present invention relates to the genetic identity of lymphatic capillary endothelial cells versus blood vascular endothelial cells using a gene profiling approach.

“Stringent hybridization conditions” or “stringent conditions” refer to conditions under which a nucleic acid such as an oligonucleotide will specifically

hybridize to its target sequence. Stringent conditions are sequence-dependent and will be different in different circumstances. Longer nucleic acids hybridize specifically at higher temperatures than shorter sequences. Generally, stringent conditions are selected to be about 5°C lower than the thermal melting point (T_m) for the specific
5 sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength, pH and nucleic acid concentration conditions) at which 50% of the nucleic acids complementary to the target sequence hybridize to the target sequence at equilibrium. The term “complementary” refers to standard Watson-Crick base pairing between nucleotides of two nucleic acid molecules. Typically, stringent
10 conditions will be those in which the salt concentration is less than about 1.0 M sodium ion, typically about 0.01 to 1.0 M sodium ion (or other salts) at pH 7.0 to 8.3 and at a temperature that is at least about 30°C for short probes, primers or oligonucleotides (*e.g.*, 10 to 50 nucleotides) and at least about 60°C for longer probes, primers or oligonucleotides. Stringent conditions also can be achieved with the
15 addition of destabilizing agents, such as formamide, as is known in the art Exemplary stringent hybridization conditions are hybridization at 42°C for 20 hours in a solution containing 50% formamide, 5xSSPE, 5x Denhardt's solution, 0.1% SDS and 0.1 mg/ml denatured salmon sperm DNA, with a wash in 1xSSC, 0.1% SDS for 30 minutes at 65°C.

20 Distinct gene expression profiles for blood vascular and lymphatic endothelial cells have been discovered (See PCT/US03/06900, incorporated herein by reference). These results provide new insights into the phenotypic diversity of endothelial cells and reveal new potential lymphatic endothelial cell molecules, some of which could provide important targets for the therapy of diseases characterized by
25 abnormal angiogenesis or lymphangiogenesis.

Differences in the expression of genes encoding proteins involved in inflammatory processes were found, as well as in those mediating cell-cell and cell-matrix interactions. Furthermore, several previously unknown genes were identified in the context of endothelial cell biology, which were differentially expressed in the
30 two cell lineages. Several of the genes were originally cloned from neural tissues, including genes involved in the uptake of synaptic macromolecules and in synapse

formation and remodeling (neuronal pentraxins I and II (Kirkpatrick, L.L., et al., *J. Biol. Chem.* 275:17786-17792. 2000), in the trafficking of synaptic vesicles (NAP-22 (Yamamoto, Y., et al., *Neurosci. Lett.* 224:127-130. 1997), piccolo (Fenster, S.D., et al., *Neuron* 25:203-214 (2000)) and in the axon growth and guidance (Nr-CAM (Grumet, M., *Cell Tissue Res.* 290:423-428 (1997), reelin (Rice, D.S. & Curran, T., *Annu. Rev. Neurosci.* 24:1005-1039 (2001)).

In addition, the LECs especially expressed a number of uncharacterized genes, which were originally cloned and highly expressed in nervous tissues (KIAA genes (Kikuno, R., et al., *Nucleic Acids Res.* 30:166-168. 2002)). The gene expression profiling data disclosed herein therefore support the view that the same molecular mechanisms that are involved in governing neural cell positioning, in guiding axonal growth cones to their specific targets and in synaptogenesis may also be commonly used in the development of the vascular system and in the establishment of BEC and LEC identity. Some other signaling molecules first described in the developing nervous system have already been implicated in the development of the vasculature and vice versa (Shima and Mailhos, *Curr. Opin. Genet. Dev.* 10:536-542 (2000); Oosthuyse, et al., *Nat. Genet.* 28:131-138 (2001); Sondell, et al., *Eur. J. Neurosci.* 12:4243-4254 (2000)).

PCT/US03/06900 identified expression of several genes in the LECs, previously shown to be expressed in smooth muscle cells (SMCs) and pericytes, such as matrix Gla, a mineral binding extracellular matrix protein involved in the inhibition of vascular and tissue calcification (Luo, G., et al., *Nature* 386:78-81 (1997)), monoamine oxidase A, the main degradative enzyme of monoamine hormones and neurotransmitters (Rodriguez, M.J., et al., *Cell Tissue Res.* 304:215-220 (2001)), integrin $\alpha 9$ (Palmer, E.L., et al., *J. Cell Biol.* 123:1289-1297 (1993)) and apolipoprotein D (Hu, C.Y., et al., *J. Neurocytol.* 30:209-218 (2001)). Some similarity of gene expression patterns between LECs and SMCs could be related to the lack of SMC around lymphatic capillaries. Instead, LECs may carry out some SMC functions by themselves. For example, lymph flow is maintained due to the intrinsic contractility of the LECs (Witte, M.H., et al., *Microsc. Res. Tech.* 55:122-145 (2001)), reminiscent of the ability of vascular SMCs to contract.

Molecular discrimination of the lymphatic and blood vessels is essential in studies of diseases involving the blood and/or lymphatic vessels and in the targeted treatment of such diseases. To date, several lymphatic endothelial specific markers have been identified, but some of them are expressed only in a subset of the lymphatic vessels, while others also occur in some blood vessel endothelia or in other cell types and their expression patterns may change in pathological conditions (for example, VEGFR-3 (Valtola, R., et al., *Am. J. Pathol.* 154:1381-1390. 1999)). Identification of new vascular markers according to the invention should provide a more reliable analysis of the blood and lymphatic vessels in pathological situations and eventually better diagnosis and treatment. Furthermore, inhibition of the function of certain molecules involved in the regulation of angiogenesis and/or lymphangiogenesis is known to prevent tumor growth and metastasis, and stimulation of the growth of blood or lymphatic vessels has been shown to be beneficial in several pathological conditions. Thus the LEC specific molecular regulators identified according to the invention may provide new targets for the treatment of diseases characterized by abnormal angiogenesis and lymphangiogenesis.

Several of the new LEC genes encode transmembrane proteins which are expected to be specific molecular markers for lymphatic endothelial cells (Table 1). These genes and encoded proteins are useful for targeted treatment of diseases that involve lymphatic vessels. They may also be useful for preparing antibodies, as antibodies against LEC-specific proteins can be used to discriminate between blood and lymphatic vessels in pathological and physiological situations. Antibodies may also be useful for the isolation of lymphatic endothelial cells. These proteins may also play a role in the regulation of lymphangiogenesis, and can provide new candidate genes for diseases that involve lymphatic vessels, such as lymphedema.

The lymphatic endothelial cell specific surface molecules can be used for molecular drug targeting with antibodies, peptides and small-molecular weight compounds, which can act as agonists or antagonists for growth factor receptor, cytokine and chemokine receptor, and hemopoietin receptor signaling, cell adhesion and cell interaction with extracellular matrix or with other cell surface molecules. Such molecules can also be used for targeting of cytotoxic or cytostatic drugs into the

lymphatic endothelial cells and for the attachment of electron-dense, radio-opaque or radioactive markers for imaging of disease processes associated with the lymphatic vessels. Such diseases include lymphedema, lymphangioma, lymphangiomyoma, lymphangiomatosis, lymphangiectasis, lymphosarcoma and lymphangiosclerosis.

5 The lymphatic endothelial cell surface molecules may be used for targeting of gene therapy for example by antibody-coated liposomes (containing proteins or genes as cargo) or by viral transducing vectors such as adenoviruses, adeno-associated viruses or lentiviruses having modified capsid/envelope proteins. The manipulation of lymphatic endothelial cell specific molecules may be applicable
10 to treatment of disease processes associated with tissue edema by increasing fluid transport across the lymphatic vessel wall for example by modulating endothelial cell-cell or cell-matrix interactions or via stimulating transendothelial transport. Targeting of the lymphatic endothelial cells for example with cytotoxic or cytostatic compounds is contemplated to be valuable in malignant tumor diseases prone to metastatic spread
15 via the lymphatic system.

 The lymphatic endothelial cell molecules may allow the improved *in vitro* growth of lymphatic endothelial cells as well as *in vitro* tissue engineering of lymphatic vessels for use in diseases where the lymphatics have been damaged, such as after surgery and in various forms of lymphedema. Ligands of the cell surface
20 proteins may further be applied to coat various polymeric matrices for the adhesion of cells in, e.g., bioimplants.

 The lymphatic endothelial-cell-specific molecules such as surface molecules can provide important tools for the modulation of inflammatory, autoimmune and infectious processes involving leukocyte migration and immune
25 recognition as well as the stimulation of secondary immune responses. Such processes include the migration of antigen presenting cells into the lymphatic system including lymph nodes as well as transendothelial cell trafficking of lymphocytes and other leukocyte subclasses and the homing, survival and function of the various classes of leukocytes.

30 These molecules may allow one to modulate the metabolism of fatty acids including fatty acid/chylomicron absorption from the gut and regulation of fat

accumulation in adipose tissue in various organs such as in the subcutaneous tissue and in the arterial wall.

Lymphatic endothelial-cell-specific molecules may further allow one to modulate the metabolism of fatty acids including fatty acid/chylomicron absorption from the gut and regulation of fat accumulation in adipose tissue in various organs such as in the skin subcutaneous tissue and in the arterial wall.

The lymphatic-cell-specific transmembrane proteins are expected to function in cell adhesion (e.g., adhesion between lymphatic endothelial cell-lymphatic endothelial cell, lymphatic endothelial cell-smooth muscle cell, lymphatic endothelial cell-immune system cell such as lymphocyte or dendritic cell), cell-extracellular matrix contacts, or as receptors such as growth factor, cytokine, chemokine or microbial receptors or ion channels. The transmembrane (TM) proteins connect to intracellular molecules that can induce cell growth, cell migration, cell apoptosis, cell differentiation or cell adhesion or other cellular functions specific for endothelial cells such as expression of adhesion receptors for leukocytes, release of nitric oxide, antithrombotic proteins, uptake of fluid and proteins from surrounding tissues and fat from gut or adipose tissues. TM proteins with short intracellular domains can function as auxiliary receptors in complex with other TM proteins.

The transmembrane proteins and their intracellular binding partner molecules can be used as molecular markers for lymphatic endothelial cells in normal and disease conditions, and to discriminate between blood and lymphatic vessels in pathological and physiological situations.

Antibodies against lymphatic specific transmembrane proteins, as well as peptides and small molecular compounds binding to extracellular domains of lymphatic-specific TM proteins can be used for the attachment of electron-dense, radio-opaque or radioactive markers for imaging of disease processes associated with the lymphatic vessels. Such diseases include lymphedema, lymphangioma, lymphangiomyoma, lymphangiomatosis, lymphangiectasis, lymphosarcoma and lymphangiosclerosis. Similarly, the lymphatic vessels can be visualized, e.g., during therapy of patients suffering from insufficient lymphatic growth, such as in lymphedema, or alternatively during treatment aiming to prevent lymphatic growth,

e.g., in tumors, thereby facilitating the monitoring of the therapeutic method of the invention.

Antibodies against LEC-specific TM proteins are also expected to be useful for the isolation of lymphatic endothelial cells.

5 Antibodies against lymphatic-specific transmembrane proteins, or peptides or small-molecule compounds binding to the extracellular domain of lymphatic-specific TM proteins are expected to be useful in targeting drug delivery to lymphatic endothelial cells, e.g., by coupling an antibody, peptide or small-molecule compound to a cytotoxic or cytostatic compound. Such coupled compounds are
10 useful as therapeutics in the treatment of malignant tumor diseases prone to metastatic spread via the lymphatic system, as well as in ameliorating a symptom associated with any such disease. The antibodies, peptides or small-molecule compounds can also be coupled to stimulatory lymphatic endothelial molecules such as growth factors, cytokines and chemokines to promote stimulation.

15 Additionally, antibodies against lymphatic-specific TM proteins or peptides, or small-molecule compounds binding to the extracellular domain of lymphatic-specific TM proteins, may be used for targeting of gene therapy, for example, by antibody-coated liposomes (containing proteins, genes or other molecules as cargo) or by viral transducing vectors such as adenoviruses, adeno-
20 associated viruses, lentiviruses, or the like, having modified capsid/envelope proteins. The manipulation of lymphatic endothelial-cell-specific molecules are expected to be applicable to the treatment of disease processes associated with tissue edema due to the relative absence, or relative dysfunction, of lymphatic vessels, which can result from an infection, surgery, radiotherapy or a genetic defect by increasing fluid
25 transport across the lymphatic vessel wall, for example by modulating endothelial cell-cell or cell-matrix interactions or by stimulating transendothelial transport.

 The lymphatic endothelial cell molecules are expected to improve the *in vitro* growth of lymphatic endothelial cells, as well as the *in vitro* tissue engineering of lymphatic vessels for use in treating disorders or diseases where the lymphatics
30 have been damaged, such as after surgery, in various forms of lymphedema, and in other applications as described herein. Ligands of the cell-surface proteins may

further be applied as a coating to various polymeric matrices for the adhesion of cells in, e.g., bioimplants.

Inflammatory, autoimmune and infectious processes involving leukocyte migration and immune recognition, such as migration of antigen-presenting
5 cells into the lymphatic system, including lymph nodes, as well as transendothelial cell trafficking of lymphocytes and other leukocyte subclasses and the homing, survival and function of the various classes of leukocytes can be modulated by targeting endothelial-cell-specific TM proteins, which mediate these cell adhesion processes.

10 Upregulation of lymphatic-specific genes in, e.g., cancer are expected to be useful as diagnostic markers, and monitoring such upregulated expression with an antibody against a lymphatic endothelial-cell-specific protein, e.g., by immunostaining of tissue(s) or by using a probe hybridizable to a lymphatic endothelial-cell-specific mRNA, e.g., under stringent hybridization conditions as
15 described herein, is contemplated.

Lymphatic endothelial-cell-specific transcription factors are expected to be useful for the differentiation of lymphatic endothelial cells from embryonic stem cells, endothelial precursor cells, or blood vascular endothelial cells.

The lymphatic endothelial transcription factors are expected to improve
20 the *in vitro* growth of lymphatic endothelial cells, as well as to facilitate *in vitro* tissue engineering of lymphatic vessels for use in treating disorders or diseases where the lymphatics have been damaged, such as after surgery, in various forms of lymphedema, and in other applications disclosed herein.

Intracellular signaling proteins participating in signaling pathways
25 regulating lymphatic endothelial cell proliferation, differentiation, apoptosis, migration or adhesion are expected to be useful targets for small-molecule compounds inhibiting these signaling events, and cellular functions dependent on such signaling. Signaling proteins are also expected to participate in VEGFR-3 signaling pathways, and will be useful in modulating cellular activities controlled, at least in part, by
30 VEGFR-3 signaling, such as lymphangiogenesis.

The lymphatic endothelial cell molecules are expected to improve the *in vitro* growth of lymphatic endothelial cells as well as *in vitro* tissue engineering of lymphatic vessels for use in treating diseases or disorders where the lymphatics have been damaged, such as after surgery, in various forms of lymphedema, and in other applications as described herein.

Lymphatic-specific transcription factors are also expected to be useful in modulating gene expression in endothelial cells to induce the expression of other lymphatic-specific genes in, for example, blood vascular endothelial cells or endothelial precursor cells.

Lymphatic-specific gene transcripts are expected to provide useful targets for RNA interference (RNAi)-induced inhibition of expression. RNAi technology is expected to be useful in the methods according to the invention, such as therapeutic methods effective in treating hyper- and hypo-proliferative endothelial-cell-associated diseases and disorders, as well as methods of ameliorating a symptom of any such disease or disorder. RNAi methodologies are known in the art and known RNAi technologies are contemplated as useful in various aspects of the invention. See Fire et al., *Nature* 391:806-811. (1998) and Sharp, P., *Genes and Dev.* 13:139-141. (1999), each of which is incorporated herein by reference. It is preferred that RNAi compounds be double-stranded RNA molecules corresponding to part or all of a coding region of a desired target for expression.

As noted, several of the new LEC genes encode transcription factors, may control cellular fate and have an important role in the differentiation of lymphatic endothelial cells. Transcription factors disclosed herein (Table 2) may control transcription of genes involved for example in the proliferation of lymphatic endothelial cells, and may be important molecular regulators of lymphatic growth. Lymphatic endothelial cell specific transcription factors can be used for the differentiation of lymphatic endothelial cells from embryonic stem cells, endothelial precursor cells or from blood vascular endothelial cells.

The lymphatic endothelial transcription factors may allow the improved *in vitro* growth of lymphatic endothelial cells as well as *in vitro* tissue

engineering of lymphatic vessels for use in diseases where the lymphatics have been damaged, such as after surgery and in various forms of lymphedema.

Polynucleotides of the Invention

In general, the isolated polynucleotides of the invention include the LEC and BEC polynucleotides exhibiting differential expression and identified in Tables 1 and 2, including sequence identifiers of these polynucleotides and their known database accession numbers, where applicable. The polynucleotide sequences may include a coding region and may include non-coding flanking sequences, which are readily identifiable by one of skill in the art. The invention contemplates polynucleotides comprising part, or all, of a coding region, with or without flanking regions, e.g., poly A sequences, 5' non-coding sequences, and the like. The polynucleotides of the present invention also include, but are not limited to, a polynucleotide that hybridizes to the complement of the nucleotide sequence of any one of SEQ ID NOS: 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 and 41 under highly stringent hybridization conditions; a polynucleotide that hybridizes to the complement of the nucleotide sequence of any one of SEQ ID NOS: 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 and 41 under moderately stringent hybridization conditions; a polynucleotide which is an allelic variant of any polynucleotide recited above; a polynucleotide which encodes a species homologue of any of the proteins recited above; or a polynucleotide that encodes a polypeptide comprising a specific domain or truncation of the polypeptide of any one of SEQ ID NOS: 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 43, 44, 45, and 46. Such polynucleotides hybridize under the above conditions to the complement of any one of SEQ ID NOS: 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 and 41 or to a fragment of any one of SEQ ID NOS: 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 and 41 wherein the fragment is greater than at least about 10 bp, and, in alternate embodiments, is about 20 to about 50 bp, or is greater than about 100 bp, 200 bp, 300 bp, 400 bp, 500 bp, 600 bp, 700 bp, or 800 bp, where appropriate. References to "bp" refer to the length of double-, or

single-, stranded nucleic acids in nucleotides or base pairs, as would be apparent from the context.

The polynucleotides of the invention also provide polynucleotides that are variants of the polynucleotides recited above. Typically, such a variant sequence varies from one of those listed herein by no more than about 20%, *i.e.*, the number of individual nucleotide substitutions, additions, and/or deletions in a similar sequence, as compared to the corresponding reference sequence, divided by the total number of nucleotides in the variant sequence is about 0.2 or less. Such a sequence is said to have 80% sequence identity to the listed sequence. Such a variant sequence can be routinely identified by applying the foregoing algorithm.

In one embodiment, a variant polynucleotide sequence of the invention varies from a listed sequence by no more than 10%, *i.e.*, the number of individual nucleotide substitutions, additions, and/or deletions in a variant sequence, as compared to the corresponding reference sequence, divided by the total number of nucleotides in the variant sequence is about 0.1 or less. Such a sequence is said to have 90% sequence identity to the listed sequence. Such a variant sequence can be routinely identified by applying the foregoing algorithm.

In an alternate embodiment a variant sequence of the invention varies from a listed sequence by no more than 5%, *i.e.*, the number of individual nucleotide substitutions, additions, and/or deletions in a variant sequence, as compared to the corresponding reference sequence, divided by the total number of nucleotides in the variant sequence is about 0.05 or less. Such a sequence is said to have 95% sequence identity to the listed sequence. Such a variant sequence can be routinely identified by applying the foregoing algorithm.

In yet another alternate embodiment, a variant sequence of the invention varies from a listed sequence by no more than 2%, *i.e.*, the number of individual nucleotide substitutions, additions, and/or deletions in a variant sequence, as compared to the corresponding reference sequence, divided by the total number of nucleotides in the variant sequence is about 0.02 or less. Such a sequence is said to have 98% sequence identity to the listed sequence. Such a variant sequence can be routinely identified.

A polynucleotide according to the invention can be joined to any of a variety of other nucleotide sequences by well-established recombinant DNA techniques (see Sambrook, J. et al. (2d Ed.; 1989) *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY). Useful nucleotide sequences for joining to polypeptides include an assortment of vectors, *e.g.*, plasmids, cosmids, lambda phage derivatives, phagemids, and the like, that are well known in the art. Accordingly, the invention also provides a vector including a polynucleotide of the invention and a host cell containing the polynucleotide. In general, the vector contains an origin of replication functional in at least one organism, convenient restriction endonuclease sites, and a selectable marker for the host cell. Vectors according to the invention include expression vectors, replication vectors, probe generation vectors, sequencing vectors, and retroviral vectors. A host cell according to the invention can be a prokaryotic or eukaryotic cell and can be a unicellular organism or part of a multicellular organism. Large numbers of suitable vectors and promoters are known to those of skill in the art and are commercially available for generating the recombinant constructs of the present invention.

The sequences falling within the scope of the present invention are not limited to the specific sequences herein described, but also include allelic variations thereof. Allelic variations can be routinely determined by comparing the polynucleotide sequence provided in any one of SEQ ID NOS: 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 and 41, a representative intermediate fragment thereof, or a nucleotide sequence at least 99.9% identical to any one of SEQ ID NOS: 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 and 41 with a sequence from another isolate of the same species. Furthermore, to accommodate codon variability, the invention includes nucleic acid molecules coding for the same amino acid sequences as do the specific open reading frames (ORFs) disclosed herein. In other words, in the coding region of an ORF, substitution of one codon for another which encodes the same amino acid is expressly contemplated.

Unless provided for otherwise here, all terms are defined as is known in the art, for example as employed in U.S. Patent No. 6,350,447, incorporated herein by reference.

Also contemplated are antisense polynucleotides based on the sequence of any of the LEC or BEC polynucleotides set forth found in any of Tables 1 and 2 herein or Tables 3 and 4 of PCT/US03/06900. Such antisense polynucleotides are substantially complementary (e.g., at least 90% complementarity), and preferably perfectly complementary, to any one of the polynucleotides of the invention, or fragments thereof, set out in any of Tables 1 and 2 herein or Tables 3 and 4 of PCT/US03/06900 (incorporated herein by reference) that are differentially expressed in LECs and BECs. These polynucleotide sequences include any of SEQ ID NOS: 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 and 41, or a fragment thereof comprising at least 10 contiguous nucleotides. Antisense nucleic acid comprises a nucleotide sequence that is complementary to a "sense" nucleic acid encoding a protein (e.g., complementary to the coding strand of a double-stranded cDNA molecule or complementary to an mRNA sequence). Methods for designing and optimizing antisense nucleotides are described in Lima *et al.*, (*J Biol Chem*, ;272:626-38. 1997) and Kurreck *et al.*, (*Nucleic Acids Res.*, ;30:1911-8. 2002). In one aspect, antisense nucleic acid molecules are provided that comprise a sequence complementary to at least about 10, 25, 50, 100, 250 or 500 nucleotides or an entire coding strand. An antisense nucleic acid of the invention can be constructed using chemical synthesis or enzymatic ligation reactions using procedures known in the art.

In one embodiment, an antisense nucleic acid molecule is antisense to a "coding region" of the coding strand of a nucleotide sequence encoding a polypeptide of the invention. The term "coding region" refers to the region of the nucleotide sequence comprising codons which are translated into amino acid residues. In another embodiment, the antisense nucleic acid molecule is antisense to a "conceding region" of the coding strand of a nucleotide sequence encoding the polynucleotide. The term "conceding region" refers to 5' and 3' sequences which flank the coding region that are not translated into amino acids (i.e., also referred to as 5' and 3' untranslated regions).

Antisense nucleic acids of the invention can be designed according to the rules of Watson and Crick or Hoogsteen base pairing. The antisense nucleic acid molecule can be complementary to the entire coding region of the mRNA of the

polynucleotide of the invention, but more preferably is an oligonucleotide that is antisense to only a portion of the coding or noncoding region of the mRNA. An antisense oligonucleotide can be, for example, about 5, 10, 15, 20, 25, 30, 35, 40, 45, or 50 nucleotides in length. An antisense nucleic acid of the invention can be
5 constructed using chemical synthesis or enzymatic ligation reactions using procedures known in the art. For example, an antisense nucleic acid (e.g., an antisense oligonucleotide) can be chemically synthesized using naturally occurring nucleotides or variously modified nucleotides designed to increase the biological stability of the molecules or to increase the physical stability of the duplex formed between the
10 antisense and sense nucleic acids (e.g., phosphorothioate derivatives and acridine substituted nucleotides can be used).

The antisense nucleic acid molecules of the invention are typically administered to a subject or generated *in situ* such that they hybridize or bind to cellular mRNA and/or genomic DNA encoding the complementary polynucleotide,
15 thereby inhibiting expression of the protein (e.g., by inhibiting transcription and/or translation). The hybridization can reflect conventional nucleotide complementarity to form a stable duplex, or, for example, in the case of an antisense nucleic acid molecule that binds to DNA duplexes, through specific interactions in the major groove of the double helix.

20 An example of a route of administration of antisense nucleic acid molecules of the invention includes direct injection at a tissue site. Alternatively, antisense nucleic acid molecules can be modified to target selected cells and then administered systemically. For example, for systemic administration, antisense molecules can be modified such that they specifically bind to receptors or antigens
25 expressed on a selected cell surface (e.g., by linking the antisense nucleic acid molecules to peptides or antibodies that bind to cell surface receptors or antigens). Additional routes of antisense therapy may be used in the invention, e.g., topical administration, transdermal administration [reviewed by Brand in *Curr. Opin. Mol. Ther.* 3:244-8, 2001] antisense administration using nanoparticulate systems [Lambert
30 et al., *Adv. Drug. Deliv. Rev.* 47:99-112, 2001], or administration of antisense

nucleotides conjugated with peptide [Juliano et al., *Curr. Opin. Mol. Ther.* 2:297-303, 2000].

The invention further contemplates use of the polynucleotides of the invention for gene therapy or in recombinant expression vectors which produce polynucleotides or polypeptides of the invention that can regulate an activity of LEC genes, and are useful in therapy of LEC disorders such as lymphedema. Delivery of a functional gene encoding a polypeptide of the invention to appropriate cells is effected *ex vivo*, *in situ*, or *in vivo* by use of vectors, including viral vectors (*e.g.*, adenovirus, adeno-associated virus, or a retrovirus), or *ex vivo* by use of physical DNA transfer methods (*e.g.*, liposomes or chemical treatments). See, for example, Anderson, *Nature*, supplement to vol. 392, no. 6679, pp. 25-20 (1998). For additional reviews of gene therapy technology see Friedmann, (*Science*, 244: 1275-1281. 1989); Verma, (*Scientific American*: 263:68-72, 81-84. 1990); and Miller, (*Nature*, 357: 455-460. 1992). Introduction of any one of the nucleotides of the present invention or a gene encoding a polypeptide of the invention can also be accomplished with extrachromosomal substrates (transient expression) or artificial chromosomes (stable expression). Cells may also be cultured *ex vivo* in the presence of proteins of the present invention in order to proliferate or to produce a desired effect on, or activity in, such cells. In another embodiment, cells comprising vectors expressing the polynucleotides or polypeptides of the invention may be cultured *ex vivo* and administered to an individual in need of treatment for an LEC disease or disorder.

Given the foregoing disclosure of the nucleic acid constructs, it is possible to produce the gene product of any of the genes comprising the sequence of any of SEQ ID NOS: 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 and 41 by routine recombinant DNA/RNA techniques. A variety of expression vector/host systems may be utilized to contain and express the coding sequence. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, phagemid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with viral expression vectors (*e.g.*, baculovirus); plant cell systems transfected with virus expression vectors (*e.g.*, Cauliflower Mosaic Virus, CaMV; Tobacco Mosaic

Virus, TMV) or transformed with bacterial expression vectors (*e.g.*, Ti or pBR322 plasmid); or even animal cell systems. Mammalian cells that are useful in recombinant protein productions include, but are not limited to, VERO cells, HeLa cells, Chinese hamster ovary (CHO) cells, COS cells (such as COS-7), WI38, BHK, HepG2, 3T3, RIN, MDCK, A549, PC12, K562 and HEK 293 cells.

Polypeptides of the invention

In general, the isolated LEC polypeptides of the invention are encoded by the above-described differentially expressed LEC polynucleotides of the invention. The sequences of the LEC polypeptides are provided in Table 1 and 2, associated with their known database accession numbers, where applicable. The isolated polypeptides of the invention include, but are not limited to, a polypeptide comprising: the amino acid sequences set forth as any one of SEQ ID NOS.: 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 43, 44, 45, and 46 or an amino acid sequence encoded by any one of the nucleotide sequences set forth in SEQ ID NOS.: 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 43, 44, 45, and 46, or the corresponding full length or mature protein. The invention also provides biologically active or immunologically active variants of any of the amino acid sequences set forth as SEQ ID NOS.: 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 43, 44, 45, and 46, or the corresponding full length or mature protein suitable variant polypeptides have sequences that are at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, 86%, 87%, 88%, 89%, at least about 90%, 91%, 92%, 93%, 94%, preferably at least about 95%, 96%, 97%, more preferably at least about 98%, or most preferably at least about 99% amino acid identity, that retain biological activity. Fragments of the proteins of the present invention which comprise at least 10 contiguous amino acids of a sequence disclosed herein and that are capable of exhibiting a biological activity of the corresponding full length protein are also encompassed by the present invention.

The protein coding sequence is identified in the sequence listing by translation of the disclosed nucleotide sequences. The mature form of such protein may be obtained by expression of a full-length polynucleotide in a suitable

mammalian cell or other host cell. The sequence of the mature form of the protein is also determinable from the amino acid sequence of the full-length form. Where proteins of the present invention are membrane bound, soluble forms of the proteins are also provided. In such forms, part or all of the regions causing the proteins to be
5 membrane bound are deleted so that the proteins are capable of being fully secreted from the cell in which it is expressed.

A variety of methodologies known in the art can be utilized to obtain any one of the isolated polypeptides or proteins of the present invention. At the simplest level, the amino acid sequence can be synthesized using commercially
10 available peptide synthesizers. The polypeptides and proteins of the present invention can alternatively be purified from cells which have been altered to express the desired polypeptide or protein. As used herein, a cell is said to be altered to express a desired polypeptide or protein when the cell, through genetic manipulation, is made to produce a polypeptide or protein which it normally does not produce or which the cell
15 normally produces at a lower level. One skilled in the art can readily adapt procedures for introducing and expressing either recombinant or synthetic sequences into eukaryotic or prokaryotic cells in order to generate a cell which produces one of the polypeptides or proteins of the present invention.

A "fragment" of a polypeptide is meant to refer to any portion of the
20 molecule, such as the peptide core, a variant of the peptide core, or an extracellular region of the polypeptide. A "variant" of a polypeptide is meant to refer to a molecule substantially similar in structure and biological activity to either the entire molecule, or to a fragment thereof. Thus, provided that two molecules possess a similar activity, they are considered variants as that term is used herein even if the composition or
25 secondary, tertiary, or quaternary structure of one of the molecules is not identical to that found in the other, or if the sequence of amino acid residues is not identical. An "analogue" of a polypeptide or genetic sequence is meant to refer to a protein or genetic sequence substantially similar in function and structure to the isolated polypeptide or genetic sequence.

30 It is understood herein that conservative amino acid substitutions can be performed to a purified and isolated polypeptide comprising any one of the

sequences of SEQ ID NOS.: 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 43, 44, 45, and 46 which are likely to result in a polypeptide that retains biological or immunological activity, especially if the number of such substitutions is small. By "conservative amino acid substitution" is meant substitution of an amino acid with an amino acid having a side chain of a similar chemical character. Similar amino acids for making conservative substitutions include those having an acidic side chain (glutamic acid, aspartic acid); a basic side chain (arginine, lysine, histidine); a polar amide side chain (glutamine, asparagine); a hydrophobic, aliphatic side chain (leucine, isoleucine, valine, alanine, glycine); an aromatic side chain (phenylalanine, tryptophan, tyrosine); a small side chain (glycine, alanine, serine, threonine, methionine); or an aliphatic hydroxyl side chain (serine, threonine).

Microarrays

Another aspect of the invention is a composition comprising a plurality of polynucleotide probes for use in detecting gene expression pattern(s) characteristic of particular cell type(s) and for detecting changes in the expression pattern of a particular cell type, e.g., lymphatic endothelial cells. For example, the invention comprehends an array, such as a microarray, comprising polynucleotides having at least 10 contiguous nucleotides selected from the polynucleotide sequences in the sequence listing, or a gene found in any of Tables 1 and 2 herein or Tables 3 and 4 of PCT/US03/06900.

Also contemplated are microarrays comprising polynucleotides having at least 10 contiguous nucleotides selected from the group of SEQ ID NOS.: 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 and 41 as well as any one of the genes found Tables 3 and 4 of PCT/US03/06900. Microarrays of the invention comprise at least 3 polynucleotides, wherein each enumerated polynucleotide has a distinct sequence selected from the group consisting of SEQ ID NOS.: 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 and 41 and a gene found in any of Tables 3 and 4 of PCT/US03/06900. Such microarrays may also have duplicate polynucleotides and additional polynucleotides, e.g., control polynucleotides for use in hybridization-based assays using the microarray. Arrays, including microarrays,

having more than three distinct polynucleotides according to the invention, such as at least five, seven, nine, 20, 50 or more such polynucleotides, will be recognized as arrays according to the invention having the capability of yielding subtle distinctions between biological samples such as various endothelial cell types, or of providing a different, and typically greater, level of confidence in the various uses of such arrays, e.g., in screening for particular endothelial cells, in screening for abnormal or diseases cells and tissues, and the like.

The term "microarray" refers to an ordered arrangement of hybridizable array elements. The array elements are arranged so that there are preferably at least three or more different array elements, more preferably at least 100 array elements, and most preferably at least 1,000 array elements, on a solid support. Preferably, the solid support is a 1 cm² substrate surface, bead, paper, nylon or other type of membrane, filter, chip, glass slide, or any other suitable solid support. The hybridization signal from each of the array elements is individually distinguishable.

In a preferred embodiment, the array elements comprise polynucleotide probes.

Hybridization means contacting two or more nucleic acids under conditions suitable for base pairing. Hybridization includes interaction between partially or perfectly complementary nucleic acids. Suitable hybridization conditions are well known to those of skill in the art. In certain applications, it is appreciated that lower stringency conditions may be required. Under these conditions, hybridization may occur even though the sequences of the interacting strands are not perfectly complementary, being mismatched at one or more positions. Conditions may be rendered less stringent by adjusting conditions in accordance with the knowledge in the art, e.g., increasing salt concentration and/or decreasing temperature. Suitable hybridization conditions are those conditions that allow the detection of gene expression from identifiable expression units such as genes. Preferred hybridization conditions are stringent hybridization conditions, such as hybridization at 42°C in a solution (*i.e.*, a hybridization solution) comprising 50% formamide, 1% SDS, 1 M NaCl, 10% dextran sulfate, and washing for 30 minutes at 65°C in a wash solution comprising 1 X SSC and 0.1% SDS. It is understood in the art that conditions of equivalent stringency can be achieved through variation of

temperature and buffer, or salt concentration, as described in Ausubel, *et al.* (Eds.), Protocols in Molecular Biology, John Wiley & Sons (1994), pp. 6.0.3 to 6.4.10.

Modifications in hybridization conditions can be empirically determined or precisely calculated based on the length and the percentage of guanosine/cytosine (GC) base pairing of the probe. The hybridization conditions can be calculated as described in
5 Sambrook, *et al.*, (Eds.), Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory Press: Cold Spring Harbor, New York (2d. Ed.; 1989), pp. 9.47 to 9.51.

One method of using probes and primers of the invention is in the
10 detection of gene expression in human cells. Normally, the target will be expressed RNAs, although genomic DNA or a cDNA library may be screened. By varying the stringency of hybridization and the target binding site (i.e., the sequence of the probe, corresponding to a subset of one of the sequences set forth at SEQ ID NOS: 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 and 41 or a gene found in any
15 of Tables 3 and 4 of PCT/US03/06900), different degrees of homology are expected to result in hybridization.

The microarray can be used for large-scale genetic or gene expression analysis of a large number of target polynucleotides. The microarray can also be used in the diagnosis of diseases and in the monitoring of treatments. Further, the
20 microarray can be employed to investigate an individual's predisposition to a disease. Furthermore, the microarray can be employed to investigate cellular responses to infection, drug treatment, and the like.

The nucleic acid probes can be genomic DNA or cDNA or mRNA polynucleotides or oligonucleotides, or any RNA-like or DNA-like material, such as
25 peptide nucleic acids, branched DNAs, and the like. The probes can be sense or antisense nucleotide probes. Where target polynucleotides are double-stranded, the probes may be either sense or antisense strands. Where the target polynucleotides are single-stranded, the probes are complementary single strands. In one embodiment, the probes are cDNAs. The size of the DNA sequence of interest may vary and is
30 preferably from 100 to 10,000 nucleotides, more preferably from 150 to 3,500 nucleotides.

The probes can be prepared using a variety of synthetic or enzymatic techniques, which are well known in the art. The probes can be synthesized, in whole or in part, using chemical methods well known in the art (Caruthers et al., *Nucleic Acids Res., Symp. Ser.*, 215-233, 1980).

5 Pharmaceutical Formulations and Routes of Administration

A protein of the present invention (from whatever source derived, such as from recombinant and non-recombinant sources) may be administered to a patient in need, by itself, or in pharmaceutical compositions where it is mixed with suitable carriers, diluents, adjuvants or excipients at doses to treat or ameliorate a variety of disorders. Such a composition may also contain (in addition to protein and a carrier) diluents, fillers, salts, buffers, stabilizers, solubilizers, and other materials well known in the art. The term "pharmaceutically acceptable" means a non-toxic material that does not interfere with the effectiveness of the biological activity of the active ingredient(s). The characteristics of the carrier will depend on the route of administration. The pharmaceutical composition of the invention may also contain cytokines, chemokines, lymphokines, growth factors, or other hematopoietic factors such as a PDGF, a VEGF (particularly a VEGF-C or a VEGF-D), VEGFR-3 (including soluble VEGFR-3 peptides comprising an extracellular domain), M-CSF, GM-CSF, TNF, IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-14, IL-15, IFN, TNF0, TNF1, TNF2, G-CSF, Meg-CSF, thrombopoietin, stem cell factor, and erythropoietin. Various forms of these polypeptides are contemplated as well, such as isolated holoproteins, subunits, fragments (*e.g.*, soluble fragments), and peptide fusions. The pharmaceutical composition may further contain other agents which either enhance the activity of the protein or complement its activity or use in treatment. Such additional factors and/or agents may be included in the pharmaceutical composition to produce a synergistic effect with a protein of the invention, or to minimize side effects. Conversely, a protein of the present invention may be included in formulations of the particular cytokine, lymphokine, other hematopoietic factor, thrombolytic or anti-thrombotic factor, or anti-inflammatory agent to minimize side effects of the cytokine,

lymphokine, other hematopoietic factor, thrombolytic or anti-thrombotic factor, or anti-inflammatory agent. A protein of the present invention may be active in multimers (*e.g.*, heterodimers or homodimers) or complexes with itself or other proteins. As a result, pharmaceutical compositions of the invention may comprise a protein of the invention in such multimeric or complexed form.

Techniques for formulation and administration of the compounds of the instant application may be found in "Remington's Pharmaceutical Sciences," Mack Publishing Co., Easton, Pa., latest edition. A therapeutically effective dose further refers to that amount of the compound sufficient to result in amelioration of symptoms, *e.g.*, treatment, healing, prevention or amelioration of the relevant medical condition, or an increase in rate of beneficial change, healing, prevention or amelioration of such conditions. When applied to an individual active ingredient, administered alone, a therapeutically effective dose refers to that ingredient alone. When applied to a combination, a therapeutically effective dose refers to combined amounts of the active ingredients that result in the therapeutic effect, whether administered in combination, serially or simultaneously.

In practicing methods of treatment or use of the present invention, a therapeutically effective amount of protein of the present invention is administered to a mammal having a condition or disorder to be treated. Protein of the present invention may be administered in accordance with the method of the invention either alone or in combination with other therapies such as treatments employing cytokines, lymphokines or other hematopoietic factors. When co-administered with one or more cytokines, lymphokines or other hematopoietic factors, a protein of the invention may be administered either simultaneously with the cytokine(s), lymphokine(s), other hematopoietic factor(s), thrombolytic or anti-thrombotic factors, or sequentially. If administered sequentially, the attending physician will decide on the appropriate sequence of administering a protein of the invention in combination with a cytokine(s), lymphokine(s), other hematopoietic factor(s), thrombolytic or anti-thrombotic factors.

Routes of Administration

- Suitable routes of administration may, for example, include oral, rectal, transmucosal, or intestinal administration; parenteral delivery, including intramuscular, subcutaneous, intramedullary injections, as well as intrathecal, direct intraventricular, intravenous, intraperitoneal, intranasal, or intraocular injections.
- Administration of protein of the present invention used in the pharmaceutical composition or to practice the method of the present invention is carried out in a variety of conventional ways, such as oral ingestion, inhalation, topical application or cutaneous, subcutaneous, intraperitoneal, parenteral or intravenous injection.
- Intravenous administration to a mammal, such as a human patient, is preferred. Alternately, one may administer the compound in a local rather than systemic manner, for example, via injection of the compound at the site of intended action.

Compositions/Formulations

- Pharmaceutical compositions for use in accordance with the present invention thus may be formulated in a conventional manner using one or more physiologically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. These pharmaceutical compositions may be manufactured in a manner that is itself known, *e.g.*, by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or lyophilizing processes. Proper formulation is dependent upon the route of administration chosen. When a therapeutically effective amount of protein of the present invention is administered orally, protein of the present invention will be in the form of a tablet, capsule, powder, solution or elixir. When administered in tablet form, the pharmaceutical composition of the invention may additionally contain a solid carrier such as a gelatin or an adjuvant. The tablet, capsule, and powder contain from about 5 to 95% protein of the present invention, and preferably from about 25 to 90% protein of the present invention. When administered in liquid form, a liquid carrier such as water, petroleum, oils of animal or plant origin such as peanut oil, mineral oil, soybean oil, or sesame oil, or synthetic oils may be added. The liquid

form of the pharmaceutical composition may further contain physiological saline solution, dextrose or other saccharide solution, or glycols such as ethylene glycol, propylene glycol or polyethylene glycol. When administered in liquid form, the pharmaceutical composition contains from about 0.5 to 90% by weight of protein of the present invention, and preferably from about 1 to 50% protein of the present invention.

When a therapeutically effective amount of protein of the present invention is administered by intravenous, cutaneous or subcutaneous injection, protein of the present invention will be in the form of a pyrogen-free, parenterally acceptable aqueous solution. The preparation of such parenterally acceptable protein solutions, having due regard to pH, isotonicity, stability, and the like, is within the skill in the art. A preferred pharmaceutical composition for intravenous, cutaneous, or subcutaneous injection should contain, in addition to protein of the present invention, an isotonic vehicle such as Sodium Chloride Injection, Ringer's Injection, Dextrose Injection, Dextrose and Sodium Chloride Injection, Lactated Ringer's Injection, or other vehicle as known in the art. The pharmaceutical composition of the present invention may also contain stabilizers, preservatives, buffers, antioxidants, or other additives known to those of skill in the art. For injection, the agents of the invention may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks' solution, Ringer's solution, or physiological saline buffer. For transmucosal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

For oral administration, the compounds can be formulated readily by combining the active compounds with pharmaceutically acceptable carriers well known in the art. Such carriers enable the compounds of the invention to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions and the like, for oral ingestion by a patient to be treated. Pharmaceutical preparations for oral use can be obtained by combination with a solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients are, in particular, fillers such as sugars, including lactose, sucrose,

mannitol, or sorbitol; cellulose preparations such as, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, gum tragacanth, methyl cellulose, hydroxypropylmethyl-cellulose, sodium carboxymethylcellulose, and/or polyvinylpyrrolidone (PVP). If desired, disintegrating agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof such as sodium alginate. Dragee cores are provided with suitable coatings. For this purpose, concentrated sugar solutions may be used, which may optionally contain gum arabic, talc, polyvinyl pyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures.

10 Dyestuffs or pigments may be added to the tablets or dragee coatings for identification or to characterize different combinations of active compound doses.

Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The push-fit capsules can contain the active ingredients in admixture with filler such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In addition, stabilizers may be added. All formulations for oral administration should be in dosages suitable for such administration. For buccal administration, the compositions may take the form of tablets or lozenges formulated in conventional manner.

15 20

For administration by inhalation, the compounds for use according to the present invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebuliser, with the use of a suitable propellant, *e.g.*, dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of, *e.g.*, gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch. The compounds may be formulated for parenteral administration by injection, *e.g.*, by bolus injection or continuous

25 30

infusion. Formulations for injection may be presented in unit dosage form, *e.g.*, in ampoules or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents.

Pharmaceutical formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble form. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions. Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, *e.g.*, sterile pyrogen-free water, before use.

The compounds may also be formulated in rectal compositions such as suppositories or retention enemas, *e.g.*, containing conventional suppository bases such as cocoa butter or other glycerides. In addition to the formulations described previously, the compounds may also be formulated as a depot preparation. Such long-acting formulations may be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds may be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

A pharmaceutical carrier for the hydrophobic compounds of the invention is a cosolvent system comprising benzyl alcohol, a nonpolar surfactant, a water-miscible organic polymer, and an aqueous phase. The cosolvent system may be the VPD co-solvent system. VPD is a solution of 3% w/v benzyl alcohol, 8% w/v of the nonpolar surfactant polysorbate 80, and 65% w/v polyethylene glycol 300, made up to volume in absolute ethanol. The VPD co-solvent system (VPD:5W) consists of

VPD diluted 1:1 with a 5% dextrose-in-water solution. This co-solvent system dissolves hydrophobic compounds well, and itself produces low toxicity upon systemic administration. Naturally, the proportions of a co-solvent system may be varied considerably without destroying its solubility and toxicity characteristics.

- 5 Furthermore, the identity of the co-solvent components may be varied: for example, other low-toxicity nonpolar surfactants may be used instead of polysorbate 80; the fraction size of polyethylene glycol may be varied; other biocompatible polymers may replace polyethylene glycol, *e.g.*, polyvinyl pyrrolidone; and other sugars or polysaccharides may substitute for dextrose. Alternatively, other delivery systems for
- 10 hydrophobic pharmaceutical compounds may be employed. Liposomes and emulsions are well known examples of delivery vehicles or carriers for hydrophobic drugs. Certain organic solvents such as dimethylsulfoxide also may be employed, although usually at the cost of greater toxicity. Additionally, the compounds may be delivered using a sustained-release system, such as semipermeable matrices of solid
- 15 hydrophobic polymers containing the therapeutic agent. Various sustained-release materials have been established and are well known by those skilled in the art. Sustained-release capsules may, depending on their chemical nature, release the compound over a time period of a few weeks up to over 100 days. Depending on the chemical nature and the biological stability of the therapeutic reagent, additional
- 20 strategies for protein stabilization may be employed.

The pharmaceutical compositions also may comprise suitable solid or gel-phase carriers or excipients. Examples of such carriers or excipients include but are not limited to calcium carbonate, calcium phosphate, various sugars, starches, cellulose derivatives, gelatin, and polymers such as polyethylene glycols. Many of

25 the proteinase-inhibiting compounds of the invention may be provided as salts with pharmaceutically compatible counterions. Such pharmaceutically acceptable base addition salts are those salts which retain the biological effectiveness and properties of the free acids and which are obtained by reaction with inorganic or organic bases such as sodium hydroxide, magnesium hydroxide, ammonia, trialkylamine, dialkylamine,

30 monoalkylamine, dibasic amino acids, sodium acetate, potassium benzoate, triethanol amine, and the like.

The pharmaceutical compositions of the invention may be in the form of a complex of a protein(s) of the present invention along with protein or peptide antigens. The pharmaceutical compositions of the invention may be in the form of a liposome in which protein of the present invention is combined, in addition to other
5 pharmaceutically acceptable carriers, with amphipathic agents such as lipids which exist in aggregated form as micelles, insoluble monolayers, liquid crystals, or lamellar layers in aqueous solution. Suitable lipids for liposomal formulation include, without limitation, monoglycerides, diglycerides, sulfatides, lysolecithin, phospholipids, saponin, bile acids, and the like. Preparation of such liposomal formulations is within
10 the level of skill in the art, as disclosed, for example, in U.S. Pat. Nos. 4,235,871; 4,501,728; 4,837,028; and 4,737,323, each of which is incorporated herein by reference.

The amount of protein of the invention in the pharmaceutical composition will depend upon the nature and severity of the condition being treated,
15 and on the nature of prior treatments which the patient has undergone. Ultimately, the attending physician will decide the amount of protein of the present invention with which to treat each individual patient. Initially, the attending physician will administer low doses of protein of the present invention and observe the patient's response. Larger doses of protein of the present invention may be administered until
20 the optimal therapeutic effect is obtained for the patient, and at that point the dosage is not increased further. It is contemplated that the various pharmaceutical compositions used to practice the methods of the invention should contain about 0.01 μ g to about 100 mg (preferably about 0.1 μ g to about 10 mg, more preferably about 0.1 μ g to about 1 mg) of protein of the invention per kg body weight. When administered, the
25 therapeutic composition for use in this invention is in a pyrogen-free, physiologically acceptable form. Therapeutically useful agents other than a protein of the invention which may also optionally be included in the composition as described above, may alternatively or additionally, be administered simultaneously or sequentially with the composition in the methods of the invention.

30 Polynucleotides of the present invention can also be used for gene therapy. Such polynucleotides can be introduced either *in vivo* or *ex vivo* into cells for

expression in a mammalian subject. Polynucleotides of the invention may also be administered by other known methods for introduction of nucleic acid into a cell or organism (including, without limitation, in the form of viral vectors or naked DNA). Cells may also be cultured *ex vivo* in the presence of proteins of the present invention
5 in order to proliferate or to produce a desired effect on or activity in such cells. Treated cells can then be introduced *in vivo* for therapeutic purposes.

Effective Dosage

Pharmaceutical compositions suitable for use in the present invention include compositions wherein the active ingredients are contained in an effective
10 amount to achieve an intended purpose. More specifically, a therapeutically effective amount means an amount effective to prevent development of, or to alleviate the existing symptoms of, the subject being treated. Suitable properties that may be used in determining effective dosages include measurements of LEC and/or BEC growth stimulation or inhibition, rates or extent of cell differentiation into LECs and/or BECs,
15 tendencies of cell expression patterns to shift towards or away from LEC- or BEC-specific expression patterns, and the like. Determination of the effective amounts is well within the capability of those skilled in the art, especially in light of the detailed disclosure provided herein. For any compound used in a method of the invention, the therapeutically effective dose can be estimated initially from cell culture assays. For
20 example, for inhibitory methods, a dose can be formulated in animal models to achieve a circulating concentration range that includes the IC_{50} as determined in cell culture (*i.e.*, the concentration of the test compound which achieves a half-maximal inhibitory concentration). Such information can be used to more accurately determine useful doses in humans.

25 A therapeutically effective dose refers to that amount of the compound that results in amelioration of symptoms or, in the case of life-threatening conditions, a prolongation of survival in a patient. Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, *e.g.*, for determining the LD_{50} (the dose lethal to 50% of the
30 population) and the ED_{50} (the dose therapeutically effective in 50% of the

population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio between LD₅₀ and ED₅₀. Compounds which exhibit high therapeutic indices are preferred. The data obtained from these cell culture assays and animal studies can be used in formulating a range of dosage for use in human. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED₅₀ with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. The exact formulation, route of administration and dosage can be chosen by the individual physician in view of the patient's condition. See, *e.g.*, Fingl et al., 1975, in "The Pharmacological Basis of Therapeutics", Ch. 1 p.1.

The amount of composition administered will, of course, be dependent on the subject being treated, on the subject's weight, the severity of the affliction, the manner of administration and the judgment of the prescribing physician.

Packaging

The compositions may, if desired, be presented in a pack or dispenser device which may contain one or more unit dosage forms containing the active ingredient. The pack may, for example, comprise metal or plastic foil, such as a blister pack. The pack or dispenser device may be accompanied by instructions for administration. Compositions comprising a compound of the invention formulated in a compatible pharmaceutical carrier may also be prepared, placed in an appropriate container, and labeled for treatment of an indicated condition.

In addition, the invention comprehends a use of such a composition to manufacture a medicament for the treatment of a cell or an organism, such as a human patient, having a hyperproliferative or hypoproliferative disorder of a LEC and/or BEC, such as lymphedema, lymphangioma, lymphangiomyeloma, lymphangiomatosis, lymphangiectasis, lymphosarcoma, or lymphangiosclerosis, comprising administering an effective amount, or dose, of a composition according to the invention to the cell or organism. Suitable compositions include, but are not limited to, any polynucleotide according to the invention (*e.g.*, an antisense polynucleotide), any polypeptide according to the invention, an antibody specifically

recognizing a polynucleotide or polypeptide according to the invention, a small molecule compound effective in modulating the expression of a polynucleotide according to the invention, and the like. Also contemplated are uses of compositions according to the invention for the manufacture of a medicament to ameliorate a symptom associated with a LEC- or BEC-associated disease or disorder.

Antibodies

Antibodies are useful for modulating the polypeptides of the invention due to the ability to easily generate antibodies with relative specificity, and due to the continued improvements in technologies for adopting antibodies to human therapy.

Thus, the invention contemplates use of antibodies (e.g., monoclonal and polyclonal antibodies, single chain antibodies, chimeric antibodies, bifunctional/bispecific antibodies, humanized antibodies, human antibodies, domain antibodies and complementary determining region (CDR)-grafted antibodies, including compounds which include CDR sequences which specifically recognize a polypeptide of the invention), specific for polypeptides of interest to the invention. Preferred antibodies are human antibodies, such as those produced in transgenic animals, which are produced and identified according to methods described in WO 93/11236, published June 20, 1993, which is incorporated herein by reference in its entirety. Antibody fragments, including Fab, Fab', F(ab')₂, and Fv, are also provided by the invention.

The term "specific for," when used to describe antibodies of the invention, indicates that the variable regions of the antibodies of the invention recognize and bind the polypeptide of interest at a detectably different, and greater, level that bind to other substances (i.e., able to distinguish the polypeptides of interest from other known polypeptides of the same family, by virtue of measurable differences in binding affinity, despite the possible existence of localized sequence identity, homology, or similarity between family members). It will be understood that specific antibodies may also interact with other proteins (for example, *S. aureus* protein A or other antibodies in ELISA techniques) through interactions with sequences outside the variable region of the antibodies, and in particular, in the constant region of the molecule. Screening assays to determine binding specificity of an antibody of the

invention are well known and routinely practiced in the art. For a comprehensive discussion of such assays, see Harlow et al. (Eds.), *Antibodies A Laboratory Manual*; Cold Spring Harbor Laboratory; Cold Spring Harbor, NY (1988), Chapter 6.

Non-human antibodies may be humanized by any method known in the art. A preferred "humanized antibody" has a human constant region, while the variable region, or at least a complementarity-determining region (CDR), of the antibody is derived from a non-human species. Methods for humanizing non-human antibodies are well known in the art. See U.S. Patent Nos. 5,585,089, and 5,693,762. Generally, a humanized antibody has one or more amino acid residues introduced into its framework region from a source which is non-human. Humanization can be performed, for example, using methods described in Jones *et al.* (*Nature* 321: 522-525, 1986), Riechmann *et al.*, (*Nature*, 332: 323-327, 1988) and Verhoeyen *et al.* [*Science* 239:1534-1536, (1988)], by substituting at least a portion of a rodent CDR for the corresponding regions of a human antibody. Numerous techniques for preparing engineered antibodies are described, *e.g.*, in Owens and Young, (*J. Immunol. Meth.*, 168:149-165, 1994). Further changes can then be introduced into the antibody framework to modulate affinity or immunogenicity.

The invention further provides a hybridoma that produces an antibody according to the invention. Antibodies of the invention are useful for detection and/or purification of the polypeptides of the invention.

Polypeptides and/or polynucleotides of the invention may also be used to immunize animals to obtain polyclonal and monoclonal antibodies which specifically react with the polypeptide. Such antibodies may be obtained using either the entire polypeptide or fragments thereof as an immunogen. The peptide immunogens additionally may contain a cysteine residue at the carboxyl terminus, and may be conjugated to a hapten such as keyhole limpet hemocyanin (KLH). Methods for synthesizing such peptides are known in the art, for example, as taught in R. P. Merrifield, *J. Amer. Chem. Soc.* 85:2149-2154 (1963); J. L. Krstenansky, et al., *FEBS Lett.* 211: 10 (1987). Monoclonal antibodies binding to the protein of the invention may be useful diagnostic agents for the immunodetection of the polypeptide. Neutralizing monoclonal antibodies binding to the polypeptide may also be useful

therapeutics for both conditions associated with the polypeptide and also in the treatment of some forms of cancer where abnormal expression of the polypeptide is involved. In the case of cancerous cells or leukemic cells, neutralizing monoclonal antibodies against the polypeptide are useful in detecting and preventing the metastatic spread of the cancerous cells mediated by the polypeptide. In general, techniques for preparing polyclonal and monoclonal antibodies as well as hybridomas capable of producing the desired antibody are well known in the art (Campbell, A. M., Monoclonal Antibodies Technology: Laboratory Techniques in Biochemistry and Molecular Biology, Elsevier Science Publishers, Amsterdam, The Netherlands (1984); St. Groth et al., *J. Immunol.* 35:1-21 (1990); Kohler and Milstein, *Nature* 256:495-497 (1975), the trioma technique, the human B-cell hybridoma technique (Kozbor et al., *Immunology Today* 4:72, 1983); Cole et al., in *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc. (1985), pp. 77-96).

Any animal (mouse, rabbit, and the like) which is known to produce antibodies can be immunized with a peptide or polypeptide of the invention. Methods for immunization are well known in the art. Such methods include subcutaneous or intraperitoneal injection of the polypeptide. One skilled in the art will recognize that the amount of the polypeptide encoded by an ORF of the invention used for immunization will vary based on the animal which is immunized, the antigenicity of the peptide and the site of injection. The protein that is used as an immunogen may be modified or administered in an adjuvant in order to increase the protein's antigenicity. Methods of increasing the antigenicity of a protein are well known in the art and include, but are not limited to, coupling the antigen with a heterologous protein (such as a globulin or β -galactosidase) or through the inclusion of an adjuvant during immunization.

For monoclonal antibodies, spleen cells from the immunized animals are removed, fused with myeloma cells, such as SP2/0-Ag14 myeloma cells, and allowed to become monoclonal-antibody-producing hybridoma cells. Any one of a number of methods well known in the art can be used to identify the hybridoma cell which produces an antibody with the desired characteristics. These include screening the hybridomas with an ELISA assay, Western blot analysis, or radioimmunoassay

(Lutz *et al.*, *Exp. Cell Research*. 175:109-124, 1988). Hybridomas secreting the desired antibodies are cloned and the class and subclass is determined using procedures known in the art (Campbell, A. M., *Monoclonal Antibody Technology: Laboratory Techniques in Biochemistry and Molecular Biology*, Elsevier Science Publishers, Amsterdam, The Netherlands (1984)). Techniques described for the
5 production of single-chain antibodies (U.S. Pat. No. 4,946,778) can be adapted to produce single-chain antibodies to polypeptide of the invention.

For polyclonal antibodies, antibody-containing antiserum is isolated from the immunized animal and is screened for the presence of antibodies with the
10 desired specificity using one of the above-described procedures. The invention further provides the above-described antibodies in detectably labeled form. Antibodies can be detectably labeled through the use of radioisotopes, affinity labels (such as biotin, avidin, and the like), enzymatic labels (such as horseradish peroxidase, alkaline phosphatase, and the like) fluorescent labels (such as FITC or
15 rhodamine, and the like), paramagnetic atoms, and the like. Procedures for accomplishing such labeling are well-known in the art; for example, see Sternberger, *et al.*, *J. Histochem. Cytochem.* 18:315, 1970; Bayer, *et al.*, *Meth. Enzym.* 62:308, 1979; Engval, *et al.*, *Immunol.* 109:129, 1972; and Goding, *J. Immunol. Meth.* 13:215, 1976.

20 The labeled antibodies of the present invention can be used for *in vitro*, *in vivo*, and *in situ* assays to identify cells or tissues in which a fragment of the polypeptide of interest is expressed. The antibodies may also be used directly in therapies or other diagnostics. The present invention further provides the above-described antibodies immobilized on a solid support. Examples of such solid supports
25 include plastics such as polycarbonate, complex carbohydrates such as agarose and sepharose, and acrylic resins such as polyacrylamide and latex beads. Techniques for coupling antibodies to such solid supports are well known in the art (Weir, D. M. *et al.*, "Handbook of Experimental Immunology" 4th Ed., Blackwell Scientific Publications, Oxford, England, Chapter 10 (1986); Jacoby, W. D. *et al.*, *Meth. Enzym.*
30 34 Academic Press, N.Y. (1974)). The immobilized antibodies of the present

invention can be used for *in vitro*, *in vivo*, and *in situ* assays as well as for immuno-affinity purification of the proteins of the present invention.

Computer-Readable Sequences

In one application of this embodiment, a nucleotide sequence of the present invention can be recorded on computer-readable media. As used herein, "computer-readable media" refers to any medium which can be read and accessed directly by a computer. Such media include, but are not limited to, magnetic storage media, such as floppy discs, hard disc storage medium, and magnetic tape; optical storage media such as CD-ROM; electrical storage media such as RAM and ROM; and hybrids of these categories such as magnetic/optical storage media. A skilled artisan can readily appreciate how any of the presently known computer-readable media can be used to create a manufacture comprising computer-readable medium having recorded thereon a nucleotide sequence of the present invention. As used herein, "recorded" refers to a process for storing information on computer-readable medium. A skilled artisan can readily adopt any of the presently known methods for recording information on a computer-readable medium to generate manufactures comprising the nucleotide sequence information of the present invention.

A variety of data storage structures are available to a skilled artisan for creating a computer-readable medium having recorded thereon a nucleotide sequence of the present invention. The choice of the data storage structure will generally be based on the means chosen to access the stored information. In addition, a variety of data processor programs and formats can be used to store the nucleotide sequence information of the present invention on computer readable medium. The sequence information can be represented in a word processing text file, formatted in commercially-available software such as WordPerfect and Microsoft Word, or represented in the form of an ASCII file, stored in a database application, such as DB2, Sybase, Oracle, or the like. A skilled artisan can readily adapt any number of data processor structuring formats (*e.g.*, text file or database) in order to obtain computer readable medium having recorded thereon the nucleotide sequence information of the present invention. By providing the nucleotide sequence of any of

SEQ ID NOS: 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 and 41 or a representative fragment thereof, or a nucleotide sequence at least 99.9% identical to any of SEQ ID NOS: 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 and 41 in computer-readable form, a skilled artisan can routinely access the
5 sequence information for a variety of purposes. Computer software is publicly available which allows a skilled artisan to access sequence information provided in a computer-readable medium. The examples which follow demonstrate how software which implements the BLAST (Altschul et al., *J. Mol. Biol.* 215:403-410, 1990) and BLAZE (Brutlag et al., *Comp. Chem.* 17:203-207, 1993) search algorithms on a
10 Sybase system is used to identify open reading frames (ORFs) within a nucleic acid sequence. Such ORFs may be protein-encoding fragments and may be useful in producing commercially important proteins such as enzymes used in fermentation reactions and in the production of commercially useful metabolites.

As used herein, "a computer-based system" refers to the hardware
15 means, software means, and data storage means used to analyze the nucleotide sequence information of the present invention. The minimum hardware means of the computer-based systems of the present invention comprises a central processing unit (CPU), input means, output means, and data storage means. A skilled artisan can readily appreciate that any one of the currently available computer-based systems are
20 suitable for use in the invention. As stated above, the computer-based systems of the present invention comprise a data storage means having stored therein a nucleotide sequence of the present invention and the necessary hardware means and software means for supporting and implementing a search means. As used herein, "data storage means" refers to memory which can store nucleotide sequence information of
25 the present invention, or a memory access means which can access manufactures having recorded thereon the nucleotide sequence information of the present invention.

As used herein, "search means" refers to one or more programs which are implemented on the computer-based system to compare a target sequence or target structural motif with the sequence information stored within the data storage means.
30 Search means are used to identify fragments or regions of a known sequence which match a particular target sequence or target motif. A variety of known algorithms are

disclosed publicly and a variety of commercially available software for conducting search means are and can be used in the computer-based systems of the present invention. Examples of such software includes, but is not limited to, MacPattern (EMBL), BLASTN and BLASTA (NPOLYPEPTIDEIA). A skilled artisan can readily recognize that any one of the available algorithms or implementing software packages for conducting homology searches can be adapted for use in the present computer-based systems. As used herein, a "target sequence" can be any nucleic acid or amino acid sequence of six or more nucleotides or two or more amino acids. A skilled artisan can readily recognize that the longer a target sequence is, the less likely a target sequence will be present as a random occurrence in the database. The most preferred sequence length of a target sequence is from about 10 to 100 amino acids or from about 30 to 300 nucleotide residues. However, it is well recognized that searches for commercially important fragments, such as sequence fragments involved in gene expression and protein processing, may be of shorter length.

As used herein, "a target structural motif," or "target motif," refers to any rationally selected sequence or combination of sequences in which the sequence(s) are chosen based on a three-dimensional configuration which is formed upon the folding of the target motif. There are a variety of target motifs known in the art. Protein target motifs include, but are not limited to, enzyme active sites and signal sequences. Nucleic acid target motifs include, but are not limited to, promoter sequences, hairpin structures and inducible expression elements (protein binding sequences).

Diagnostic Assays and Kits

The present invention further provides diagnostic assays, and related kits, for hyper- and/or hypo-proliferative disorders or diseases of endothelial cells such as LECs or BECs. These assays comprise methods to identify the presence or expression of one of the ORFs of the present invention, or homolog thereof, in a test sample, using a nucleic acid probe or an antibody according to the invention.

In general, methods for detecting a polynucleotide of the invention can comprise contacting a sample with a compound that binds to and forms a complex

with, the polynucleotide for a period sufficient to form the complex, and detecting the complex, so that if a complex is detected, a polynucleotide of the invention is detected in the sample.

Such methods can also comprise contacting a sample under stringent
5 hybridization conditions with nucleic acid primers that anneal to a polynucleotide of the invention under such conditions, and amplifying annealed polynucleotides, so that if a polynucleotide is amplified, a polynucleotide of the invention is detected in the sample.

In general, methods for detecting a polypeptide of the invention can
10 comprise contacting a sample with a compound that binds to and forms a complex with the polypeptide for a period sufficient to form the complex, and detecting the complex, so that if a complex is detected, a polypeptide of the invention is detected in the sample. In detail, such methods comprise incubating a test sample with one or more of the antibodies or one or more of the nucleic acid probes of the invention and
15 assaying for binding of the nucleic acid probes or antibodies to components within the test sample.

Conditions for incubating a nucleic acid probe or antibody with a test sample vary. Incubation conditions depend on the format employed in the assay, the detection methods employed, and the type and nature of the nucleic acid probe or
20 antibody used in the assay. One skilled in the art will recognize that any one of the commonly available hybridization, amplification or immunological assay formats can readily be adapted to employ the nucleic acid probes or antibodies of the present invention. Examples of such assays can be found in Chard, T., An Introduction to Radioimmunoassay and Related Techniques, Elsevier Science Publishers,
25 Amsterdam, The Netherlands (1986); Bullock, G. R. et al., Techniques in Immunocytochemistry, Academic Press, Orlando, Fla. Vol. 1 (1982), Vol. 2 (1983), Vol. 3 (1985); Tijssen, P., Practice and Theory of immunoassays: Laboratory Techniques in Biochemistry and Molecular Biology, Elsevier Science Publishers, Amsterdam, The Netherlands (1985). The test samples of the present invention
30 include cells, protein or membrane extracts of cells, or biological fluids such as sputum, blood, serum, plasma, or urine. The test sample used in the above-described

method will vary based on the assay format, nature of the detection method and the tissues, and cells or extracts used as the sample to be assayed. Methods for preparing protein extracts or membrane extracts of cells are well known in the art and can be readily be adapted in order to obtain a sample which is compatible with the system
5 utilized.

In another embodiment of the present invention, kits are provided which contain the necessary reagents to carry out the assays of the present invention. In one embodiment, the invention provides a compartment kit to receive, in close confinement, one or more containers which comprises: (a) a first container comprising
10 one of the probes or antibodies of the present invention; and (b) one or more other containers comprising one or more of the following: wash reagents, reagents capable of detecting presence of a bound probe or antibody.

In detail, a compartment kit includes any kit in which reagents are contained in separate containers. Such containers include small glass containers,
15 plastic containers or strips of plastic or paper. Such containers allows one to efficiently transfer reagents from one compartment to another compartment such that the samples and reagents are not cross-contaminated, and the agents or solutions of each container can be added in a quantitative fashion from one compartment to another. Such containers will include a container which will accept the test sample, a
20 container which contains the antibody or antibodies used in the assay, containers which contain wash reagents (such as phosphate-buffered saline, Tris buffers, and the like), and containers which contain the reagents used to detect the bound antibody or probe. Types of detection reagents include labeled nucleic acid probes, labeled
25 enzymatic, or antibody binding reagents which are capable of reacting with the labeled antibody. One skilled in the art will readily recognize that the disclosed probes and antibodies of the present invention can be readily incorporated into one of the established kit formats which are well known in the art.

EXAMPLES

Each reference cited in the methods described below is incorporated by reference herein. Methods used in the examples are as follows:

Antibodies

5 Monoclonal antibodies against human VEGFR-3 (clone 2E11D11; see International Patent Application No. PCT/US02/22164, published as WO 03/006104), PAL-E (Monosan), CD31 (Dako), N-cadherin, VE-cadherin, β -catenin and plakoglobin and polyclonal rabbit anti-human podoplanin were used (Breiteneder-Geleff, S., et al., *Am. J. Pathol.* 154:385-394, 1999). Mouse anti-human integrin α 9
10 was provided by Dr. Dean Sheppard (University of California at San Francisco, San Francisco) and Dr. Curzio Rüegg (University of Lausanne Medical School, Lausanne, Switzerland). The fluorochrome-conjugated secondary antibodies were obtained from Jackson Immunoresearch.

Cell Culture and Transfection

15 Human amniotic epithelial cells are cultured in Med199 medium in the presence of 5% fetal calf serum. Human dermal microvascular endothelial cells are obtained from PromoCell (Heidelberg, Germany). Anti-Podoplanin antibodies, MACS colloidal super-paramagnetic MicroBeads conjugated to goat anti-rabbit IgG antibodies (Miltenyi Biotech, Bergisch Gladbach, Germany), LD and MS separation
20 columns and Midi/MiniMACS separators (Miltenyi Biotech) are used for cell sorting according to the instructions of the manufacturer. The isolated cells are cultured on fibronectin-coated (10 μ g/ml, Sigma, St. Louis, MO) plates as described (Mäkinen, T., et al., *EMBO J.* 20:4762-4773, 2001).

Immunofluorescence and Immunohistochemistry

25 To carry out immunofluorescence on isolated epithelial cells, the cells are cultured on coverslips, fixed with 4% paraformaldehyde, permeabilized with 0.1% Triton-X100 in phosphate-buffered saline (PBS) and stained with primary antibodies. For integrin α 9, staining live cells were incubated with the antibody for 15 minutes on

ice before fixation. The cells were further stained with FITC- or TRITC-conjugated secondary antibodies. F-actin is stained using TexasRed-conjugated phalloidin (Molecular Probes, Eugene, OR). Cells are counterstained with Hoechst 33258 fluorochrome (Sigma) and viewed using a Zeiss Axioplan 2 fluorescent microscope.

5 Normal human skin obtained after surgical removal is embedded in Tissue-Tek® (Sakura, The Netherlands), frozen and sectioned. The sections (6 µm) are fixed in cold acetone for 10 minutes and stained with the primary antibodies followed by peroxidase staining using Vectastain Elite ABC kit (Vector Laboratories, Burlingame, CA) and 3-amino-9-ethyl carbazole (Sigma, St. Louis, MO).

10 RNA isolation, Northern blotting and microarray analyses

 Total RNA was isolated and DNaseI treated in RNeasy columns (Qiagen, Valencia, CA). ³²P-labeled probes for hybridization with the Atlas filters (Clontech) were prepared using 2-5 µg of total RNA according to the manufacturer's instructions with the exception that the probe was purified using Nick-25 columns
15 (Pharmacia Biotech, Uppsala, Sweden). Following hybridizations and washes, the membranes were analyzed using a Fuji BAS 100 phosphoimager. For the Affymetrix® analysis, 2-5 RNA samples isolated from LEC and BEC cell cultures were hybridized separately using RNA extracted from four lots of cells isolated from different individuals. For the Affymetrix® expression analysis, 5 µg of total RNA
20 was used for the synthesis of double-stranded cDNA using Custom SuperScript ds-cDNA Synthesis Kit (Invitrogen, Carlsbad, CA). Biotin-labeled cRNA was then prepared using the Enzo BioArray™ HighYield™ RNA Transcript Labeling Kit (Affymetrix, Santa Clara, CA), and the unincorporated nucleotides were removed using RNeasy columns (Qiagen, Valencia, CA). The hybridization, washing and
25 staining of Human Genome U133A microarrays were done according to the instructions of the manufacturer (Affymetrix, GeneChip Expression Analysis Technical Manual). The probe arrays were scanned at 570 nm using an Agilent GeneArray® Scanner and the readings from the quantitative scanning were analyzed by the Affymetrix® Microarray Suite version 5.0 and Data Mining Tool version 3.0.

For the comparison analyses, the hybridization intensities were calculated using a global scaling intensity of 100.

The differentially expressed sequences were used for searching EST contigs in the GenBank database of the National Center for Biotechnology Information and the National Library of Medicine. (NCBI/NLM), and open reading frames were predicted using the orf finder software available at NCBI/NLM. The SOSUI system was used for prediction of transmembrane helices and signal sequences from the protein sequences, and other protein domain architectures were analyzed using Pfam (Protein families database of alignments and HMMs).

It is expected that the amino acid sequence of a given polypeptide, whether obtained from the Swissprot database, which provides the sequences for typical Pfam analyses, or a Genbank sequence, would be substantially, if not exactly, identical over a typically predominating overlap between those sequences. Moreover, the polypeptides characterized by either of the sequences would be expected to have the same or substantially similar biological activities. Differences in the two sequences may arise due to the fact that some polypeptides identified and used in the Pfam analyses may contain domains designated as "low complexity" or "coiled coil" that are not included in the sequence of a final polypeptide product, as provided by Genbank. Additionally, the Genbank polynucleotide sequences provided herein, which encode the polypeptides disclosed herein, are expected to correspond to polynucleotides encoding the polypeptides utilized in the Pfam analysis.

EXAMPLE 1

IDENTIFICATION OF CELL SURFACE PROTEINS EXPRESSED DIFFERENTIALLY IN LEC

Lymphatic endothelial cells (LEC) were isolated from cultures of human dermal microvascular endothelial cells using magnetic microbeads and antibodies against the lymphatic endothelial cell surface marker podoplanin (Breiteneder-Geleff, et al., *Am. J. Pathol.* 154:385-394 (1999); Mäkinen, et al., *EMBO J.* 20:4762-4773 (2001)), as described in PCT/US03/06900. The purities of the isolated LEC populations were confirmed to be over 99% as assessed by immunofluorescence using antibodies against VEGFR-3 or podoplanin. The isolated

cells were cultured for a couple of passages, and RNA was extracted from the cultures and used for hybridization with oligonucleotide microarrays containing sequences from about 12,000 known genes, *i.e.*, approximately 1/3 of the total number of all predicted human transcripts.

5 PCT/US03/06900, which examined LEC specific genes based on the 95Av2 and 9513-E arrays, identified approximately 165 LEC specific genes in several different protein families, such as adhesion molecules, cytoskeletal proteins, extracellular matrix proteins transcription factors and growth factors.

Experiments were performed to characterize LEC gene expression
10 profiles revealed by the U133A microarray. The Human Genome U133 (HG-U133) Set, consisting of two GeneChip® arrays, contains almost 45,000 probe sets representing more than 39,000 human gene transcripts derived from approximately 33,000 well-substantiated human genes. This set design uses sequences selected from GenBank®, dbEST, and RefSeq. The HG-U133A Array includes representation of
15 the RefSeq database sequences and probe sets. The HG-U133B Array contains primarily probe sets representing EST clusters. The U133A array utilized herein provides a greater sampling of human genes and ESTs (approximately 20,000 more genes) than the arrays described previously in PCT/US03/06900.

Differentially expressed genes were delineated from the U133A array
20 if their signal ratio was over 1.1 and p-value below 0.05, resulting in over 600 genes found to be differentially expressed between LECs and BECs.

LEC-specific genes were further analyzed using a subtraction library involving LEC and BEC genes. To construct the library, total RNA was isolated as previously described and 5µg of total RNA was pre-amplified using a SMART™
25 PCR cDNA synthesis kit (BD Biosciences Clontech). After RsaI-digestion, PCR-Select cDNA subtraction was carried out in both directions, resulting in selective amplification of differentially expressed sequences, and subtracted LEC and BEC cDNA libraries were prepared (BD Biosciences Clontech). Subtractive hybridization was performed with a 1 (tester): 30 (driver) ratio in both directions and subtracted
30 cDNA pools were amplified by PCR. Forty nanograms of the purified PCR-amplified product were cloned into the pAtlas vector (PUC-based vector) for the construction of

subtracted libraries, although a number of other vectors could be used in the construction, as would be known in the art.

Differential screening of the subtracted LEC-specific library was carried out as described in the PCR-Select Differential Screening Kit User Manual (BD Biosciences, Clontech). The LEC-specific subtracted library was plated and individual bacterial clones were picked and grown. After DNA extraction, the inserts were amplified by PCR and used for sequencing. An aliquot of each PCR-amplified insert was also arrayed onto a nylon membrane and used for hybridization with ³²P-labeled cDNA probes. The results from the hybridizations with subtracted LEC-specific (tester) and subtracted BEC-specific (driver) cDNA probes were used for the differential expression analyses.

BLAST (The Basic Local Alignment Search Tool) was used to compare the sequences against nucleotide, protein and EST sequence databases. For unknown sequences, EST contigs were searched and open reading frames were predicted using ORF finder. Protein domain architectures were analyzed using Pfam (Protein families database of alignments and HMMs) and Smart (Simple Modular Architecture Research Tool).

The nucleotide sequences of clones that were differentially expressed in LECs versus BECs were analyzed in the manner described above. Several of the EST or unknown gene fragments detected in the first screen have been investigated further to determine their sequence similarities to known gene sequences and to identify any open reading frames and functional domain similarities.

Several of the LEC-specific genes have been found to correspond to KIAA gene sequences, which are large nucleotide EST clones encoding unknown human proteins. (Kazusa DNA Research Institute, 1532-3, Yana Kisarazu, Chiba, 292-0812, Japan). These LEC-specific genes were further analyzed in several available databases to determine the existence of species homologs and the percent similarity in these homologs and also to reveal amino acid sequences that demonstrate similarity to conserved protein domains.

Analyses of the LEC clone sequences were performed using the HomoloGene database maintained by the U.S. National Center for Biotechnology

Information offered by the National Institutes of Health to determine species homologs and orthologs and their percent similarity to the newly isolated human LEC-specific genes. Analyses of the sequences were performed using a resource of curated and calculated homologs for genes as represented by UniGene or by
 5 annotation of genomic sequences, generally comparing EST and mRNA sequences from UniGene, as well as transcripts extracted from annotated genomic sequences. (Zhang, *et al.*, *J. Comp. Biol.* 7:203-14. 2000). The best match for a nucleotide
 sequence in one organism to a nucleotide sequence in a second organism is based on the degree of similarity between the two sequences, with a minimum alignment of 100
 10 base pairs. The similarity between the two sequences was determined by an alignment score. The alignment score for a sequence pair is the sum of the similarity scores of the sections of the two sequences that aligned.

BLAST and Pfam analysis of the isolated genes revealed numerous LEC-specific transmembrane proteins, including KIAA1972, KIAA0472, KIAA0494,
 15 ATP10D, KIAA1724, DKFZp564K142, KIAA1573, DKFZp761N1114, and FLJ25916, set out in SEQ ID NOS: 6, 8, 10, 12, 14, 16, 20, 22, 24, 42 and 43 (see Table 1).

Table 1

LEC-Specific Transmembrane Proteins

Protein	Genbank Accession Number	nucleotide SEQ ID NO:	polypeptide SEQ ID NO:
KIAA1972	XM_166279	SEQ ID NO: 5	SEQ ID NO: 6
KIAA0472	XM_290898	SEQ ID NO: 7	SEQ ID NO: 8
KIAA0494	NM_014774	SEQ ID NO: 9	SEQ ID NO: 10
ATP10D	NM_020453	SEQ ID NO: 11	SEQ ID NO: 12
KIAA1724	XM_040280	SEQ ID NO: 13	SEQ ID NO: 14
DKFZp564K142	NM_032121	SEQ ID NO: 15	SEQ ID NO: 16
KIAA1573	NM_020925	SEQ ID NO: 19	SEQ ID NO: 20
DKFZp761N1114	AI332934/ NM_181644	SEQ ID NO: 21	SEQ ID NO: 22

FLJ25916	AK098782/ NM 207347	SEQ ID NO: 23	SEQ ID NO: 24
----------	------------------------	---------------	---------------

KIAA1972: LEC-specific gene KIAA1972 (Genbank Accession No. XM_166279; SEQ ID NOS: 5 and 6), a 3540 bp mRNA, has been detected by Northern blot expression at high levels in liver, kidney, testis, spleen and ovary, at moderate levels in heart, lung, smooth muscle, pancreas, spinal cord, and fetal liver, and at low levels in brain and fetal brain. Northern blot data was derived from the electronic database of Human Unidentified Gene-Encoded (HUGE) Large Proteins maintained by Kazusa DNA Research Institute, Chiba, Japan (see Ohara et al., *DNA Res.* 4:53-59, 1997).

Unigene comparison of the KIAA1972 sequence indicates the protein is 36% identical [over a string of 96 amino acids (aa)] to human Ryanodine receptor 2, RYR2, which is a cardiac muscle-type ryanodine receptor calcium release channel. KIAA1972 also shows 34% identity (over a string of 110 aa) to a fragment of mouse (*Mus musculus*) ryanodine receptor, demonstrates 29% identity (over a sequence of 152 aa) to a *Drosophila melanogaster* ryanodine receptor/calcium release channel, shows 48% identity (over a string of 497 aa) to a *Caenorhabditis elegans* zinc finger, C3HC4 type (RING finger) protein, and shows 39% identity (over 140 aa) to an *Arabidopsis thaliana* hypothetical protein. KIAA1972 also demonstrates similarity to species orthologs *R. norvegicus* LOC291860 (92.4% similarity) and *M. musculus* 4930470D19Rik (87% similarity).

Protein domain mapping for the KIAA1972 sequence using the Pfam database (SwissProt designation Q96DX4) predicts a protein of 576 amino acids with a signal peptide from amino acids 1-16, a SPRY domain from approximately residues 358-482 and a RING domain from approximately residues 527-561. The SPRY domain is derived from similarities to the *spla* motif and the ryanodine receptor. The SPRY domains have homologues in butyrophilin/marenostrin/pyrin proteins. Ca^{2+} -release from the sarcoplasmic or endoplasmic reticulum, the site of intracellular Ca^{2+} storage, is mediated by the ryanodine receptor (RyR) and/or the inositol trisphosphate receptor (IP3R).

The RING-finger domain is a specialized type of Zn-finger region of 40 to 60 amino acid residues that binds two zinc atoms, and is probably involved in mediating protein-protein interactions. RING domains contain cysteine and histidine residues for binding to the Zinc molecules in either a C3HC4-or a C3H2C3 configuration. E3 ubiquitin-protein ligase activity is intrinsic to the RING domain of c-Cbl and is likely to be a general function of the RING domain. E3 ubiquitin-protein ligase Mdm2 expression is amplified in certain tumors (including soft tissue sarcomas, osteosarcomas and gliomas), indicating that ubiquitin-ligase function may be important in maintaining cell homeostasis and regulating cell cycle activities.

Therefore, KIAA1972 is expected to act to regulate calcium storage in lymphatic endothelial cells and to function in the regulation of cell homeostasis. Deregulation of KIAA1972 is expected to result in aberrant LEC proliferation and lymphatic disorders such as lymphedema, lymphangioma, lymphangiomyeloma, lymphangiomatosis, lymphangiectasis, lymphosarcoma, and lymphangiosclerosis.

KIAA0472: Full-length KIAA0472, first isolated from the human adult brain genome, is a 2589 bp mRNA sequence. Unigene comparison of the KIAA0472 gene (Genbank Accession No. XM_290898; SEQ ID NOS: 7 and 8) indicates that the predicted protein is 65% identical (within a sequence of 102 aa) to human neuronal thread protein, shares 29% identity (over 245 aa) to *M. musculus* death domain-containing protein RIP; and demonstrates 33% identity (over 139 aa) with rat *R. norvegicus*) protein kinase C, delta type, NPKC-delta. KIAA0472 also demonstrates significant identity to several other protein kinases such as *D. melanogaster* TGF-beta activated kinase 1, DmTak1 (27% identity over 262 aa), a *C. elegans* tyrosine-protein kinase (25% identity over 293 aa) and an *A. thaliana* protein kinase (38% identity over 161 aa).

Protein domain mapping of the predicted 390-amino-acid KIAA0472 gene (SwissProt designation O75060) reveals that a 365 amino acid fragment of the polypeptide (SEQ ID NO: 42) possesses a protein kinase domain from approximately residues 88-342. Pfam analysis predicts the domain lies at residues 88-333 while SMART analysis predicts a STYKc domain, from approximately residues 88-342. There is also a region of similarity to a potential transmembrane domain from amino acids 262-284. The designation of the STYKc domain leads to the expectation that

KIAA0472 is a protein kinase of unclassified specificity, and is more specifically expected to be a phosphotransferase exhibiting dual-specificity as a Ser/Thr/Tyr kinase. Both serine/threonine and tyrosine protein kinases share a common catalytic core comprising a glycine-rich region of amino acids in proximity to a lysine residue which has been shown to be involved in ATP binding, as well as a conserved aspartic acid residue which is important for the catalytic activity of the enzyme.

KIAA0494: KIAA0494 is a mRNA sequence of 5766 base pairs first isolated from human adult brain. Unigene comparison of the KIAA0494 sequence (Genbank Accession No. NM_014774; SEQ ID NOS: 9 and 10) indicates the predicted protein is 23% identical (over a sequence of 306 amino acids) to *Arabidopsis thaliana* hypothetical protein F5E19_70. Analysis of potential species orthologs revealed that KIAA0494 is similar to *R. norvegicus* LOC298425 (85.7% similarity) and *M. musculus* 4732418C07Rik (82.2% similarity).

Protein domain mapping using the Pfam database indicates that the 495-amino-acid KIAA0494 protein (SwissProt designation O75071) exhibits a transmembrane domain from approximately amino acids 71-93; and, coiled-coil regions from amino acids 95-115, 209-229 and 292-312. Both transmembrane and coiled coil domains are found in proteins with diverse functions. Presence of the transmembrane domains indicates the protein is membrane bound. Coiled-coil domains are structural domains composed of two or three alpha helices that wind around each other and are found in a variety of proteins.

ATP10D: Screening for differential expression of LEC- and BEC-specific genes identified the ATP10D gene (Genbank Accession No. NM_020453; SEQ ID NOS: 11 and 12) as a gene upregulated in lymphatic endothelia. ATP10D was isolated as a 6550 bp mRNA sequence. Analysis of related species orthologs detects an ATP10D ortholog in *R. norvegicus* LOC305314 (87.2 %).

Unigene comparison of the ATP10D gene sequence indicates that the predicted ATP10D protein demonstrates 99% identity, over a 650 aa length of aligned region, to human phospholipid-transporting ATPase VD (ATPVD); demonstrates 56% identity, over a 580 aa alignment, to a mouse potential phospholipid-transporting ATPase VA; demonstrates 23% identity (over a 365 amino acid alignment) to rat ATPase isoform 2, Na⁺K⁺ transporting, beta polypeptide 2; shows 38% identity (604

aa alignment) to an *A. thaliana* putative chromatin granule ATPase II homolog; and is 41% identical (over a 570 aa alignment) to a *C. elegans* ATPase.

Human ATP10D is predicted to be a 1426-amino-acid protein. Protein domain mapping of the 1416 amino acid mouse homolog of human ATP10D (SwissProt designation Q8K2X1) demonstrates five transmembrane domains, from amino acids 7-25, 53-75, 88-105, 115-137, and 157-176, and also to possess a signal peptide from residues 1-23. The mouse ATP10D also contains a hydrolase domain from amino acids 805-1074. Hydrolase enzymes catalyze the hydrolysis of various molecular bonds and include such proteins as epoxide hydrolases and phosphatases.

Human ATP10D also contains predicted regions of similarity to MgtA domains involved in cation transport, from approximately residues 70-457 and residues 722-1329 of SEQ ID NO: 12, and is expected to be a cation transport ATPase, similar to other ATPase transporters that are involved in inorganic ion transport and metabolism. Other possible functions include cation transport, integral to membrane, magnesium ion binding, hydrolase activity, ATPase activity, and phospholipid-translocating ATPase activity. ATP10D is expected to play a significant role in transporting ions across the LEC membrane and regulating varied cellular activities.

KIAA1724: KIAA1724 (Genbank Accession No. XM_040280; SEQ ID NOS: 13 and 14) was isolated as a LEC-specific, 8083 bp mRNA sequence. Tissue expression analysis by Northern blot (data derived from HUGE protein database maintained by Kazusa DNA Research Institute, Chiba, Japan) detected KIAA1724 transcripts at highest levels in *B. amygdala*, *B. cerebellum*, *B. substantia nigra*, and *B. thalamus*, with moderate expression in overall brain, liver, *B. corpus callosum*, *B. caudate nucleus*, *B. hippocampus* and the *B. subthalamic nucleus*. Analysis of species similarities led to the identification of species orthologs in *R. norvegicus* LOC298856 (88.4 %) and *M. musculus* 4933402G07Rik (84.4 %)

Unigene analysis of the KIAA1724 sequence reveals that the sequence demonstrates 64% identity, over a string of 49 amino acids, to a probable human thromboxane A2 receptor isoform beta, and 34% identity, over a 377 amino acid alignment, to *C. elegans* hypothetical protein F54D7.2.

Protein domain mapping of a 400-residue fragment comprising the predicted 386 residue KIAA1724 protein (SwissProt designation Q9COD9) indicates the protein contains a CDP-alcohol phosphatidyltransferase (CDP-OH P transf) domain from amino acids 108-252, and 10 transmembrane domains located at residues 61-83; 98-117; 137-159; 164-186; 193-215; 235-257; 270-292; 305-324; 331-53; and 358-380. Phosphatidyltransferases are involved in phospholipid biosynthesis. Generally, these enzymes are proteins of from 200 to 400 amino acid residues exhibiting a conserved region that contains three aspartic acid residues and is located in the N-terminal region of the proteins. KIAA1724 is expected to function in phospholipid biosynthesis in LECs.

DKFZp564K142: Analysis of the genome for LEC-specific genes identified hypothetical gene DKFZp564K142 (Genbank Accession No. NM_032121; SEQ ID NOS: 15 and 16), which comprises a nucleotide sequence of 2788 bp, and exhibits higher expression in LECs versus blood endothelial cells. Initial analysis of the sequence revealed DKFZp564K142 species orthologs in *S. scrofa* Ssc.16415 (93.9%), *R. norvegicus* IAG2 (87.8 %) and *M. musculus*, 2610529C04Rik (79.7 %).

Unigene analysis of DKFZp564K142 indicates that the amino acid sequence of the protein is 69% identical over a 323 aa region to the protein encoded by human gene N33; is 93% identical, over 332 aa, to the rat implantation-associated protein, IAG2; and is 42% identical (over a 303 aa sequence) to the *C. elegans* ZK686.3 protein.

Pfam analysis of the 335 amino acid hypothetical DKFZp564K142 protein (SwissProt designation Q9HOU3) indicates that the protein comprises an OST3-OST6 domain located from approximately residues 20-330. The proteins in the yeast OST3-OST6 family are part of a complex of eight endoplasmic reticulum (ER) proteins that transfer core oligosaccharide from dolichol carrier to Asn-X-Ser/Thr motifs. This family includes both OST3 and OST6, each of which contains four predicted transmembrane helices. Disruption of OST3 and OST6 leads to a defect in the assembly of the protein complex. Hence, the function of these genes is expected to be essential for recruiting a fully active complex necessary for efficient N-glycosylation.

The DKFZp564K142 protein also contains a signal peptide at amino acids 1-29 and multiple transmembrane domains, similar to either the OST3 or OST6 protein, which are located at approximately amino acids 7-29, 186-205, 212-234, 266-288, and 301-320. Both rat IAG2 and DKFZp564K142 exhibit five transmembrane domains. Based on the high similarity to rat IAG2, DKFZp564K142 is expected to be a human homolog to implantation-associated protein involved in pregnancy and may play a role in establishing lymphatic endothelial cells during pregnancy and embryo implantation.

KIAA1573: KIAA1573 (Genbank Accession No. NM_020925; SEQ ID NOS: 19 and 20) encodes a LEC-specific, hypothetical human protein and is approximately 5565 base pairs in length. Northern blot analysis of mRNA expression (data derived from HUGE protein database maintained by Kazusa DNA Research Institute, Chiba, Japan) shows that KIAA1573 is expressed at high levels in B. thalamus, with lower expression in other tissues, such as B. caudate nucleus, B. hippocampus, B. subthalamic nucleus, B. substantia nigra, ovary, brain and kidney. Comparison of KIAA1573 to other species leads to the expectation of orthologs such as *R. norvegicus* LOC298267 (90.6%), *B. taurus* Bt.1428 (88.6%) and *M. musculus* 1190007F10Rik (83.2%).

Unigene analysis of the predicted amino acid sequence of KIAA1573 reveals that the protein demonstrates 22% identity (over a 598 aa sequence) to human calcium channel, voltage-dependent, alpha 2 delta calcium channel subunit; demonstrates 22% identity (within a sequence of 676 aa) to mouse calcium channel, voltage-dependent, alpha 2 delta calcium channel subunit; and 19 % identity (over 677 aa) to rat CIC2_RAT Dihydropyridine-sensitive L-type, calcium channel alpha-2/delta subunit precursor.

Pfam protein domain analysis of a 1185 residue polypeptide sequence (SEQ ID NO: 43) comprising the predicted 985 amino acid KIAA1573 protein (Swissprot designation Q9HCJ9) reveals two Cache domains in KIAA1573, a full Cache domain from amino acids 364-443 and a partial domain from 683-764. The protein also possesses a single transmembrane domain spanning residues 1006-1028 of SEQ ID NO: 43, and is expected to be expressed on the cell surface. The cache

motif is a signaling domain that is found in animal calcium channel subunits and a certain class of prokaryotic chemotaxis receptors.

Analysis of the protein sequence using the Smart [Simple Modular Architecture Research Tool] database predicts the presence of a VWA domain from residues 137-333. The VWA domain is derived from von Willebrand factor A, which is a large multimeric glycoprotein found in blood plasma. This vWF domain is found in various plasma proteins: complement factors B, C2, CR3 and CR4; the integrins (I-domains); collagen types VI, VII, XII and XIV; and other extracellular proteins. The majority of VWA-containing proteins are extracellular. A common feature of proteins with this domain appears to be involvement in multiprotein complexes. Proteins that incorporate vWF domains participate in numerous biological events (e.g. cell adhesion, migration, homing, pattern formation, and signal transduction), involving interaction with a large array of ligands. VWA domains in extracellular eukaryotic proteins mediate adhesion via metal ion-dependent adhesion sites (MIDAS). A number of human diseases arise from mutations in VWA domains, including diseases of the lymphatic endothelium, such as bleeding disorders. The similarity of KIAA1573 to extracellular, adhesion type proteins leads to the expectation that it acts as a mediator of cell signaling, while the presence of the VWA domain implicates KIAA1573 as a useful target in regulating LEC function in lymphatic endothelium mediated bleeding disorders.

DKFZp761N1114: DKFZp761N1114 (Genbank Accession No. AI332934, NM_181644; SEQ ID NOS: 21 and 22) has a nucleotide sequence of 3352 bp. Tissue expression analysis of human tissues, as detected by Multiple Tissue Northern Blot (Clontech) of adult human RNA, indicates that DKFZp761N1114 produces a 4.5 kb transcript, which is expressed at high levels in the stomach, kidney, and brain, with lower expression in other tissues, such as skeletal muscle, colon, placenta, lung, tongue and trachea. Comparison of the sequence with other species leads to expected orthologs in *R. norvegicus* LOC304787 (85.9 %) and *M. musculus* - A930031D07Rik (80.3 %).

Unigene analysis of the predicted protein sequence shows that DKFZp761N1114 is 60% identical (over a 100 aa sequence) to human neuronal thread protein.

Protein domain mapping reveals the 403 amino acid DKFZp761N1114 (SwisProt designation Q8N468) protein contains a signal sequence and 7 apparent transmembrane domains, from amino acids 27-49; 108-130; 194-216; 236-258; 265-287; 291-313; 326-348; and 353-375. DKFZp761N1114 may also contain a fucose permease domain from approximately residues 202-376, which is involved in carbohydrate transport and metabolism. As such, DKFZp761N1114 is expected to act as a unique seven-transmembrane receptor on the surface of LECs and to modulate or regulate carbohydrate activity in the cell.

Further analysis revealed that the DKFZp761N1114 gene encodes an alternative predicted protein structure which includes a transmembrane protein with only a single transmembrane domain and a MSF-1 domain (Major Facilitator Superfamily), a domain present for example in glycerol-3-phosphate transporter from *E. coli* bacteria. An additional alternative predicted domain structure indicates DKFZp761N1114 may encode twelve transmembrane domains and an amino-terminal signal peptide.

The gene is expressed in lymphatic endothelial cells (LECs), but not in blood vascular endothelial cells (BECs), in an *in vitro* cell culture of primary endothelial cells (HDMEC, PromoCell).

In vivo, the gene is expressed, as revealed by non-radioactive *in situ* hybridization (NR-ISH), in lymphatic vessels of mouse embryonic mesenterium at E17.5, but is not detectable in blood vessels. The antisense transcript, as detected by NR-ISH using the sense probe, is also expressed in the mesenteric lymphatic vessels, but to a lesser extent. Thus, the expression of DKFZp761N1114 is expected to be modulated at least in part, by its antisense transcript. Using frozen sections from E16.5 mouse embryos, very little expression in embryonic mouse tissues was detected.

A Multiple Tissue Northern Blot (Clontech) of adult mouse RNA, indicates the DKFZp761N1114 gene is expressed in kidney and to a lesser extent in liver, brain and heart, but not in lung, testis, spleen or skeletal muscle. NR-ISH analysis showed that the gene was expressed in collecting tubules in the kidney of adult mice.

The expression of DKFZp761N1114 in kidney tubules and its domain structure, involving a Major Facilitator Superfamily (MFS) domain, indicate a function in uptake of substances from the extracellular space in kidney, as well as in lymphatic vessels.

5 *FLJ25916*: Screening of the LEC library also identified hypothetical protein FLJ25916 (Genbank Accession No. AK098782; SEQ ID NOS: 23 and 24) as a LEC-specific gene. FLJ25916 was isolated from fis, clone CBR04903 as a 1893 bp fragment.

10 Comparison of the putative coding region of the FLJ25916 mRNA to other sequences reveals that FLJ25916 is similar to human hypothetical protein KIAA0233 (44% identity over a 572 amino acid sequence) and is 28% identical (over 315 aa) to *C. elegans* protein T20D3.11.p. KIAA0233, to which FLJ25916 shows significant identity, demonstrates homology to both the mouse collagen alpha chain precursor and rat procollagen (26% identity with both proteins, over a sequence of
15 approximately 1005 aa). Ortholog analysis for FLJ 25916 predicts similarity to mouse protein ENSMUSP00000059624 (novel Ensembl prediction).

 Protein domain mapping of the hypothetical 583 amino acid FLJ25916 protein (SwissProt designation Q8N787) identified a signal peptide from residues 1-18, nine transmembrane domains located at approximately amino acids 2-19; 50-72;
20 220-242; 247-264; 269-291; 340-362; 401-423; 427-444; and 457-479, and also predicted a coiled-coil region from amino acids 169-189. The similarity of FLJ25916 to other mammalian collagen-like proteins leads to the expectation that it acts as a structural protein and/or interacts with the extracellular matrix surrounding the LEC.

 Further analysis shows that FLJ25916, also identified as C18orf30, is
25 expressed in lymphatic endothelial cells (LECs) and to a lesser extent in blood vascular endothelial cells (BECs) in an *in vitro* cell culture of primary endothelial cells (HDMEC, PromoCell). *In vivo*, the gene is expressed in the trigeminal ganglion in the brain, in the eye and in bone and is expressed weakly in submandibular gland, kidney and lung, as revealed by non-radioactive *in situ* hybridization (NR-ISH) of
30 frozen sections from E16.5 mouse embryos.

The above transmembrane and cell-surface-expressed proteins are useful for the identification and isolation of LEC, and useful for regulation of LEC activity.

EXAMPLE 2

IDENTIFICATION OF INTRACELLULAR PROTEINS EXPRESSED DIFFERENTIALLY IN LEC

Assays using the microarray and LEC cDNA libraries identified several LEC-specific genes encoding proteins expressed on the cell surface and which are suitable targets for identification and localization of lymphatic endothelial cells, as well as being suitable targets for the manipulation of their function and activity. In addition to these cell-surface LEC proteins, intracellular proteins were identified as being LEC-specific; these proteins are expected to contribute to the regulation of the intracellular function of LEC genes, such as the transcription of particular genes and the transport of necessary proteins through the intracellular space.

By way of example, the analysis of LEC-specific intracellular proteins identified the following genes as being expressed at higher levels in LECs versus BECs: HERC3, KIAA1340, protein FLJ20051, protein FLJ13910, protein FLJ11029, KIAA1102, protein FLJ32029, protein FLJ23594, and protein FLJ10532 (see Table 2).

Table 2

LEC-Specific Transcription Factors or Intracellular Proteins

Protein	Genbank Accession Number	nucleotide SEQ ID NO:	polypeptide SEQ ID NO:
HERC3	NM_014606	SEQ ID NO: 27	SEQ ID NO: 28
KIAA1340	XM_044836	SEQ ID NO: 29	SEQ ID NO: 30
FLJ20051	NM_019087	SEQ ID NO: 31	SEQ ID NO: 32
FLJ13910	NM_022780	SEQ ID NO: 17	SEQ ID NO: 18
FLJ11029	NM_018304	SEQ ID NO: 33	SEQ ID NO: 34

KIAA1102	XM_044461	SEQ ID NO: 35	SEQ ID NO: 36
FLJ32029	NM_173582	SEQ ID NO: 37	SEQ ID NO: 38
FLJ23594	NM_024781	SEQ ID NO: 39	SEQ ID NO: 40
FLJ10532	AK001394	SEQ ID NO: 41	
BIKE protein	NM_017593	SEQ ID NO: 25	SEQ ID NO: 26

HERC3: Screening for LEC-specific genes identified the *HERC3* gene as a gene upregulated preferentially in LECs versus BECs. Unigene comparison of the *HERC3* sequence (Genbank Accession No. NM_014606; SEQ ID NOS: 27 and 28) indicates that the predicted protein is 95% identical (over 891 aa) to human hypothetical protein 2 (fragment); 35% identical (over 337 aa) to mouse rjs protein; shares 31% identity (over 184 aa) to rat ubiquitin ligase Nedd4; demonstrates 37% identity (over 309 aa) to the *A. thaliana* UVB-resistance protein, UVR8; is 24% identical (over a 330 aa sequence) to *D. melanogaster* chromatin-binding protein BJ1; and is 30% identical (over 891 aa) to *C. elegans* hypothetical protein Y48G8AL.1.p.

HERC3 protein domain mapping (SwissProt designation Q15034) reveals four regulator-of-chromosome-condensation (RCC1)-like protein domains found from residues 52-99, 102-152, 155-205, and 208-257. Regulator-of-chromosome-condensation (RCC1) is a eukaryotic protein that binds to chromatin and interacts with ran, a nuclear GTP-binding protein, to promote the loss of bound GDP and the uptake of fresh GTP, thus acting as a guanine-nucleotide dissociation stimulator (GDS). The interaction of RCC1 with ran is believed to play an important role in the regulation of gene expression. RCC1 is a protein that contains seven tandem repeats of a domain of about 50 to 60 amino acids. The RCC1-type of repeat is also found in the X-linked retinitis pigmentosa GTPase regulator.

Mapping of the *HERC3* protein also reveals a HECT domain spanning approximately amino acids 721-1050, Pfam predicting the domain from amino acids 752-1050 and SMART analysis detecting the domain at residues 721-1050. HECT (Homologous to the E6-AP Carboxyl Terminus) domains are typically involved in ubiquitin ligase activity. Proteins containing this domain at the C-terminus include

ubiquitin-protein ligase, which regulates ubiquitination of CDC25, and directly transfers ubiquitin to targeted substrates. Human thyroid receptor interacting protein 12 also contains this domain and is a component of an ATP-dependent multi-subunit protein that interacts with the ligand-binding domain of the thyroid hormone receptor.

5 This protein may have similar function to an E3 ubiquitin-protein ligase such as human ubiquitin-protein ligase E3A, which interacts with the E6 protein of the cancer-associated human papillomavirus types 16 and 18. A cysteine residue in the domain is required for ubiquitin-thiolester formation. The HECT domain is known to form complexes with the ubiquitin-conjugating domain of other proteins, and this
10 domain is involved in ubiquinating proteins to signal their degradation by the proteasome.

Further analysis of protein domains in HERC3 indicate that a region from residues 333-481 is typically associated with proteins containing the classical zinc finger C2H2 domain; and residues 482-605 represent a domain associated with
15 proteins containing lectin c domains.

Based on the foregoing information, HERC3 is expected to function as a regulator of protein degradation in LECs, given the similarity to ubiquitin-ligase proteins. HERC3 is also expected to be involved in chromatin binding, based on the presence of the RCC1 domain.

20 *KIAA1340*: Northern blot expression of KIAA1340 (Genbank Accession No. XM_044836; SEQ ID NOS: 29 and 30) indicates that KIAA1340 mRNA (approximately 4363 bp) is expressed at the highest levels in brain and at moderate levels in lung, ovary and spinal cord. *In situ* expression measured in day 16.5 mouse embryo indicates that the transcript is expressed at the highest level in the
25 thymus and the intestine of developing mice. Expected orthologs of KIAA1340 include *R. norvegicus*, LOC312860 (86.9%), and *M. musculus*, C230080I20Rik (82.4%)

Unigene comparison of the KIAA1340 gene sequence indicates that the protein product of KIAA1340 demonstrates 33% identity, over a 155-aa length of
30 aligned region, to mouse Kelch-like ECH-associated protein 1 (Cytosolic inhibitor of Nrf2); shares 32% identity, over a 155-amino-acid alignment, with rat Kelch-like ECH-associated protein 1 (Cytosolic inhibitor of Nrf2, INrf2); has 36% identity, over

98 aa, with the *A. thaliana* hypothetical protein F7A7.180; has 25% identity, over a 278-aa alignment, with *D. melanogaster* long-form kelch protein; and is 27% identical (over a 151-aa alignment) to *C. elegans* hypothetical protein W02G9.2.

Pfam analysis using a 410 amino acid fragment (SEQ ID NO: 44) of the predicted 505 amino acid KIAA1340 protein (SwissProt designation Q8N334) indicates KIAA1340 exhibits similarity to Kelch-related proteins and contains Kelch domains from amino acids 147-193 and 195-236, and partial Kelch domains from residues 238-277 and 291-340. Comparison domain mapping using the SMART database predicts Kelch domains from 159-206, 207-249, and 250-296. Identification of Kelch domains using SMART analysis reveals the Kelch domains to be located from residues 159-206, 207-249 and 250-296. The Kelch motif is an approximately 50-residue domain, named after the *Drosophila* mutant in which it was first identified, and is found six times in *Drosophila* egg-chamber regulatory protein, in mouse protein MIPP, in a number of poxviruses, in α - and β -scruin, and in galactose oxidase from the fungus *Dactylium dendroides*. The known functions of kelch-containing proteins include: scruin functioning as an actin cross-linking protein; galactose oxidase catalyzing the oxidation of the hydroxyl group at the C6 position in D-galactose; neuraminidase hydrolyzing sialic acid residues from glycoproteins; and kelch from *Drosophila* having a cytoskeletal function in the developing oocyte.

KIAA1340 also demonstrates a BTB/POZ domain at approximately amino acids 6-66 of SEQ ID NO: 44. The BTB (for BR-C, tkk and bab) or POZ (for Pox virus and Zinc finger) domain is present near the N-terminus of a fraction of zinc-finger proteins and in proteins that contain the motif, such as Kelch and a family of pox virus proteins. The BTB/POZ domain mediates homomeric dimerization and, in some instances, heteromeric dimerization. These domains are known to be protein-protein interaction domains found at the N-termini of several C2H2-type transcription factors, as well as Shaw-type potassium channels. POZ domains from several zinc-finger proteins have been shown to mediate transcriptional repression and to interact with components of histone deacetylase co-repressor complexes, including N-CoR and SMRT.

The presence of the Kelch domains in KIAA1340 indicates the protein may have several different functions, while the presence of the Zinc-binding domain

implicates KIAA1340 in protein-protein interactions, which are expected to be involved in the regulation of DNA binding and transcription.

5 *FLJ20051*: FLJ20051 exhibits a coding region of 2501 bp (Genbank Accession No. NM_019087; SEQ ID NOS: 31 and 32) and is expressed as a 4.3 kb transcript. Multiple Tissue Northern Blot (Clontech) of adult human RNA demonstrates that the FLJ20051 transcript is expressed at highest levels in kidney, placenta, heart, and skeletal muscle, and lower levels in the brain, lung, liver, colon, small intestine, thymus, peripheral blood leukocytes, and spleen.

10 Whole mount *in situ* analysis, as well as non-radioactive *in situ* hybridization (NR-ISH), of mouse mesenterium (E17.5, E18.5 or P6) detected FLJ20051 expression in lymphatic vessels of the gut mesenterium but not in arteries or veins, i.e., blood vessels. The antisense transcript, as detected by NR-ISH using the sense probe, is also expressed in the mesenteric lymphatic vessels, but to a lesser extent. These data indicate that the expression of FLJ20051 is controlled, at least in
15 part, by its antisense transcript. These data also indicate that FLJ20051 plays a role in the development of the lymphatic endothelium.

Using *in situ* hybridization of frozen sections from E16.5 mouse embryos, FLJ20051 expression was detected in lung, kidney, brain and hair follicles in the skin. Multiple Tissue Northern Blot (Clontech) of adult mouse RNA revealed
20 that the gene is expressed in kidney, and to a lesser extent in heart, brain and liver, and in still lesser extent in lung and testis, with no detectable expression in spleen or skeletal muscle.

Unigene comparison of the FLJ20051 gene sequence indicates that the predicted protein demonstrates 35% identity (over a 159-amino-acid alignment
25 region) to a mouse homolog of human ADP-ribosylation factor 6; demonstrates 35% identity (over 159 aa) to the rat homolog of ADP-ribosylation factor 6; is 33% identical (over a 160-aa sequence alignment) to *A. thaliana* probable ADP-ribosylation factor At2g18390; shows 35% identity (over a 154-aa alignment) to ADP-ribosylation factor 3 of *D. melanogaster*; and is 34% identical (over a 168-aa
30 sequence) to *C. elegans* ADP-ribosylation factors of the Arf family, which contains an ATP/GTP-binding P-loop. FLJ20051 orthologs are *R. norvegicus* LOC310106, (86.4% similarity), and *M. musculus* C230032K13Rik, (85.1% similarity).

Homology/Unigene data indicates that FLJ20051 acts as an ADP ribosylase. Domain mapping revealed that the 204-amino-acid protein (SwissProt designation Q9NXUS) contains GTP binding site from approximately residues 8-198. Pfam predicts an ADP-ribosylation factor (arf) domain from approximately residues 17 to 194 while SMART analysis predicts the GTPase region lies from amino acids 8-198. Small ADP ribosylation factor GTP-binding proteins are key regulators of vesicle biogenesis in intracellular trafficking. Members of the Arf family include Arl (Arf-like), Arp (Arf-related proteins) and the related Sar (Secretion-associated and Ras-related) proteins. Arf proteins cycle between inactive GDP-bound and active GTP-bound forms that bind selectively to effectors. The Arf domains can form complexes with GMP-PDE delta domains, which are believed to be specific soluble transport factors for certain prenylated proteins; Arl2-GTP is a regulator of PDE-mediated transport. Arf6 is found to be involved in cell motility, membrane dynamics, actin reorganization and tumor cell invasion.

Based on the foregoing information, FLJ20051 is expected to be an ADP Ribosylase active in lymphatic endothelial cells and to regulate GDP/GTP binding and intracellular trafficking, membrane dynamics, cell motility, or actin reorganization. FLJ20051 has recently been named ARFRP2: ADP-ribosylation factor related protein 2 (NM_019087).

FLJ13910: Microarray analysis of LEC-specific genes identified hypothetical protein FLJ13910 as having greater expression levels in LEC versus BEC. This gene was also detected in a putative LEC library. Further, the gene is expressed in lymphatic endothelial cells (LECs), but not in blood vascular endothelial cells (BECs) in an *in vitro* cell culture of primary endothelial cells (HDMEC, PromoCell), confirming that FLJ13910 is a LEC-specific gene.

Protein FLJ13910 (Genbank Accession No. NM_022780; SEQ ID NOS: 17 and 18) is encoded by an mRNA sequence of 3239 base pairs. Unigene analysis of the amino acid sequence of this mRNA indicates that FLJ13910 shows identity to the mouse erythroblast macrophage protein (26% identity over a 261-aa sequence), shows 37% identity (over 320 aa) to a putative *A. thaliana* WD repeat protein; and demonstrates 25% identity (within a 340-aa sequence) to *C. elegans*

T07D1.2.p. Species orthologs of FLJ13910 are found in *R. norvegicus* LOC312439 (94.5 %), and in *M. musculus* 1110007A06Rik (83.8 %).

Protein domain analysis of the 391-residue FLJ13910 protein (SwissProt designation Q9H871) revealed a LisH domain from amino acids 114-146 and a CTLH domain from approximately amino acids 153-210. The lissencephaly type-1-like homology motif (LisH) is an alpha-helical motif present in Lis1, treacle, Nopp140, some katanin p60 subunits, muskelin, tonneau, LEUNIG and WD40 repeat-containing proteins. It is believed that LisH motifs contribute to the regulation of microtubule dynamics, either by mediating dimerization, or by binding either a cytoplasmic dynein heavy chain or microtubules. The CTLH domain forms an alpha-helix. This domain is found in the mouse and human erythroblast macrophage proteins, both mouse and rat muskelin proteins, in the *A. thaliana* putative WD repeat protein. Muskelin is a mediator of cell-adhesive behavior and cytoskeletal organization (Adams *et al.*, *EMBO J.* 17:4964-74, 1998), whereas WD-repeat proteins are a group of structurally related proteins that participate in a wide range of cellular functions, including transmembrane signaling, mRNA modification, vesicle formation, and vesicular trafficking. Protein domain mapping predicts an E3 ubiquitin ligase domain (KOG2817), which is indicative of post-translational modification, protein turnover or chaperone function.

Based on the protein domain mapping analysis, which indicates structural similarity to several actin binding or cytoskeleton related proteins, FLJ13910 is expected to act as a modulator of LEC migration and localization. FLJ13910 is also expected to act as an internal regulator of vesicle formation and vesicle trafficking, based on structural similarity to WD domain-containing proteins. Given the similarity to the muskelin protein, FLJ13910 is also expected to participate in structural and adhesion activities in LECs and plays a role in development and migration of LECs.

In vivo, NR-ISH of frozen sections from E16.5 mouse embryos detected gene transcript in the trigeminal ganglion, brain, nasal cavities, intestine, lung, submandibular gland and kidney.

FLJ11029: Protein FLJ11029 (Genbank Accession No. NM_018304; SEQ ID NOS: 33 and 34) is encoded by a 2334 bp mRNA sequence. Unigene

analysis of the amino acid sequence indicates that FLJ11029, which is upregulated in LEC compared to BEC, is a protein that shows no sequence similarity to any gene disclosed in a public database. Database exploration for species orthologs indicates that FLJ11029 is similar to *R. norvegicus* - LOC287597 (81.8 %), and *M. musculus* B930067F20Rik (81.6 %).

Protein domain mapping of the predicted 360 amino acid FLJ11029 (SwissProt designation Q9NUZ7) indicates that the amino acid sequence contains a domain at approximately residues 1-150 that is similar to domains associated with the PmbA_TldD family of proteins. It has been suggested that the TldD and PmbA proteins modulate the activity of DNA gyrase, or that PmbA may be involved in secretion. Pfam mapping also indicates FLJ11029 amino acids 151-360 are associated with domains found in multiple species of bacteria, the Bac surface Antigen, and are related to the UPF0140 family of eukaryotic proteins. Additional analysis indicates that this protein contains one of the following domains: Zinc finger, C3HC4 type (RING finger), ZPR1 zinc-finger domain, or C5HC2 zinc finger, which is expected to confer a DNA-binding function.

Protein domain mapping leads to the expectation that protein FLJ11029 acts as a DNA-binding/DNA-regulating protein involved in LEC transcriptional regulation.

KIAA1102: KIAA1102 (XM_044461; SEQ ID NO: 35 and 36) is encoded by a 6743 bp mRNA sequence. The gene is expressed in lymphatic endothelial cells (LECs), but not in blood vascular endothelial cells (BECs) in an *in vitro* cell culture of primary endothelial cells (HDMEC, PromoCell).

KIAA1102 is expressed as a 6.5 kb transcript in most tissues and as a 8.5 kb transcript in skeletal muscle. Expression in human tissue, as assayed by Northern blot analysis using Multiple Tissue Northern Blot (Clontech), is highest in heart, brain, skeletal muscle, and lung, with intermediate levels of expression found in prostate, thyroid, spleen, and placenta, and lower levels in kidney, small intestine, tongue and trachea. Assessment of KIAA1102 homologs in other species identified orthologs in *B. taurus* Bt.10388 (87.1%), *R. norvegicus* LOC305332 (84.0 %), and *M. musculus* LOC269654 (81.6%).

Analysis of protein similarity using the Unigene database reveals that KIAA1102 is 28% identical (over 308 amino acids) to human LIM domain only 7 isoform; is 23% identical (over a sequence of 197 aa) to mouse sciellin; is 24% identical (over 293 aa) to *C. elegans* F28F5.3b.p; and is 19% identical (over a
5 sequence of 1371 aa) to the *D. melanogaster* T13564 microtubule-associated protein homolog.

Protein domain mapping of a 1101 fragment (SEQ ID NO. 45) of the predicted 1427 residue KIAA1102 protein (SwissProt designation Q9UPQ0) reveals similarity to a Calponin-like actin-binding domain (CH domain) from approximately
10 residues 41-156; more particularly, Pfam analysis predicts the domain lies at residues 41-156 and SMART predicts a CH domain from amino acids 41-142. Mapping also detected a Zn-binding protein domain (LIM) spanning approximately amino acids 1030-1094 of SEQ ID NO: 45, and more specifically with Pfam analysis from aa 1031-1094, or with SMART analysis from residues 1030-1088.

15 The LIM domain is a conserved cysteine-rich domain of about 60 amino acids. This type of zinc-binding domain is present in diverse proteins, such as the homeobox protein Lim-1, the insulin gene enhancer binding protein Isl-1, and mammalian LH-2, a transcriptional regulatory protein involved in the control of cell differentiation in developing lymphoid and neural cell types. Some LIM domains
20 bind protein partners via tyrosine-containing motifs. LIM domains are found in many key regulators of developmental pathways.

The calponin homology domain defines a superfamily of actin-binding domains found in both cytoskeletal proteins and signal transduction proteins, and is similar to a region in many actin-binding proteins, including spectrin, alpha-actinin
25 and fimbrin, which contain a 250-amino-acid stretch called the actin binding domain(ABD). The ABD is believed to have arisen from duplication of a domain that is also found in as a single copy in a number of other proteins, like calponin or the vav proto-oncogene, and has been called the calponin homology (CH) domain. CH-domain-containing proteins are divided into monomeric actin cross-linking molecules,
30 dimeric cross-linking proteins (e.g., alpha-actinin, beta-spectrin, and filamin) and monomeric F-actin binding proteins (dystrophin, utrophin). Each single ABD, comprising two CH domains, is able to bind one actin monomer. The CH domain

also occurs in a number of proteins not known to bind actin, a notable example being the vav-protooncogene. Examples of proteins containing CH domains include alpha-actinins, calponins, human dystrophin, mammalian vav proto-oncogene, human ras GTPase activating-like protein IQGAP1, Spectrin beta chain., Transgelins, and T-plastin.

Given the protein domain homology data above, it is expected that KIAA1102 functions as a DNA control protein using its Zinc binding domain and this protein may regulate development of LECs. Additionally, the KIAA1102 protein comprises a domain similar to those in actin-binding proteins and is expected to act as a regulator of cell motility.

In vivo, non-radioactive *in situ* hybridization (NR-ISH) of frozen sections from E16.5 mouse embryos showed KIAA1102 gene expression in lung, kidney glomerulus, thymus, brain, lips and bone marrow. *FLJ32029*: The protein-coding region of protein FLJ32029 (phosphoglucomutase 2-like 1 (PGM2L1), NM_173582; SEQ ID NOS: 37 and 38) lies within a 8456 bp mRNA sequence. The FLJ32029 mRNA is expressed as a 9.5 kb mRNA that is detected in brain and skeletal muscle. Comparison of the sequence to other mammalian species identified orthologs in *R. norvegicus* LOC293145 (84.9% similar) and *M. musculus* 4931406N15Rik (83.9%).

Unigene analysis revealed that FLJ32029 is 59% identical (over 596 aa) to human protein FLJ10983; 44% identical (over 600 aa) to a *C.elegans* phosphoglucomutase and phosphomannomutase phosphoserine; and 39% identical (over 576 aa) to *S. cerevisiae* protein YMR278w.

Mapping for potential protein family domain similarities indicates the predicted 622-amino-acid FLJ32029 protein (SwissProt designation Q96MQ7) comprises multiple phosphoglucomutase/phosphomannomutase domains (PGM-PMM-I and PGM-PMM II) at approximately amino acid residues 61-217 (alpha/beta/alpha domain I); amino acids 237-349 (alpha/beta/alpha domain II); amino acids 428-519 (alpha/beta/alpha domain III); and amino acids 520-617, similar to the phosphomannomutase C-terminal domain, which is involved in intramolecular transferase activity (e.g., phosphotransferase activity) and carbohydrate transport and metabolism. FLJ32029 is expected to function in carbohydrate regulation in

lymphatic endothelial cells and a dysfunctional FLJ32029 gene is expected to be involved in the development of lymphedema as a result of improper carbohydrate metabolism (Drubaix et al., *Int J Exp Pathol.* 78:117-21. 1997).

The identity of FLJ32029 as an LEC-specific gene was confirmed in
5 an *in vitro* culture, where the gene was detected in lymphatic endothelial cells (LECs), and to a lesser extent in blood vascular endothelial cells (BECs) in cell culture of primary endothelial cells (HDMEC, PromoCell).

In vivo analysis of frozen sections from E16.5 mouse embryos (NR-ISH), detected gene expression in brain, bone marrow, glomeruli of the kidney cortex,
10 thymus and intestine. *FLJ23594*: The protein-coding region of FLJ23594 (C18orf14, chromosome 18 open reading frame 14, NM_024781; SEQ ID NOS: 39 and 40) lies within a 2015 bp mRNA sequence. The 3 kb FLJ23594 transcript has been found to be expressed in human tissue at high levels in the placenta, lung, lymph node, and adrenal gland, as detected using a Multiple Tissue Northern Blot (Clontech) of adult
15 human RNA. *In vitro* analysis shows that FLJ23594 was detected in lymphatic endothelial cells (LECs) and, to a lesser extent, in blood vascular endothelial cells (BECs) in cell culture of primary endothelial cells (HDMEC, PromoCell).

Comparison of the amino acid sequence with known amino acid sequences reveals that FLJ23594 shows 23% identity (over 271 aa) to human myosin
20 heavy chain, nonmuscle type B (Cellular myosin heavy chain, type B); demonstrates 22% identity (over 271 aa) to mouse myosin heavy chain, nonmuscle type B; is 22% identical (over a 271-aa sequence) to rat myosin heavy chain, nonmuscle type B; shows 24% identity (over 300 aa) to *D. melanogaster* myosin heavy chain 2; shows 23% identity (over 266 aa) to *S. cerevisiae* myosin-1 isoform (Type II myosin); shows
25 22% identity (over 269 aa) to an *A. thaliana* probable centromere protein; and is 22% identical (over a sequence of 284 aa) to *C. elegans* protein F52B10.1.

Protein domain mapping of FLJ23594 (SwissProt designation Q9H5C1) reveals a protein of 297 amino acids having two regions of similarity to bacterial M protein, from amino acids 148-168 and 216-236. Mapping also indicates
30 regions of similarity, at aa 153-296, to the bZIP leucine zipper. The bZIP superfamily of eukaryotic DNA-binding transcription factors groups together proteins that contain a basic region mediating sequence-specific DNA-binding followed by a leucine zipper

required for dimerization. bZIP domains usually bind a pallindromic six-nucleotide site, but the specificity can be altered by interaction with an accessory factor. Some proteins that contain the bZIP domain include AP-1, oncogene v-jun, Jun-B and jun-D, probable transcription factors which are highly similar to jun/AP-1, and
5 mammalian cAMP response element (CRE) binding proteins CREB, CREM, ATF-1, ATF-3, ATF-4, ATF-5, ATF-6 and LRF-1.

The similarity to bZIP-containing proteins indicates FLJ23594 acts as a DNA-binding protein and functions in the regulation of DNA transcription. The ability to regulate LEC-specific DNA-binding proteins or transcription factors allows
10 for manipulation of LEC development, wherein the LEC can be induced to grow and proliferate by increased transcription or where the LEC can be induced to halt growth by blocking DNA regulatory functions. The structural similarity to cellular myosin also implicates FLJ23594 in cytokinesis, cell shaping, and specialized functions such as secretion and capping.

15 *FLJ10532*: Protein FLJ10532 (Genbank Accession No. AK001394; SEQ ID NO: 41) is encoded by a nucleic acid that was isolated as a 2372 bp mRNA fragment from *fis*, clone NT2RP2001044. The 2.3 kb transcript may be part of the message encoding protein DKFZp762K222. Cross-species comparison identified possible orthologs for DKFZp762K222 as *R. norvegicus* LOC306459 (88.3%), and
20 *M. musculus* 4933409N07Rik (87.7 %). Cell culture of primary endothelial cells (HDMEC, PromoCell) indicates the FLJ10532 transcript is expressed in lymphatic endothelial cells (LECs) and is undetectable blood vascular endothelial cells (BECs).

BIKE: Identified herein as a LEC-specific gene, the human BIKE gene (Genbank Accession No. NM_017593; SEQ ID NOS: 25 and 26), a homolog of
25 mouse BMP-2 kinase, exhibits a 2763 bp nucleotide sequence expressed as a 7.5 kb transcript. Northern blot analysis indicates that BIKE is expressed at high levels in skeletal muscle, tongue, lymph node, colon, and placenta, and at lower levels in heart, thyroid and stomach. BIKE demonstrates 93% identity, over a region of 329 aligned amino acids, to mouse Bmp2-inducible kinase; 39% identity, over a 289-residue
30 alignment, to rat cyclin G-associated kinase; 40% identity, over a 270-aa alignment, to *Arabidopsis thaliana* hypothetical protein T21L14.21; 65% identity, over a 289-aa

region, to *D. melanogaster* Nak-P1; and 41% identity, over a 291-residue alignment, to a *C. elegans* serine/threonine kinase.

Protein domain mapping by Pfam analysis of a 661 residue human polypeptide sequence (SEQ ID NO: 46) comprising the predicted 355 amino acid
5 BIKE protein [SwissProt designation Q9NSY1 (human), Q91Z96 (Mouse, 1138 residue fragment)] indicates the protein contains a protein kinase domain located from approximately residues 51 to 314. SMART domain predicts this domain lies from approximately residues 51-317. Thus, the protein kinase domain lies at approximately amino acids 51-317. SMART analysis indicates the domain is a
10 STYKc domain, as in KIAA0472, which are expected to be phosphotransferases that exhibit dual-specificity as Ser/Thr/Tyr kinases.

The BIKE gene product is the human homolog of mouse BMP-2-inducible kinase. Bone morphogenic proteins (BMPs) play a key role in skeletal development and patterning. Expression of the mouse gene is increased during BMP-
15 2-induced differentiation and the gene product is a putative serine/threonine protein kinase containing a nuclear localization signal. Therefore, the protein encoded by the human homolog is expected to be a protein kinase with a putative regulatory role in osteoblast differentiation. The unexpected identification of the BIKE protein as a LEC-specific gene reveals that the BMP homolog is involved in development of the
20 lymphatic system, as well as osteogenesis.

The manipulation of lymphatic endothelial-cell-specific molecules is expected to be applicable to treatments of LEC diseases disorders associated with tissue edemas. Without wishing to be bound by theory, manipulation of such molecules is expected to modulate endothelial cell-cell or cell-matrix protein
25 interactions and/or to affect transendothelial transport, thereby altering the state of fluid transport across the lymphatic vessel wall. Further, such molecules provide targets for the delivery of therapeutic compounds, such as growth factors, mitogens, and the like, as well as cytostatic or cytotoxic agents known in the art. These therapeutic compounds are targeted to such cells by associating a therapeutic agent
30 with, e.g., a binding partner (such as an antibody) of the LEC surface marker. The

transmembrane proteins identified herein also provide useful targets for modulating cell adhesion or extracellular events integral to lymphatic endothelial cell function.

EXAMPLE 3

5 **EX-VIVO CELL STIMULATION AND GENE THERAPY FOR LYMPHEDEMA WITH LEC-SPECIFIC GENE-TRANSFECTED CELLS**

The ability of LEC-specific transcription factors to regulate genes specifically involved in LEC development provides a basis for treatment of individuals exhibiting a LEC disorder or condition resulting from either an increase or
10 decrease in LEC gene expression levels. For example, mouse pups with a homozygous mutation in transcription factor Sox18 that disrupts formation of the Sox-18/MEF2C complex (Hosking, *et al.*, *Biochem. Biophys. Res. Commun.* 287:493-500. 2001) develop chylous ascites in some genetic backgrounds (Pennisi, *et al.*, *Nat. Genet.* 24:434-437. 2000), suggesting that both proteins may be involved in the
15 regulation of lymphatic development. Additionally, upregulation of VEGFR-3 expression by transcription factor Prox-1 has been shown to be involved in development of LEC (See PCT/US03/06900 and Petrova *et al.*, *EMBO J.* 21:4593-4599, 2002).

Similar to the effects of Prox-1, it is contemplated that upregulation of
20 the LEC-specific genes disclosed herein is useful in promoting LEC development as a treatment for LEC disorders characterized by an under-developed lymphatic system. Conversely, decreasing activity of LEC-specific genes and downregulating LEC development is contemplated as a treatment for LEC disorders characterized by an over-developed lymphatic system such as lymphedema. Additionally, use of the
25 genes or gene products disclosed herein as prophylactic therapeutics designed to serve as a therapeutic antagonist in therapies involving other primary agents is also contemplated, such as the use of a material disclosed herein to counter the influence of the primary agent on LEC gene expression. It is known in the art that *ex vivo* transfection of cells and subsequent transfer of these cells to patients is an effective
30 method to upregulate *in vivo* levels of the transferred gene and to provide relief from a

disease resulting from under-expression of the gene(s) (Gelse et al., *Arthritis Rheum.* 48:430-41. 2003; Huard et al, *Gene Ther.* 9:1617-26. 2002; Kim et al., *Mol. Ther.* 6:591-600 2002).

To develop a therapy for treating irregularities of LEC development, endothelial cells, such as CAECs, SAVECs, LECs or BECs, are isolated from individuals experiencing a LEC disorder (e.g. lymphedema) and then placed in an appropriate culture medium (see above) to promote the growth and viability of the cells. The cells are then transfected with a viral vector designed to express any one of the LEC-specific genes to initiate LEC differentiation of the non-LECs *in vitro* and to promote growth of the LECs in culture. Transfection of cells with a viral vector, such as a replication-deficient adenovirus, adeno-associated virus, lentivirus or retrovirus, is well known to those of skill in the art (see, e.g., Laitinen et al., *Hum. Gene Ther.* 9:1481-1486, 1998; Petrova et al., *supra*) and is described above.

Cells transfected with a LEC-specific gene are then transferred into an affected patient in therapeutically effective numbers to promote LEC expansion *in vivo*. In preferred embodiments, the manipulated cells are autologous cells. These cells are delivered by one or more administrations typically involving injection. The cells are delivered at a local site of an LEC disease or disorder such as lymphedema; alternatively, the cells are delivered systemically, e.g., by injection remote to the perceived site in need of such cells or their product(s).

Administration of transfected cells to patients with lymphedema provides supplementary LECs that are incorporated into the lymphatic network, thereby promoting lymphatic development and lymph clearance to relieve the symptoms of lymphedema. It is contemplated that a method comprising LEC-specific gene transfection into endothelial cells and administration of transfected cells is useful in the treatment of any disease characterized by an alteration in LEC numbers or activity, such as lymphedema, lymphangioma, lymphangiomyeloma, lymphangiomatosis, lymphangiectasis, lymphosarcoma, and lymphangiosclerosis. Additionally, such methods are useful in ameliorating a symptom (e.g., lymph-induced swelling in the case of lymphedema) associated with such diseases.

EXAMPLE 4
MICROARRAY ANALYSIS TO DETECT LEC-AND LYMPH-RELATED DISORDERS

All LEC-specific genes identified, specifically any gene found in any of Tables 1 and 2 herein, are contemplated to be useful in the detection of LEC *in vivo* and in determining the extent of the lymphatic vasculature in a sample. The LEC-specific genes are also expected to be useful in diagnosing lymphedema and other LEC-related disorders. Further, the invention comprehends the use of a plurality of LEC-specific genes or their gene products for any of the applications identified herein. In this latter context, it is further contemplated that one or more LEC-specific genes, or their gene products, may be used in combination (regardless of whether such materials are physically or temporally combined or not) with at least one of the LEC-specific genes, or gene products, identified in Tables 3 and/or 4 of PCT/US03/06900.

Another aspect of the invention is a composition comprising a plurality of polynucleotide probes for use in detecting gene expression pattern(s) characteristic of particular cell type(s) and for detecting changes in the expression pattern of a particular cell type, *e.g.*, lymphatic endothelial cells. The term "polynucleotide probe" is used herein to refer to any one of the nucleic acid sequences listed in SEQ ID NOS: 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 and 41, or any fragment thereof or a nucleic acid having a sequence encoding an amino acid sequence set forth in SEQ ID NOS: 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 43, 44, 45, and 46, or a fragment thereof. Preferably, the fragment is at least 10 nucleotides in length; more preferably, it is at least 20 nucleotides in length. Such a composition is employed for the diagnosis and/or treatment of any condition or disease in which the dysfunction or non-function of lymphatic endothelial cells is implicated or suspected. The composition may alternatively include any one of genes found in Tables 3 and 4 of PCT/US03/06900 (specifically incorporated herein by reference). In one embodiment, the present invention provides a composition comprising a plurality of polynucleotide probes, wherein at least a subset of the polynucleotide probes comprises at least a portion of an expressed gene isolated from a population of LEC-specific genes identified above. Also contemplated is a composition comprising a plurality of polynucleotide probes, with

at least a subset of such probes each comprising a unique sequence selected from the group consisting of SEQ ID NOS: 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, and those sequences identified in any one of Tables 3 and 4 of PCT/US03/06900. Preferably, the composition comprises a subset of at least 3
5 polynucleotides, each having a different sequence selected from the group consisting of SEQ ID NOS: 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, and those sequences identified in any one of Tables 3 and 4 of PCT/US03/06900. Also preferred are compositions comprising at least 5, at least 7, at least 9, at least 15, at least 20, or at least 25 distinct polynucleotides having distinct sequences selected
10 from the group of SEQ ID NOS: 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, and those identified in any one of Tables 3 and 4 of PCT/US03/06900.

The composition is particularly useful as a set of hybridizable array elements in a microarray for monitoring the expression of a plurality of target polynucleotides. The microarray comprises a substrate and the hybridizable array
15 elements. The microarray is used, for example, in the diagnosis and/or prognosis of a disease derived from aberrant lymphatic endothelial cell activity, such as lymphedema, lymphangioma, lymphangiomyeloma, lymphangiomatosis, lymphangiectasis, lymphosarcoma, and lymphangiosclerosis. Compositions may be useful in identifying more than one cell type and may be useful in the diagnosis
20 and/or prognosis of more than one disease, disorder or condition. Further, useful information is obtained from those probes yielding a signal and from those probes not yielding a signal.

A polynucleotide comprising the sequence of any one of SEQ ID NOS: 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, and any one of the
25 genes found in any of Tables 3 and 4 of PCT/US03/06900 may be used for the diagnosis of a condition, disorder, or disease in which the abnormal expression of any one of the genes encoded by SEQ ID NOS: 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, and any gene found in any one of Tables 3 and 4 of PCT/US03/06900, is associated. For example, a polynucleotide comprising any one
30 of the sequences set forth in SEQ ID NOS: 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, and a gene found in any of Tables 3 and 4 of

PCT/US03/06900 may be used in hybridization or PCR assays of a fluid or tissue (e.g., obtained from a biopsy) to detect abnormal gene expression in patients with lymphedema or another lymph-associated disease, or suspected of having such a disease. In addition, a polynucleotide comprising a sequence encoding any of the amino acid sequences set forth in SEQ ID NOS: 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 43, 44, 45, and 46, used in conjunction with at least one gene found in any of Tables 3 and 4 of PCT/US03/06900 is useful for the diagnosis of conditions or diseases associated with aberrant expression of a polypeptide having any one of those amino acid sequences. Fragments comprising at least 10 nucleotides are also useful in these diagnostic methods.

Expression profiles are generated using the compositions of the invention comprising, e.g., at least one sequence as set forth in SEQ ID NOS: 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 or 41. The expression profile generated from a microarray is used to detect changes in the expression of genes implicated in disease.

EXAMPLE 5

COMPOSITIONS OF THE INVENTION USEFUL FOR MODULATING VEGF-C/VEGFR-3 INTERACTIONS AND LEC DEVELOPMENT

VEGF-C and VEGF-D knockout mice demonstrate aberrant vascular development that can be overcome by administration of exogenous VEGF-C and/or VEGF-D polypeptide. To determine if transcriptional regulation by any one of the proteins having an amino acid sequence set forth in SEQ ID NOS: 18, 26, 28, 30, 32, 34, 36, 38 or 40 (which may act as LEC-specific transcription factors) can overcome this vascular defect due to its potential interaction with, and transcriptional effect on, the VEGFR-3 promoter, VEGF-C or VEGF-D knockout mice are genetically crossed by interbreeding with mice overexpressing a polypeptide having an amino acid sequence of any one of SEQ ID NOS: 18, 26, 28, 30, 32, 34, 36, 38, 40, 44, 45 or 46 from a cell-specific-promoter (e.g., a K-14 keratin promoter). The effects of transcription factor (TF) activity on lymphedema are assessed through measurement of lymphedema and vascular development, as described in Karkkainen et al. (*Proc. Natl. Acad. Sci. USA*. 98:12677-12682. 2001).

Survival of the knockout mice and detection of lymphatic development in the VEGF-C and/or VEGF-D knockout/TF-overexpressing mice indicates that the transcription factor of interest induces VEGFR-3 signaling and plays a role in lymphangiogenesis.

5 VEGF-C overexpressing mice (K-14-VEGF-C Tg) exhibit an extensive network of lymphatic vasculature, are prone to tumor metastasis, and demonstrate upregulated VEGFR-3 expression and symptoms of lymphedema (US Patent No. 6,361,946). To determine if the LEC-specific genes identified herein aid in regulating VEGF-C signaling through VEGFR-3, K-14-VEGF-C Tg mice are
10 administered a composition of the invention, comprising at least one LEC-specific polypeptide or polypeptide fragment. Particularly useful are those fragments from extracellular proteins having sequences set forth in SEQ ID NOS: 6, 8, 10, 12, 14, 16, 20, 22, 24, 42 and 43.

Decreased lymphangiogenesis, decreased incidence of tumor
15 metastasis, and decreased levels of VEGFR-3 exhibited by the LEC-specific-composition-treated K-14-VEGF-C mutant animals, as compared to the K-14-VEGF-C Tg single mutant animal, indicates that the administered composition interferes with VEGF-C signaling through VEGFR-3.

A result indicating that the composition of the invention is a negative
20 regulator of lymphangiogenesis identifies a composition useful in a method of treating disorders mediated by extensive lymphatic vasculature, such as lymphangiogenesis in tumor development or lymphangiosarcoma. Such a composition, comprising at least one LEC-specific polypeptide, may be administered to a patient or an animal in need, thereby inhibiting or preventing the induction of lymphangiogenesis by interfering
25 with lymphangiogenic signaling. A result indicating that a composition of the invention is a positive regulator of lymphatic endothelial cells identifies a composition suitable for use in a method of treating disorders characterized by insufficient LEC development, e.g., insufficient development of a lymphatic network.

Additionally, to determine the effects of the invention *in vivo*,
30 compositions of the invention are administered to animals affected by lymphedema in the *Chy* mouse model of lymphedema (Lyon *et al.*, *Mouse News Lett.* 71: 26, 1984). Lymphedema symptoms, such as footpad swelling and fluid retention/ascites, are

measured after treatment with either a soluble LEC-specific polypeptide, as described above, or with a LEC-specific gene therapy vector designed to regulate either cell surface proteins or intracellular proteins, comprising at least one of SEQ ID NOS: 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 or 41, as disclosed
5 previously. Improvement of lymphedema in the *Chy* mouse model after administration of compositions of the invention indicate that a composition which affects LEC-specific genes is capable of regulating LEC disorders, and is expected to be a useful therapeutic in the treatment of human lymphatic disorders.

Additionally contemplated is the treatment of subjects affected by
10 lymphedema with a combination of therapeutics that includes a composition of the invention in conjunction with a lymphatic growth factor such as VEGF-C and/or VEGF-D to overcome impaired lymphatic vascular development.

From the foregoing it will be appreciated that, although specific
embodiments of the invention have been described herein for purposes of illustration,
15 various modifications may be made without deviating from the spirit and scope of the invention.

All of the above-cited U.S. patents, U.S. patent application
publications, U.S. patent applications, foreign patents, foreign patent applications and
non-patent publications referred to in this specification are incorporated herein by
20 reference in their entireties.

CLAIMS

We claim:

1. A method for differentially modulating the growth or differentiation of blood endothelial cells (BEC) or lymphatic endothelial cells (LEC), comprising contacting endothelial cells with a composition comprising an agent that differentially modulates blood or lymphatic endothelial cells, said agent selected from the group consisting of:

- (a) a polypeptide that comprises an amino acid sequence of a BEC polypeptide or a LEC polypeptide, or an active fragment of said polypeptide ;
- (b) a polynucleotide that comprises a nucleotide sequence that encodes a polypeptide according to (a);
- (c) an antibody that specifically binds to a polypeptide according to (a);
- (d) a polypeptide comprising a fragment of the antibody of (c), wherein the fragment and the antibody bind to the polypeptide;
- (e) an antisense nucleic acid to a human gene or mRNA encoding the polypeptide of (a);
- (f) an interfering RNA (RNAi) to a human gene or mRNA encoding the polypeptide of (a).

2. A method according to claim 1, wherein the endothelial cells are contacted with the composition *ex vivo*.

3. A method according to claim 1, wherein the composition comprises a pharmaceutically acceptable diluent, adjuvant, or carrier, and the contacting step comprises administering the composition to a mammalian subject to differentially modulate BECs or LECs in the mammalian subject.

4. A method according to claim 3, comprising:
identifying a human subject with a disorder characterized by hyperproliferation of LECs; and
administering to the human subject the composition, wherein the agent differentially inhibits LEC growth compared to BEC growth.

5. A method according to claim 3, comprising:
identifying a human subject with a disorder characterized by hyperproliferation of LECs;
screening LECs of the subject to identify overexpression of a polypeptide set forth in any of Tables 1 and 2 herein; and
administering to the human subject the composition, wherein the agent differentially inhibits LEC growth compared to BEC growth by inhibiting expression of the polypeptide identified by the screening step.

6. A method according to claim 3 of modulating the growth of lymphatic endothelial cells in a human subject, comprising steps of:
identifying a human subject with a hypoproliferative lymphatic disorder;
screening the subject to identify underexpression or underactivity of a LEC polypeptide set forth in any of Tables 1 and 2 herein;
administering to the human subject said composition, wherein the agent comprises the LEC polypeptide (a) identified by the screening step or an active fragment of said polypeptide, or comprises the polynucleotide (b) that comprises a nucleotide sequence that encodes the polypeptide.

7. Use of an agent for the manufacture of a medicament for the differential modulation of blood vessel endothelial cell (BEC) or lymphatic vessel endothelial cell (LEC) growth or differentiation, said agent selected from the group consisting of:

- (a) a polypeptide that comprises an amino acid sequence of a BEC polypeptide or a LEC polypeptide, or an active fragment of said polypeptide ;
- (b) a polynucleotide that comprises a nucleotide sequence that encodes a polypeptide according to (a);
- (c) an antibody that specifically binds to a polypeptide according to (a);
- (d) a polypeptide comprising a fragment of the antibody of (c), wherein the fragment and the antibody bind to the polypeptide;
- (e) an antisense nucleic acid to a human gene or mRNA encoding the polypeptide of (a);
- (f) an interfering RNA (RNAi) to a human gene or mRNA encoding the polypeptide of (a).

8. A method or use according to any one of claims 1-7, wherein the polypeptide is a LEC polypeptide selected from the LEC polypeptides set forth in any of Tables 1 and 2 herein, and the agent differentially modulates LEC growth or differentiation over BEC growth or differentiation..

9. A method or use according to any one of claims 1-7, wherein the polypeptide is a BEC polypeptide selected from the BEC polypeptides set forth in Tables 1 and 2 herein, and the agent differentially modulates BEC growth or differentiation over LEC growth or differentiation.

10. The method according to any of claims 5 or 6, further comprising screening LECs of the subject to identify overexpression of a second polypeptide set forth in any of Tables 3 and 4 of PCT/US03/06900.

11. A method or use according to claim 8, wherein the LEC polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 6, 8, 10, 12, 14, 16, 20, 22, 24, 42 and 43.
12. A method or use according to claim 8, wherein the LEC polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 20.
13. A method or use according to claim 7, wherein the agent comprises an antibody according to (c) or polypeptide according to (d).
14. A method according to claim 7, wherein the agent comprises an extracellular domain fragment of the polypeptide of (a), or a polynucleotide encoding said extracellular domain fragment.
15. A method or use according to any one of claims 1-10, wherein the agent comprises an antisense molecule.
16. A method of treating hereditary lymphedema comprising:
identifying a human subject with lymphedema and with a mutation in at least one allele of a gene encoding a LEC protein identified in any of Tables 1 and 2, wherein the mutation correlates with lymphedema in human subjects, and with the proviso that said LEC protein is not VEGFR-3; and
administering to said subject a composition comprising a lymphatic growth agent selected from the group consisting of VEGF-C polypeptides, VEGF-D polypeptides, VEGF-C polynucleotides, and VEGF-D polynucleotides.

17. Use of a lymphatic growth agent selected from the group consisting of VEGF-C polypeptides, VEGF-D polypeptides, VEGF-C polynucleotides, and VEGF-D polynucleotides in the manufacture of a medicament for the treatment of hereditary lymphedema resulting from a mutation in a LEC gene identified in any of Tables 1 and 2.

18. A method of screening for an endothelial cell disorder or predisposition to said disorder, comprising:
obtaining a biological sample containing endothelial cell mRNA from a human subject; and
measuring expression of a BEC or LEC gene from the amount of mRNA in the sample transcribed from said gene, wherein the BEC or LEC gene encodes a polypeptide identified in any of Tables 1 and 2.

19. A method of monitoring the efficacy or toxicity of a drug on endothelial cells, comprising the steps of:
measuring expression of at least one BEC or LEC gene in endothelial cells of a mammalian subject before and after administering a drug to the subject, wherein the at least one BEC or LEC gene encodes a polypeptide set forth in any of Tables 1 and 2, and wherein changes in expression of the BEC or LEC gene correlates with efficacy or toxicity of the drug on endothelial cells.

20. A method of identifying compounds that modulate growth of endothelial cells, comprising
culturing endothelial cells in the presence and absence of a compound;
and
measuring expression of at least one BEC or LEC gene in the cells, wherein the BEC or LEC gene is selected from the genes encoding polypeptides set forth in any of Tables 1 and 2, wherein a change in expression of at least one BEC gene in the presence compared to the absence of the compound identifies the compound as a modulator of BEC growth, and wherein a change in expression of at

least one LEC gene in the presence compared to the absence of the compound identifies the compound as a modulator of LEC growth.

21. A method according to claim 20 of screening for a compound that selectively modulates BEC or LEC growth or differentiation, wherein the measuring step comprises measuring expression of at least one BEC gene and at least one LEC gene in the cells, and wherein the method comprises screening for a compound that selectively modulates BEC or LEC growth or differentiation by selecting a compound that differentially modulates expression of the at least one BEC gene compared to expression of the at least one LEC gene.

22. A composition comprising an isolated polynucleotide that comprises a nucleotide sequence that encodes a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 43, 44, 45, and 46; and a pharmaceutically acceptable diluent, carrier or adjuvant.

23. A composition according to claim 22, comprising a polynucleotide that comprises a nucleotide sequence selected from the group consisting of SEQ ID NOS: 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 and 41, or a fragment thereof that encodes the polypeptide.

24. An expression vector comprising an expression control sequence operably linked to a polynucleotide that comprises a nucleotide sequence that encodes a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 43, 44, 45, and 46.

25. An expression vector according to claim 24 that is a replication-deficient adenoviral or adeno-associated viral vector containing the polynucleotide.
26. A composition comprising an expression vector according to claim 24 or claim 25 and a pharmaceutically acceptable diluent, carrier, or adjuvant.
27. A kit comprising the composition according to any one of claims 22, 23, or 26 packaged with a protocol for administering the composition to a mammalian subject to modulate the lymphatic system in said subject.
28. A host cell transformed or transfected with an expression vector according to claim 24.
29. A method for producing a LEC polypeptide comprising the steps of growing a host cell according to claim 28 under conditions in which the cell expresses the polypeptide encoded by the polynucleotide.
30. A purified and isolated polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 43, 44, 45, and 46.
31. A purified and isolated polypeptide comprising an amino acid sequence selected from the group consisting of:
- (a) SEQ ID NOS: 6, 8, 10, 12, 14, 16, 20, 22, 24, 42 and 43; and
 - (b) an extracellular domain fragment of at least 10 amino acids of an amino acid sequence of (a).

32. A purified and isolated, soluble polypeptide according to claim 31 comprising an extracellular domain fragment of an amino acid sequence selected from the group consisting of : SEQ ID NOS: 6, 8, 10, 12, 14, 16, 20, 22, 24, 42 and 43, wherein the polypeptide lacks any transmembrane domain.

33. A polypeptide according to claim 32 that lacks any intracellular domain.

34. A fusion protein comprising a polypeptide according to claim 32 or 33 fused to an immunoglobulin fragment comprising an immunoglobulin constant region.

35. A composition comprising a polypeptide or protein according to any one of claims 30-34 and a pharmaceutically acceptable diluent, carrier or adjuvant.

36. A kit comprising the composition according to claim 35 and a protocol for administering said pharmaceutical composition to a mammalian subject to modulate the lymphatic system in said subject.

37. An antibody that specifically binds to a polypeptide according to any one of claims 30-34.

38. An antibody according to claim 37 that is a humanized antibody, a chimerized antibody, a human antibody, or a Fab fragment thereof.

39. A protein comprising an antigen binding domain of an antibody that specifically binds to a polypeptide according to any one of claim 30-34, wherein said protein specifically binds to said polypeptide.

40. A method of identifying a LEC nucleic acid comprising:
(a) contacting a biological sample containing a candidate LEC nucleic acid with a polynucleotide comprising a fragment of at least 14 contiguous nucleotides of a sequence selected from the group consisting of SEQ ID NOS: 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 and 41, or a complement thereof, under the following stringent hybridization conditions:

(i) hybridization at 42°C for 20 hours in a solution containing 50% formamide, 5xSSPE, 5x Denhardt's solution, 0.1% SDS and 0.1 mg/ml denatured salmon sperm DNA, and

(ii) washing for 30 minutes at 65°C in 1xSSC, 0.1% SDS; and
(b) detecting hybridization of said candidate LEC nucleic acid and said polynucleotide, thereby identifying a LEC nucleic acid.

41. A method of identifying a LEC protein comprising:
(a) contacting a biological sample containing a candidate LEC protein with a LEC protein binding partner selected from the group consisting of an antibody according to claim 37 or a protein according to claim 39, under conditions suitable for binding therebetween; and

(b) detecting binding between said candidate LEC protein and said LEC binding partner, thereby identifying a LEC protein.

42. A method of identifying a LEC comprising:
(a) contacting a biological sample comprising cells with a LEC binding partner under conditions suitable for binding therebetween, wherein said LEC binding partner comprises an antibody that binds to a polypeptide comprising a sequence selected from the group consisting of SEQ ID NOS: 6, 8, 10, 12, 14, 16, 20, 22, 24, 42 and 43, or comprises an antigen binding fragment of said antibody; and
(b) identifying a LEC by detecting binding between a cell and said LEC binding partner, where binding of the LEC binding partner to the cell identifies a LEC.

43. A method of assaying for risk of developing hereditary lymphedema, comprising

(a) assaying nucleic acid of a human subject for a mutation that correlates with a hereditary lymphedema phenotype and alters the encoded amino acid sequence of at least one gene allele of the human subject when compared to the amino acid sequence of the polypeptide encoded by a corresponding wild-type gene allele, wherein the wild-type polypeptide is a polypeptide identified in any of Tables 1 and 2.

44. A method of assaying for risk of developing hereditary lymphedema, comprising

(a) assaying nucleic acid of a human subject for a mutation that correlates with a hereditary lymphedema phenotype and alters the encoded amino acid sequence of at least one gene allele of the human subject when compared to the amino acid sequence of the polypeptide encoded by a corresponding wild-type gene allele, wherein the wild-type polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NOS: 6, 8, 10, 12, 14, 16, 20, 22, 24, 42 and 43;

(b) correlating the presence or absence of said mutation in the nucleic acid to a risk of developing hereditary lymphedema, wherein the presence of said mutation in the nucleic acid correlates with an increased risk of developing hereditary lymphedema, and wherein the absence of said mutation in the nucleic acid correlates with no increased risk of developing hereditary lymphedema.

45. A method of assaying for risk of developing hereditary lymphedema, comprising

(a) assaying nucleic acid of a human subject for a mutation that alters the encoded amino acid sequence of at least one transcription factor allele of the human subject and alters transcription modulation activity of the transcription factor polypeptide encoded by the allele, when compared to the transcription modulation activity of a transcription factor polypeptide encoded by a wild-type allele,

wherein the wild-type transcription factor polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 18, 26, 28, 30, 32, 34, 36, 38, 40, 44, 45, and 46; and

(b) correlating the presence or absence of said mutation in the nucleic acid to a risk of developing hereditary lymphedema, wherein the presence of said mutation in the nucleic acid correlates with an increased risk of developing hereditary lymphedema, and wherein the absence of said mutation in the nucleic acid correlates with no increased risk of developing hereditary lymphedema.

46. The method according to claim 45 further comprising assaying said nucleic acid for a mutation that alters the expression of Sox18.

47. The method according to claim 46 wherein the assaying identifies a mutation altering activity of a transactivating or DNA binding domain of Sox18.

48. The method according to claim 46, wherein said mutation reduces transcriptional activation of a SOX18-responsive gene compared to transcriptional activation of said gene in the absence of said mutation.

49. A method of assaying for risk of developing hereditary lymphedema, comprising

(a) assaying nucleic acid of a human subject for a mutation that alters the encoded amino acid sequence of at least one LEC gene allele of the human subject and alters the binding affinity of the adhesion polypeptide encoded by the LEC gene allele, when compared to the binding affinity of an adhesion polypeptide encoded by a wild-type allele,

wherein the wild-type adhesion polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 20 and SEQ ID NO: 43; and

(b) correlating the presence or absence of said mutation in the nucleic acid to a risk of developing hereditary lymphedema, wherein the presence of said mutation in the nucleic acid correlates with an increased risk of developing hereditary lymphedema, and wherein the absence of said mutation in the nucleic acid correlates with no increased risk of developing hereditary lymphedema.

50. The method according to any one of claims 43-49, wherein the assaying identifies the presence of the mutation, and the correlating step identifies the increased risk of said patient developing hereditary lymphedema.

51. The method according to claim 49, further comprising assaying said nucleic acid for a mutation that alters the expression of Sox18.

52. A method of screening a human subject for an increased risk of developing hereditary lymphedema comprising assaying nucleic acid of a human subject for a mutation that alters the encoded amino acid sequence of at least one polypeptide comprising an amino acid sequence found in any of Tables 1 and 2.

53. The method according to claim 52, wherein the polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NOS: 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 43, 44, 45, and 46 in a manner that correlates with the risk of developing hereditary lymphedema.

54. The method according to claim 52 further comprising assaying said nucleic acid for a mutation that alters the expression of Sox18.

55. The method according to any one of claims 43-54 wherein said method comprises at least one procedure selected from the group consisting of:

- (a) determining a nucleotide sequence of at least one codon of at least one polynucleotide of the human subject;
- (b) performing a hybridization assay to determine whether nucleic acid from the human subject has a nucleotide sequence identical to or different from one or more reference sequences;
- (c) performing a polynucleotide migration assay to determine whether nucleic acid from the human subject has a nucleotide sequence identical to or different from one or more reference sequences; and
- (d) performing a restriction endonuclease digestion to determine whether nucleic acid from the human subject has a nucleotide sequence identical to or different from one or more reference sequences.

56. The method according to any one of claims 43-54 wherein said method comprises: performing a polymerase chain reaction (PCR) to amplify nucleic acid comprising the coding sequence of said LEC polynucleotide, and determining nucleotide sequence of the amplified nucleic acid.

57. A method of screening for a hereditary lymphedema genotype in a human subject, comprising:

(a) providing a biological sample comprising nucleic acid from said subject, and

(b) analyzing said nucleic acid for the presence of a mutation altering the encoded amino acid sequence of the at least one allele of at least one gene in the human subject relative to a human gene encoding an amino acid sequence selected from the group consisting of SEQ ID NOs: 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38 40, 42, 43, 44, 45, and 46, wherein the presence of a mutation altering the encoded amino acid sequence in the human subject in a manner that correlates with lymphedema in human subjects identifies a hereditary lymphedema genotype.

58. The method according to claim 57 wherein said biological sample is a cell sample.

59. The method according to claim 57 wherein said analyzing comprises sequencing a portion of said nucleic acid.

60. The method according to claim 57 wherein the human subject has a hereditary lymphedema genotype identified by the method of screening.

61. A method of inhibiting lymphangiogenesis comprising administering to a subject an inhibitor of a LEC transmembrane polypeptide,
wherein the LEC transmembrane polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 6, 8, 10, 12, 14, 16, 20, 22, 24, 42 and 43; and
wherein the inhibitor is selected from the group consisting of
(a) a soluble extracellular domain fragment of the LEC transmembrane polypeptide;
(b) an antibody that binds to the extracellular domain of the LEC transmembrane polypeptide;
(c) a polypeptide comprising an antigen binding domain of the antibody according to (b); and
(d) an antisense nucleic acid complementary to the nucleic acid encoding the LEC transmembrane polypeptide or its complement.

62. A method according to claim 61, wherein the inhibitor is a polypeptide comprising an extracellular domain fragment of an LEC polypeptide, wherein the sequence of said extracellular domain is selected from the group consisting of amino acids 1-1005 of SEQ ID NO:43.

63. The method according to claim 61 wherein said subject is a human containing a tumor.

64. A method for modulating lymphangiogenesis in a mammalian subject comprising: administering to a mammalian subject in need of modulation of lymphangiogenesis an antisense molecule to a LEC polynucleotide, in an amount effective to inhibit transcription or translation of the polypeptide encoded by the LEC polynucleotide, wherein the LEC polynucleotide comprises a nucleotide sequence selected from the group consisting of SEQ ID NOS: 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 and 41.

65. A method of treating hereditary lymphedema, comprising:
(a) identifying a human subject with hereditary lymphedema and with a mutation that alters the encoded amino acid sequence of at least one polypeptide of the human subject, relative to the amino acid sequence of a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 6, 8, 10, 12, 14, 16, 20, 22, 24, 42 and 43; and
(b) administering to said subject a lymphatic growth factor selected from the group consisting of a VEGF-C polynucleotide, a VEGF-C polypeptide, a VEGF-D polynucleotide, and a VEGF-D polypeptide.

66. The method according to claim 8, further comprising a second LEC polypeptide selected from the LEC polypeptides set forth in any of Tables 3 and 4 of PCT/US03/06900.

67. The method according to any of claims 16 or 17, further comprising a mutation in at least one allele of a second gene encoding a LEC protein identified in any of Tables 3 and 4 of PCT/US03/06900, wherein the mutation in said second gene correlates with lymphedema in human subjects.

68. The method according to any of claims 18, 19, or 20, further comprising a second BEC or LEC gene that encodes a polypeptide set forth in any of Tables 3 and 4 of PCT/US03/06900.

69. The method according to any of claims 43 or 52, further comprising assaying the nucleic acid for a second mutation that correlates with a hereditary lymphedema phenotype and alters the encoded amino acid sequence of a second gene allele of the human subject when compared to the amino acid sequence of the polypeptide encoded by a corresponding wild-type allele of said second gene, wherein the wild-type polypeptide encoded by said second gene is a polypeptide identified in any of Tables 3 and 4 of PCT/US03/06900.

70. An isolated polypeptide comprising an amino acid sequence at least 95% identical to amino acids 1-1005 of SEQ ID NO: 43.

71. A soluble polypeptide comprising a fragment of the amino acid sequence set forth in SEQ ID NO: 43, wherein said fragment lacks the transmembrane and intracellular amino acids from approximately residues 1006-1028 of SEQ ID NO: 43.

72. An isolated polypeptide comprising an adhesion domain having an amino acid sequence at least 95% identical to a fragment of a polypeptide comprising the amino acid sequence set forth in SEQ ID NO: 20 or SEQ ID NO: 43, wherein said fragment includes at least one vonWillebrand factor domain.

73. An isolated polypeptide comprising a Cache domain and comprising an amino acid sequence at least 95% identical to a fragment of a polypeptide comprising the amino acid sequence set forth in SEQ ID NO: 20 or SEQ ID NO: 43, wherein said fragment includes at least one Cache domain.

74. A fusion protein comprising a polypeptide according to any one of claims 70-73 and a heterologous polypeptide.

75. An antibody that specifically binds to a polypeptide according to any one of claims 70-73.

76. An isolated polypeptide comprising an amino acid sequence at least 95% identical to a protein kinase domain selected from the group consisting of a protein kinase domain set out in SEQ ID NO: 8, in SEQ ID NO: 26, at approximately amino acids 88-342 of SEQ ID NO: 42 and at approximately amino acids 51-317 of SEQ ID NO: 46.

77. An isolated polypeptide comprising an amino acid sequence at least 95% identical to a Zinc binding domain selected from the group consisting of a Zinc-binding domain set out in SEQ ID NO: 36, a Zinc-binding domain at approximately amino acids 527-561 of SEQ ID NO: 6, and a Zinc-binding domain at approximately amino acids 1030-1094 of SEQ ID NO: 45.
78. An isolated polypeptide comprising an amino acid sequence at least 95% identical to a transcription factor selected from the group consisting of SEQ ID NO: 30, SEQ ID NO: 38, SEQ ID NO: 40, and SEQ ID NO: 44.
79. An isolated polypeptide comprising an amino acid sequence at least 95% identical to a cytoskeleton regulation protein selected from the group consisting of SEQ ID NO: 18, SEQ ID NO: 36, and SEQ ID NO: 45.
80. An isolated polypeptide comprising a SPRY domain having an amino acid sequence at least 95% identical to a fragment of a polypeptide comprising the amino acid sequence set forth in SEQ ID NO: 6, wherein said fragment includes at least one SPRY domain.
81. An isolated polypeptide comprising a cation transport activity and comprising an amino acid sequence at least 95% identical to a fragment of a polypeptide comprising the amino acid sequence set forth in SEQ ID NO: 12, wherein said fragment includes at least one cation transport domain.
82. An isolated polypeptide comprising a guanine-nucleotide dissociation stimulator activity and comprising an amino acid sequence at least 95% identical to a fragment of a polypeptide comprising the amino acid sequence set forth in SEQ ID NO: 28, wherein said fragment includes at least one RCC1-like domain.

83. An isolated polypeptide comprising a ubiquitin ligase activity and comprising an amino acid sequence at least 95% identical to a fragment of a polypeptide comprising the amino acid sequence set forth in SEQ ID NO: 28, wherein said fragment includes at least one Hect domain.

84. An isolated polypeptide comprising a Kelch domain and comprising an amino acid sequence at least 95% identical to a fragment of a polypeptide comprising the amino acid sequence set forth in SEQ ID NO: 30 or SEQ ID NO: 44, wherein said fragment includes at least one Kelch domain

85. An isolated polypeptide comprising a GTP-binding activity and comprising an amino acid sequence at least 95% identical to a fragment of a polypeptide comprising the amino acid sequence set forth in SEQ ID NO: 32, wherein said fragment includes at least one GTP-binding domain.

86. A polynucleotide comprising a nucleotide sequence that encodes a polypeptide according to any one of claims 70-85.

87. An expression vector comprising a polynucleotide according to claim 85 operatively linked to an expression control sequence.

88. An expression vector according to claim 87 that is a replication-deficient adenoviral vector or adeno-associated virus vector.

89. A composition comprising the polypeptide of any one of claims 70-87 for use in the methods of the invention.

SEQUENCE LISTING

<110> Alitalo, et al.

<120> Lymphatic and Blood Endothelial Cell Genes

<130> 28967/39230

<160> 46

<170> PatentIn version 3.2

<210> 1

<211> 1997

<212> DNA

<213> Homo sapiens

<400> 1

```

ccccccccgc ctctccaaaa agctacaccg acgcggaccg cggcggcgctc ctccctcgcc      60
ctcgcttcac ctgcggggct ccgaatgcgg ggagctcgga tgtccggttt cctgtgaggc      120
ttttacctga caccgcgcgc ctttccccgg cactggctgg gagggcgccc tgcaaagtgt      180
ggaacgcgga gccccggacc cgctcccgcc gcctccggct cgcccagggg gggtcgccgg      240
gaggagcccc ggggagaggg accaggaggg gcccgcggcc tcgcaggggc gcccgcgccc      300
ccaccctgc ccccgccagc ggaccggtcc cccacccccg gtccttcac catgcacttg      360
ctgggcttct tctctgtggc gtgttctctg ctgcgcgtg cgctgctccc gggtcctcgc      420
gaggcgcccc cgccgcgcgc cgccttcgag tccggactcg acctctcgga cgcgagcccc      480
gacgcgggcg aggccacggc ttatgcaagc aaagatctgg aggagcagtt acggtctgtg      540
tccagtgtag atgaactcat gactgtactc taccagaat attggaaaat gtacaagtgt      600
cagctaagga aaggaggctg gcaacataac agagaacagg ccaacctcaa ctcaaggaca      660
gaagagacta taaaatttgc tgcagcacat tataatacag agatcttgaa aagtattgat      720
aatgagtgga gaaagactca atgcatgcca cgggaggtgt gtatagatgt ggggaaggag      780
tttgagtgcg cgacaaacac cttctttaa cctccatgtg tgtccgtcta cagatgtggg      840
ggttgctgca atagtgaggg gctgcagtgc atgaacacca gcacgagcta cctcagcaag      900
acgttatattg aaattacagt gcctctctct caaggcccca aaccagtaac aatcagtttt      960
gccaatcaca cttcctgccg atgcatgtct aaactggatg ttacagaca agttcattcc     1020
attattagac gttccctgcc agcaacacta ccacagtgtc aggcagcgaa caagacctgc     1080
cccaccaatt acatgtggaa taatcacatc tgcagatgcc tggctcagga agattttatg     1140
ttttcctcgg atgctggaga tgactcaaca gatggattcc atgacatctg tggaccaaac     1200
aaggagctgg atgaagagac ctgtcagtgt gtctgcagag cggggcttcg gcctgccagc     1260
tgtggacccc acaaagaact agacagaaac tcatgccagt gtgtctgtaa aaacaaactc     1320
ttccccagcc aatgtggggc caaccgagaa tttgatgaaa acacatgcca gtgtgtatgt     1380

```

```

aaaagaacct gccccagaaa tcaacccta aatcctggaa aatgtgcctg tgaatgtaca 1440
gaaagtccac agaaatgctt gttaaaagga aagaagttcc accaccaaac atgcagctgt 1500
tacagacggc catgtacgaa ccgccagaag gcttgtgagc caggattttc atatagttaa 1560
gaagtgtgtc gttgtgtccc ttcattattgg aaaagaccac aaatgagcta agattgtact 1620
gttttccagt tcatcgattt tctattatgg aaaactgtgt tgccacagta gaactgtctg 1680
tgaacagaga gacccttgtg ggtccatgct aacaaagaca aaagtctgtc tttcctgaac 1740
catgtggata actttacaga aatggactgg agctcatctg caaaaggcct cttgtaaaga 1800
ctggttttct gccaatgacc aaacagccaa gattttcctc ttgtgatttc tttaaaagaa 1860
tgactatata atttatttcc actaaaaata ttgtttctgc attcattttt atagcaacaa 1920
caattggtaa aactcactgt gatcaatatt tttatatcat gcaaaatatg tttaaaataa 1980
aatgaaaatt gtattat                                     1997

```

```

<210> 2
<211> 419
<212> PRT
<213> Homo sapiens

```

```

<400> 2

```

```

Met His Leu Leu Gly Phe Phe Ser Val Ala Cys Ser Leu Leu Ala Ala
1           5           10           15

```

```

Ala Leu Leu Pro Gly Pro Arg Glu Ala Pro Ala Ala Ala Ala Ala Phe
20           25           30

```

```

Glu Ser Gly Leu Asp Leu Ser Asp Ala Glu Pro Asp Ala Gly Glu Ala
35           40           45

```

```

Thr Ala Tyr Ala Ser Lys Asp Leu Glu Glu Gln Leu Arg Ser Val Ser
50           55           60

```

```

Ser Val Asp Glu Leu Met Thr Val Leu Tyr Pro Glu Tyr Trp Lys Met
65           70           75           80

```

```

Tyr Lys Cys Gln Leu Arg Lys Gly Gly Trp Gln His Asn Arg Glu Gln
85           90           95

```

```

Ala Asn Leu Asn Ser Arg Thr Glu Glu Thr Ile Lys Phe Ala Ala Ala
100          105          110

```

```

His Tyr Asn Thr Glu Ile Leu Lys Ser Ile Asp Asn Glu Trp Arg Lys
115          120          125

```

```

Thr Gln Cys Met Pro Arg Glu Val Cys Ile Asp Val Gly Lys Glu Phe

```

- 3 -

Arg Arg Pro Cys Thr Asn Arg Gln Lys Ala Cys Glu Pro Gly Phe Ser
 385 390 395 400

Tyr Ser Glu Glu Val Cys Arg Cys Val Pro Ser Tyr Trp Lys Arg Pro
 405 410 415

Gln Met Ser

<210> 3
 <211> 2781
 <212> DNA
 <213> Homo sapiens

<400> 3
 atggatatgt ttcctctcac ctgggttttc ttagccctct acttttcaag acaccaagtg 60
 agaggccaac cagacccacc gtgcggaggt cgtttgaatt ccaaagatgc tggctatatc 120
 acctctcccg gttaccccca ggactacccc tcccaccaga actgcgagtg gattgtttac 180
 gccccgaac ccaaccagaa gattgtcctc aacttcaacc ctcactttga aatcgagaag 240
 cagcactgca agtatgactt tatcgagatt cgggatgggg acagtgaatc cgcagacctc 300
 ctgggcaaac actgtgggaa catcgccccg cccaccatca tctcctcggg ctccatgctc 360
 tacatcaagt tcacctccga ctacgccccg cagggggcag gcttctctct gcgctacgag 420
 atcttcaaga caggctctga agattgctca aaaaacttca caagcccaa cgggaccatc 480
 gaatctctg ggtttcctga gaagtatcca cacaacttg actgcacctt taccatcctg 540
 gccaaaccca agatggagat catcctgcag ttctgatct ttgacctgga gcatgacct 600
 ttgcaggtgg gagaggggga ctgcaagtac gattggctgg acatctggga tggcattcca 660
 catgttggcc ccctgattgg caagtactgt gggacaaaa caccctctga acttcgttca 720
 tcgacgggga tcctctccct gacctttcac acggacatgg cgggtggcaa ggatggcttc 780
 tctgcgcgtt actacctggt ccaccaagag ccactagaga actttcagt caatgttcct 840
 ctgggcatgg agtctggccg gattgctaata gaacagatca gtgcctcatc tacctactct 900
 gatgggaggt ggacccctca acaaagccgg ctccatggtg atgacaatgg ctggaccccc 960
 aacttggatt ccaacaagga gtatctccag gtggacctgc gctttttaac catgctcacg 1020
 gccatcgcaa cacagggagc gatttccagg gaaacacaga atggctacta cgtcaaatcc 1080
 tacaagctgg aagtcagcac taatggagag gactggatgg tgtaccggca tggcaaaaac 1140
 cacaaggtat ttcaagccaa caacgatgca actgaggtgg ttctgaacaa gctccacgct 1200
 ccactgctga caaggtttgt tagaatccgc cctcagacct ggcactcagg tatcgccctc 1260
 cggctggagc tcttcggctg ccgggtcaca gatgctccct gctccaacat gctggggatg 1320
 ctctcaggcc tcattgcaga ctcccagatc tccgcctctt ccaccagga atacctctgg 1380

```

agccccagtg cagccccgct ggtcagcagc cgctcgggct ggttccctcg aatccctcag 1440
gccagccccg gtgaggagtg gcttcaggta gatctgggaa caccaagac agtgaaaggt 1500
gtcatcatcc agggagccccg cggaggagac agtatcactg ctgtggaagc cagagcattt 1560
gtgcgcaagt tcaaagtctc ctacagccta aacggcaagg actgggaata cattcaggac 1620
cccaggaccc agcagccaaa gctgttcgaa gggaacatgc actatgacac ccctgacatc 1680
cgaaggtttg accccattcc ggcacagtat gtgcgggtat acccggagag gtggtcgccc 1740
gcgggggattg ggatgcggct ggaggtgctg ggctgtgact ggacagactc caagcccacg 1800
gtaaaaacgc tgggaccacac tgtgaagagc gaagagacaa ccacccccta cccaccgaa 1860
gaggaggcca cagagtgtgg ggagaactgc agctttgagg atgacaaaga tttgcagctc 1920
ccttcgggat tcaattgcaa cttcgatttc ctcgaggagc cctgtggttg gatgtatgac 1980
catgccaaagt ggctccggac cacctgggac agcagctcca gcccaaacga ccggacgttt 2040
ccagatgaca ggaatttctt gcggctgcag agtgacagcc agagagaggg ccagtatgcc 2100
cggctcatca gccccctgt ccacctgcc cgaagcccg tgtgcatgga gttccagtac 2160
caggccacgg gcggcccgcg ggtggcgctg caggtggtgc ggaagccag ccaggagagc 2220
aagttgctgt gggcatccg tgaggaccag ggccgagct ggaagcacgg gcggatcatc 2280
ctgcccagct acgacatgga gtaccagatt gtgttcgagg gagtgatagg gaaaggacgt 2340
tccggagaga ttgccattga tgacattcgg ataagcactg atgtccact ggagaactgc 2400
atggaacca tctcggcttt tgcaaggac atcccagaaa tacatgagag agaaggatat 2460
gaagatgaaa ttgatgatga atacgaggtg gactggagca attcttcttc tgcaacctca 2520
gggtctggcg cccctcgcac cgacaaagaa aagagctggc tgtacaccct ggatcccatc 2580
ctcatcacca tcatcgccat gagctcactg ggcgtcctcc tgggggccac ctgtgcaggc 2640
ctctgtctct actgcacctg ttcctactcg ggctgagct cccgaagctg caccacactg 2700
gagaactaca acttcgagct ctacgatggc cttaagcaca aggtcaagat gaaccaccaa 2760
aagtgctgct ccgaggcatg a 2781

```

```

<210> 4
<211> 926
<212> PRT
<213> Homo sapiens

```

```

<400> 4

```

```

Met Asp Met Phe Pro Leu Thr Trp Val Phe Leu Ala Leu Tyr Phe Ser
1           5           10           15

```

```

Arg His Gln Val Arg Gly Gln Pro Asp Pro Pro Cys Gly Gly Arg Leu
          20           25           30

```

Asn Ser Lys Asp Ala Gly Tyr Ile Thr Ser Pro Gly Tyr Pro Gln Asp
 35 40 45
 Tyr Pro Ser His Gln Asn Cys Glu Trp Ile Val Tyr Ala Pro Glu Pro
 50 55 60
 Asn Gln Lys Ile Val Leu Asn Phe Asn Pro His Phe Glu Ile Glu Lys
 65 70 75 80
 His Asp Cys Lys Tyr Asp Phe Ile Glu Ile Arg Asp Gly Asp Ser Glu
 85 90 95
 Ser Ala Asp Leu Leu Gly Lys His Cys Gly Asn Ile Ala Pro Pro Thr
 100 105 110
 Ile Ile Ser Ser Gly Ser Met Leu Tyr Ile Lys Phe Thr Ser Asp Tyr
 115 120 125
 Ala Arg Gln Gly Ala Gly Phe Ser Leu Arg Tyr Glu Ile Phe Lys Thr
 130 135 140
 Gly Ser Glu Asp Cys Ser Lys Asn Phe Thr Ser Pro Asn Gly Thr Ile
 145 150 155 160
 Glu Ser Pro Gly Phe Pro Glu Lys Tyr Pro His Asn Leu Asp Cys Thr
 165 170 175
 Phe Thr Ile Leu Ala Lys Pro Lys Met Glu Ile Ile Leu Gln Phe Leu
 180 185 190
 Ile Phe Asp Leu Glu His Asp Pro Leu Gln Val Gly Glu Gly Asp Cys
 195 200 205
 Lys Tyr Asp Trp Leu Asp Ile Trp Asp Gly Ile Pro His Val Gly Pro
 210 215 220
 Leu Ile Gly Lys Tyr Cys Gly Thr Lys Thr Pro Ser Glu Leu Arg Ser
 225 230 235 240
 Ser Thr Gly Ile Leu Ser Leu Thr Phe His Thr Asp Met Ala Val Ala
 245 250 255
 Lys Asp Gly Phe Ser Ala Arg Tyr Tyr Leu Val His Gln Glu Pro Leu
 260 265 270
 Glu Asn Phe Gln Cys Asn Val Pro Leu Gly Met Glu Ser Gly Arg Ile
 275 280 285

Ala Asn Glu Gln Ile Ser Ala Ser Ser Thr Tyr Ser Asp Gly Arg Trp
 290 295 300
 Thr Pro Gln Gln Ser Arg Leu His Gly Asp Asp Asn Gly Trp Thr Pro
 305 310 315 320
 Asn Leu Asp Ser Asn Lys Glu Tyr Leu Gln Val Asp Leu Arg Phe Leu
 325 330 335
 Thr Met Leu Thr Ala Ile Ala Thr Gln Gly Ala Ile Ser Arg Glu Thr
 340 345 350
 Gln Asn Gly Tyr Tyr Val Lys Ser Tyr Lys Leu Glu Val Ser Thr Asn
 355 360 365
 Gly Glu Asp Trp Met Val Tyr Arg His Gly Lys Asn His Lys Val Phe
 370 375 380
 Gln Ala Asn Asn Asp Ala Thr Glu Val Val Leu Asn Lys Leu His Ala
 385 390 395 400
 Pro Leu Leu Thr Arg Phe Val Arg Ile Arg Pro Gln Thr Trp His Ser
 405 410 415
 Gly Ile Ala Leu Arg Leu Glu Leu Phe Gly Cys Arg Val Thr Asp Ala
 420 425 430
 Pro Cys Ser Asn Met Leu Gly Met Leu Ser Gly Leu Ile Ala Asp Ser
 435 440 445
 Gln Ile Ser Ala Ser Ser Thr Gln Glu Tyr Leu Trp Ser Pro Ser Ala
 450 455 460
 Ala Arg Leu Val Ser Ser Arg Ser Gly Trp Phe Pro Arg Ile Pro Gln
 465 470 475 480
 Ala Gln Pro Gly Glu Glu Trp Leu Gln Val Asp Leu Gly Thr Pro Lys
 485 490 495
 Thr Val Lys Gly Val Ile Ile Gln Gly Ala Arg Gly Gly Asp Ser Ile
 500 505 510
 Thr Ala Val Glu Ala Arg Ala Phe Val Arg Lys Phe Lys Val Ser Tyr
 515 520 525
 Ser Leu Asn Gly Lys Asp Trp Glu Tyr Ile Gln Asp Pro Arg Thr Gln
 530 535 540

Gln Pro Lys Leu Phe Glu Gly Asn Met His Tyr Asp Thr Pro Asp Ile
 545 550 555 560

Arg Arg Phe Asp Pro Ile Pro Ala Gln Tyr Val Arg Val Tyr Pro Glu
 565 570 575

Arg Trp Ser Pro Ala Gly Ile Gly Met Arg Leu Glu Val Leu Gly Cys
 580 585 590

Asp Trp Thr Asp Ser Lys Pro Thr Val Lys Thr Leu Gly Pro Thr Val
 595 600 605

Lys Ser Glu Glu Thr Thr Thr Pro Tyr Pro Thr Glu Glu Glu Ala Thr
 610 615 620

Glu Cys Gly Glu Asn Cys Ser Phe Glu Asp Asp Lys Asp Leu Gln Leu
 625 630 635 640

Pro Ser Gly Phe Asn Cys Asn Phe Asp Phe Leu Glu Glu Pro Cys Gly
 645 650 655

Trp Met Tyr Asp His Ala Lys Trp Leu Arg Thr Thr Trp Ala Ser Ser
 660 665 670

Ser Ser Pro Asn Asp Arg Thr Phe Pro Asp Asp Arg Asn Phe Leu Arg
 675 680 685

Leu Gln Ser Asp Ser Gln Arg Glu Gly Gln Tyr Ala Arg Leu Ile Ser
 690 695 700

Pro Pro Val His Leu Pro Arg Ser Pro Val Cys Met Glu Phe Gln Tyr
 705 710 715 720

Gln Ala Thr Gly Gly Arg Gly Val Ala Leu Gln Val Val Arg Glu Ala
 725 730 735

Ser Gln Glu Ser Lys Leu Leu Trp Val Ile Arg Glu Asp Gln Gly Gly
 740 745 750

Glu Trp Lys His Gly Arg Ile Ile Leu Pro Ser Tyr Asp Met Glu Tyr
 755 760 765

Gln Ile Val Phe Glu Gly Val Ile Gly Lys Gly Arg Ser Gly Glu Ile
 770 775 780

Ala Ile Asp Asp Ile Arg Ile Ser Thr Asp Val Pro Leu Glu Asn Cys
 785 790 795 800

Met Glu Pro Ile Ser Ala Phe Ala Val Asp Ile Pro Glu Ile His Glu
805 810 815

Arg Glu Gly Tyr Glu Asp Glu Ile Asp Asp Glu Tyr Glu Val Asp Trp
820 825 830

Ser Asn Ser Ser Ser Ala Thr Ser Gly Ser Gly Ala Pro Ser Thr Asp
835 840 845

Lys Glu Lys Ser Trp Leu Tyr Thr Leu Asp Pro Ile Leu Ile Thr Ile
850 855 860

Ile Ala Met Ser Ser Leu Gly Val Leu Leu Gly Ala Thr Cys Ala Gly
865 870 875 880

Leu Leu Leu Tyr Cys Thr Cys Ser Tyr Ser Gly Leu Ser Ser Arg Ser
885 890 895

Cys Thr Thr Leu Glu Asn Tyr Asn Phe Glu Leu Tyr Asp Gly Leu Lys
900 905 910

His Lys Val Lys Met Asn His Gln Lys Cys Cys Ser Glu Ala
915 920 925

<210> 5

<211> 3540

<212> DNA

<213> Homo sapiens

<400> 5

```

ggaggggtgg ggcggtacga agcgggggtg ggctctgcgc gtaatggcag cgccgtggcc      60
tcgcgtccat ctttgccgtt ctctcgacc tgcacaaag gagtcgcgcc gccgccgccg      120
ccccctccct ccggtggggc cgggaggtag agaaaaactg ccattggatg tccagaatcc      180
cctgtagttg ataatgttgg gaataagctc tgcaactttc tttggcattc agttgttaaa      240
aacaaatagg atgcaaattc ctcaactcca ggttatgaaa acagtacttg gaaaactgaa      300
aactacctaa atgatcgtct ttggttgggc cgtgttctta gcgagcagaa gccttggcca      360
gggtctgttg ttgactctcg aagagcacat agcccacttc ctagggactg gaggtgccgc      420
tactaccatg ggtaattcct gtatctgccg agatgacagt ggaacagatg acagtgttga      480
caccacacag caacaggccg agaacagtgc agtaccact gctgacacaa ggagccaacc      540
acgggaccct gttcggccac caaggagggg ccgaggacct catgagccaa ggagaaagaa      600
acaaaatgtg gatgggctag tgttggacac actggcagta atacggactc ttgtagataa      660
tgatcaggaa cctccctatt caatgataac attacacgaa atggcagaaa cagatgaagg      720
atggttggat gttgtccagt ctttaattag agttattcca ctggaagatc cactgggacc      780

```

agctgttata	acattgttac	tagatgaatg	tccattgccc	actaaagatg	cactccagaa	840
attgactgaa	attctcaatt	taaatggaga	agtagcttgc	caggactcaa	gccatcctgc	900
caaacacagg	aacacatctg	cagtcctagg	ctgcttggcc	gagaaactag	caggtcctgc	960
aagtataggt	ttacttagcc	caggaatact	ggaatacttg	ctacagtgtc	tgaagttaca	1020
gtcccacccc	acagtcatgc	tttttgcact	tatcgcactg	gaaaagtttg	cacagacaag	1080
tgaaaataaa	ttgactattt	ctgaatccag	tattagtgtg	cggcttgtca	cattggagtc	1140
ctgggcta	gacatgatt	atctgaaacg	tcaagttggg	ttctgtgccc	agtggagctt	1200
agacaatctc	tttttaaaag	aaggtagaca	gctgacctat	gagaaagtga	acttgagtag	1260
cattagggcc	atgctgaata	gcaatgatgt	cagcgagtac	ctgaagatct	cacctcatgg	1320
cttagaggct	cgctgtgatg	cctcctcttt	tgaaagtgtg	cgttgcacct	tttgtgtgga	1380
tgccggggta	tggtactatg	aagtaacagt	ggcacttctt	ggcgatcatg	agattggctg	1440
ggccactcga	gacagcaaat	tcctcaatca	tgaaggctac	ggcattgggg	atgatgaata	1500
ctcctgtgcg	tatgatggct	gccggcagct	gatttggtac	aatgccagaa	gtaagcctca	1560
catacaccca	tgctggaaag	aaggagatac	agtaggattt	ctgttagact	tgaatgaaaa	1620
gcaaatgatc	ttctttttta	atggcaacca	gctgcctcct	gaaaagcaag	tcttttcctc	1680
tactgtatct	ggattttttg	ctgcagctag	tttcatgtca	tatcaacaat	gtgagttcaa	1740
ttttggagca	aaaccattca	aatacccacc	atctatgaaa	tttagcactt	ttaatgacta	1800
cgcttccta	acagctgaag	aaaaaatcat	tttgccaagg	cacaggcgct	ttgctctgtt	1860
gaagcaagtc	agtatccgag	aaaactgctg	ttccctttgt	tgtgatgagg	tagcagacac	1920
acaattgaag	ccatgtggac	acagtgcact	gtgcatggat	tgtgccttgc	agctggagac	1980
ctgcccattg	tgtcgtaaag	aaatagtatc	tagaatcaga	cagatttctc	atatttcattg	2040
acacatgtga	agaggcatcg	tggacttttt	tctactcaat	tccagccaat	gttgaaaaga	2100
aaaagaaaaa	aaaaactctc	taatcagttg	tacacacatt	gaaacttata	gccatggcca	2160
gattttatgc	taaaaatggg	agtttgtcaa	agacaaaatt	ctcttagaat	ctaattccaac	2220
ttgccagccc	tgagaaaatc	ccttttaagg	ccaaggaaaag	ctgaatgcta	gcagccaggg	2280
ctgtgggtact	tccatgagaa	accatagcag	acaatgccct	cccaagtact	gaaatcacac	2340
tggaatcccc	cttgttgggt	tcatttgatt	gtttaacaca	ggatgtgttg	tgtcattctg	2400
aagtttttat	ttggggcaga	agtctttatg	gagatgtaaa	tgacagcggt	tctgggttat	2460
gcataacttc	tactgggtca	gagacaccgg	tgtgtcaagc	atggatattg	cattgcaaga	2520
cttgaatcta	taaaaattag	aatcacacag	tcagtactac	aagcaaaaca	gagaacctga	2580
aagaagggtgc	acagactgta	agaaaaaacc	caagtttgtg	atatttcagt	gattccaaag	2640
aacattctag	gttttttgtt	tgtttttttg	ttttttgggt	tttttttttt	tactgcagaa	2700

```

aattggtggt attttcacat tcatagtgtt tctatccaat ttcagtaccc acatttaatg 2760
aggaaaaaat gttttaccaa tgaaggagga attcttaa at tagctgtaat gttagggttg 2820
agaaaatttg gtatttaggg tattttcaag gtaccatcaa atcagatttc tgtttttttg 2880
ttaaaaaaaaa tttttttaat cagtattgtt tttacaagta atatactttg aaactcttga 2940
actaatagtc tcaaaaaactc tagaggacag tctgagaaca cgtatttcta ttgttctaaa 3000
taaatacatg tttttgaata gttcaatcat gaattattga ctatgtcttc atcaaaagtg 3060
ttaatccctc tcagggtctc tgggtgaagac cttcaagagt ttgggttttt ctcccaggaa 3120
attggaaggt agaattgtaa attcatagaa cttcttttat aatgggtgtac ctcagcagct 3180
gcctttcaat ttatgccaa gtccttacaga gtttatactt gaatagtaaa tatgtcttct 3240
gagttttaca gtgtcttaaa ctcaatgcac attttttttt cttctttttc cacccttct 3300
tgtttgtagt tcattatacc tgcctatta cagaactgat ttccttcttg gctgtacatg 3360
ttgggggtgct ggattttttt ccgtgtcttt agtcttccat aaatccacac acacacacac 3420
acacaaaaaaa tatatatata tataaatata tatgtaggat acatgttctc ttctttagct 3480
tgtggtgaat acagtaattt gcattgaaga ataaaacatc tgttgccctt tttgactaag 3540

```

```

<210> 6
<211> 576
<212> PRT
<213> Homo sapiens

```

```

<400> 6

```

```

Met Ile Val Phe Gly Trp Ala Val Phe Leu Ala Ser Arg Ser Leu Gly
1           5           10           15

```

```

Gln Gly Leu Leu Leu Thr Leu Glu Glu His Ile Ala His Phe Leu Gly
20           25           30

```

```

Thr Gly Gly Ala Ala Thr Thr Met Gly Asn Ser Cys Ile Cys Arg Asp
35           40           45

```

```

Asp Ser Gly Thr Asp Asp Ser Val Asp Thr Gln Gln Gln Gln Ala Glu
50           55           60

```

```

Asn Ser Ala Val Pro Thr Ala Asp Thr Arg Ser Gln Pro Arg Asp Pro
65           70           75           80

```

```

Val Arg Pro Pro Arg Arg Gly Arg Gly Pro His Glu Pro Arg Arg Lys
85           90           95

```

```

Lys Gln Asn Val Asp Gly Leu Val Leu Asp Thr Leu Ala Val Ile Arg
100          105          110

```

Thr Leu Val Asp Asn Asp Gln Glu Pro Pro Tyr Ser Met Ile Thr Leu
 115 120 125

His Glu Met Ala Glu Thr Asp Glu Gly Trp Leu Asp Val Val Gln Ser
 130 135 140

Leu Ile Arg Val Ile Pro Leu Glu Asp Pro Leu Gly Pro Ala Val Ile
 145 150 155 160

Thr Leu Leu Leu Asp Glu Cys Pro Leu Pro Thr Lys Asp Ala Leu Gln
 165 170 175

Lys Leu Thr Glu Ile Leu Asn Leu Asn Gly Glu Val Ala Cys Gln Asp
 180 185 190

Ser Ser His Pro Ala Lys His Arg Asn Thr Ser Ala Val Leu Gly Cys
 195 200 205

Leu Ala Glu Lys Leu Ala Gly Pro Ala Ser Ile Gly Leu Leu Ser Pro
 210 215 220

Gly Ile Leu Glu Tyr Leu Leu Gln Cys Leu Lys Leu Gln Ser His Pro
 225 230 235 240

Thr Val Met Leu Phe Ala Leu Ile Ala Leu Glu Lys Phe Ala Gln Thr
 245 250 255

Ser Glu Asn Lys Leu Thr Ile Ser Glu Ser Ser Ile Ser Asp Arg Leu
 260 265 270

Val Thr Leu Glu Ser Trp Ala Asn Asp Pro Asp Tyr Leu Lys Arg Gln
 275 280 285

Val Gly Phe Cys Ala Gln Trp Ser Leu Asp Asn Leu Phe Leu Lys Glu
 290 295 300

Gly Arg Gln Leu Thr Tyr Glu Lys Val Asn Leu Ser Ser Ile Arg Ala
 305 310 315 320

Met Leu Asn Ser Asn Asp Val Ser Glu Tyr Leu Lys Ile Ser Pro His
 325 330 335

Gly Leu Glu Ala Arg Cys Asp Ala Ser Ser Phe Glu Ser Val Arg Cys
 340 345 350

Thr Phe Cys Val Asp Ala Gly Val Trp Tyr Tyr Glu Val Thr Val Val
 355 360 365

Thr Ser Gly Val Met Gln Ile Gly Trp Ala Thr Arg Asp Ser Lys Phe
 370 375 380

Leu Asn His Glu Gly Tyr Gly Ile Gly Asp Asp Glu Tyr Ser Cys Ala
 385 390 395 400

Tyr Asp Gly Cys Arg Gln Leu Ile Trp Tyr Asn Ala Arg Ser Lys Pro
 405 410 415

His Ile His Pro Cys Trp Lys Glu Gly Asp Thr Val Gly Phe Leu Leu
 420 425 430

Asp Leu Asn Glu Lys Gln Met Ile Phe Phe Leu Asn Gly Asn Gln Leu
 435 440 445

Pro Pro Glu Lys Gln Val Phe Ser Ser Thr Val Ser Gly Phe Phe Ala
 450 455 460

Ala Ala Ser Phe Met Ser Tyr Gln Gln Cys Glu Phe Asn Phe Gly Ala
 465 470 475 480

Lys Pro Phe Lys Tyr Pro Pro Ser Met Lys Phe Ser Thr Phe Asn Asp
 485 490 495

Tyr Ala Phe Leu Thr Ala Glu Glu Lys Ile Ile Leu Pro Arg His Arg
 500 505 510

Arg Leu Ala Leu Leu Lys Gln Val Ser Ile Arg Glu Asn Cys Cys Ser
 515 520 525

Leu Cys Cys Asp Glu Val Ala Asp Thr Gln Leu Lys Pro Cys Gly His
 530 535 540

Ser Asp Leu Cys Met Asp Cys Ala Leu Gln Leu Glu Thr Cys Pro Leu
 545 550 555 560

Cys Arg Lys Glu Ile Val Ser Arg Ile Arg Gln Ile Ser His Ile Ser
 565 570 575

<210> 7

<211> 2589

<212> DNA

<213> Homo sapiens

<400> 7

ttggagacgg agtctcactc tgtcacccat tctggagtac aatggcatga tctcagctca 60

ctgcaacctc ctcttctggt gttcaagtga ttcttctgcc tcagcctcct gagtagctgg 120

gactacaggc ttgtgccacc aactcagct aatttttgta ttttcagtag agacgggggc 180

tccccatatt	ggccaggctg	gtcttgaact	cctgacctca	aacgatctgc	ccaccccagc	240
ctcccaaagt	gctgggatta	caagtgtgag	ccaccacacc	cagcctaata	taactgtttt	300
cctttccttt	tagtctttat	ctcattgtta	tatgtgatta	taatgttgtc	atttatgcat	360
tcacatcacc	agtaattatc	tcaaacagat	cttaaagtct	gcctatcatg	ttgaagtcac	420
gtttcactca	gggtcgctcag	ttacaaggat	gctatgggag	caaatacaac	agatcatcca	480
gcgcatacaca	tgggtgagcc	cacctgccat	cactctggaa	tggagagga	aggtggccca	540
ggaagccatt	gagagcctca	gcgcctccaa	attggctaag	agcatttgca	gccaattccg	600
gactcggctc	aatagttccc	acgaggcttt	tgcagcctcc	ttgcggcagc	tggagactgg	660
ccactcaggc	cggttagaga	aaacggaaga	tctatggctg	agggttcgga	aagatcatgc	720
tccccgcctg	gcccgctttt	ctctggaaag	ctgttcttta	caggatgtct	tgcttcacgc	780
taaacctaaa	ctgggacagg	aactgggccc	gggccagtat	ggtgtggtat	acctgtgtga	840
caactgggga	ggacacttcc	cttgtgcctt	caaatacagt	gtccctccag	atgagaagca	900
ctggaatgat	ctggcttttg	aatttcacta	tatgaggtct	ctgccgaagc	atgagcgatt	960
ggtggatctc	catggttcag	tcattgacta	caactatggt	ggtggctcca	gcattgctgt	1020
gctcctcatt	atggagcggc	tacaccggga	tctctacaca	gggctgaagg	ctgggctgac	1080
cctggagaca	cgtttgcaga	tagcactaga	tgtggtggag	ggaatccgct	tctctcacag	1140
ccagggaact	gtccatcgtg	atatcaaact	gaaaaatgtg	ctgctggata	agcagaaccg	1200
tgccaagatc	actgacttag	gattctgcaa	gccagaggcc	atgatgtcag	gcagcattgt	1260
ggggacacca	atccatatgg	cccctgaact	tttcacaggg	aagtacgata	attccgtgga	1320
tgtctacgct	tttggaaatt	ttttctggta	tatctgtctc	ggctctgtca	agctccctga	1380
ggcattttgag	aggtgtgcta	gcaaagacca	tctctggaac	aatgtgcgga	ggggggctcg	1440
cccagaacgt	cttcctgtgt	ttgatgagga	gtgctggcag	ttgatggaag	cctgttgga	1500
tggcgacccc	ttgaagaggc	ctctcttggg	cattgtccag	cccatgctcc	agggcatcat	1560
gaatcggctc	tgcaagtcca	attctgagca	gccaaacaga	ggactagatg	attctacttg	1620
aaagcaaaga	cctttctctt	tcactctcta	gttatttcct	tccccctcac	cttttgcca	1680
tggggagaaat	ttgacattta	ttcactatag	gacacactcc	caagggaact	ggtgcttgct	1740
gggaaacttg	gaaccttccc	aggcagggat	gactcctgga	cagtgaagag	ttgaatgact	1800
gagcatattc	agcagctcac	tgaagcgcca	agctatccct	ttagcaaaaa	agtgtctcag	1860
atgtgtaaaa	gctgaggaat	gtggtgttct	ggcttcacaa	atgaaaagga	ggcagatgtt	1920
accattgtct	tttactgtta	tatacttcta	agacagcaag	cgggacactg	cagtggcaat	1980
agtgttaaaa	aatctcattc	tcattgattt	tggctctagc	taggaatagt	ttagtcagga	2040
ctcagaaaat	ttgagcttag	ttctgggatt	atgaaaacat	ggggacaaaa	acaataactt	2100

```

gtggatgtct ggttcctgct gtctgcagcc aggtatctca ggttgcaatg gtcagaagtc 2160
ccagtgaggg agctagaaac agagctatct gttcctcata gtgccagtgt gtttacattt 2220
ataggaccac atatccctgg tttctggggg aggggtttcac tgttacatca ccaagaggcc 2280
ctgaaaagag gaaaaaaaaa aaaaaagagg ccaggcatgg tggctcacgc ctgtaatctc 2340
agcactttgg gaggctgagg cgggcggatc acaagggtcca gagttcaaga gcagcctggc 2400
caacatgggtg aaaccccgtc tctactaaag atacaaaaaa ttagccgggc gtggtggtgc 2460
atgcctgtaa tccaagatac tcaggaggct gaggcaggag aatcgcttga acctgggagg 2520
cggaggttgc agtgagccga gatcatgcca ttgcactcca gcctgggcga cagagtgaga 2580
ctctgtctc 2589

```

```

<210> 8
<211> 390
<212> PRT
<213> Homo sapiens

```

```

<400> 8

```

```

Met Leu Trp Glu Gln Ile Lys Gln Ile Ile Gln Arg Ile Thr Trp Val
1           5           10           15

```

```

Ser Pro Pro Ala Ile Thr Leu Glu Trp Lys Arg Lys Val Ala Gln Glu
          20           25           30

```

```

Ala Ile Glu Ser Leu Ser Ala Ser Lys Leu Ala Lys Ser Ile Cys Ser
          35           40           45

```

```

Gln Phe Arg Thr Arg Leu Asn Ser Ser His Glu Ala Phe Ala Ala Ser
          50           55           60

```

```

Leu Arg Gln Leu Glu Ala Gly His Ser Gly Arg Leu Glu Lys Thr Glu
65           70           75           80

```

```

Asp Leu Trp Leu Arg Val Arg Lys Asp His Ala Pro Arg Leu Ala Arg
          85           90           95

```

```

Leu Ser Leu Glu Ser Cys Ser Leu Gln Asp Val Leu Leu His Arg Lys
          100          105          110

```

```

Pro Lys Leu Gly Gln Glu Leu Gly Arg Gly Gln Tyr Gly Val Val Tyr
          115          120          125

```

```

Leu Cys Asp Asn Trp Gly Gly His Phe Pro Cys Ala Leu Lys Ser Val
          130          135          140

```

```

Val Pro Pro Asp Glu Lys His Trp Asn Asp Leu Ala Leu Glu Phe His

```


145		150		155		160
Tyr Met Arg Ser	Leu Pro Lys His Glu Arg	Leu Val Asp	Leu His Gly			
	165		170			175
Ser Val Ile Asp	Tyr Asn Tyr Gly Gly Gly	Ser Ser Ile	Ala Val Leu			
	180		185			190
Leu Ile Met Glu	Arg Leu His Arg Asp	Leu Tyr Thr	Gly Leu Lys Ala			
	195		200			205
Gly Leu Thr Leu	Glu Thr Arg Leu Gln Ile	Ala Leu Asp	Val Val Glu			
	210		215			220
Gly Ile Arg Phe	Leu His Ser Gln Gly	Leu Val His Arg	Asp Ile Lys			
	225		230			235
Leu Lys Asn Val	Leu Leu Asp Lys Gln Asn	Arg Ala Lys Ile	Thr Asp			
	245		250			255
Leu Gly Phe Cys	Lys Pro Glu Ala Met Met	Ser Gly Ser Ile	Val Gly			
	260		265			270
Thr Pro Ile His	Met Ala Pro Glu Leu Phe	Thr Gly Lys Tyr	Asp Asn			
	275		280			285
Ser Val Asp Val	Tyr Ala Phe Gly Ile Leu	Phe Trp Tyr Ile	Cys Ser			
	290		295			300
Gly Ser Val Lys	Leu Pro Glu Ala Phe Glu	Arg Cys Ala Ser	Lys Asp			
	305		310			315
His Leu Trp Asn	Asn Val Arg Arg Gly	Ala Arg Pro Glu	Arg Leu Pro			
	325		330			335
Val Phe Asp Glu	Glu Cys Trp Gln Leu Met	Glu Ala Cys Trp	Asp Gly			
	340		345			350
Asp Pro Leu Lys	Arg Pro Leu Leu Gly	Ile Val Gln Pro	Met Leu Gln			
	355		360			365
Gly Ile Met Asn	Arg Leu Cys Lys Ser Asn	Ser Glu Gln Pro	Asn Arg			
	370		375			380
Gly Leu Asp Asp	Ser Thr					
	385					390

<210> 9
 <211> 5766
 <212> DNA
 <213> Homo sapiens

<400> 9
 cgccgcccga gccgcccga ccaacgcca caacaacgtg tcgggagcag ccgcgcccga 60
 agccggagca ggggcccggag tcgcggtctg agtggggcct gcgcgtgtga gtggccgaga 120
 gttgcccgtg ccggagacct caaactccgg gcgggagcgc tttcgggtggg ccgagccgcc 180
 gtcgcccgcg gcaccgtgtc cccaaggccc agacagcggg gcgggggtct atcaggcgcg 240
 ggactggctt ctctcaggcc gaccttcta gccgggtcc ggatcttct cggcctcggg 300
 ccccttcac gttcaggacg gaggcggag aacgggtgat atcacacacc ccagccggct 360
 tcctggccag ggcattttct ctgcggcca gcagtgttg gattgtagct gctgggagcc 420
 cctcctgcc gccacctctc ctgcaaggaa gggactgacg cgttccact ttcttactct 480
 gcacctccc actctcttc cattcaagaa cgagtttctc ttccctctcc attggaaatc 540
 ctgctccgga ggtccaaaac acgtatttta agcgtgccct tcctcacctc ctggagcacc 600
 cttagaagta ttgcctactt atccgggtct agaatccttc atttttgacc tggctttttt 660
 ttcgcccttt gggagataaa ggtccctctc caccctctac taacactctg caccgaaggc 720
 cttatccttt ggggtcacca gtccttggc catttctatg tgatttcccc caccatctg 780
 agttccagtt tcctctgggc tccaatctcc agtccctggc ggatctggtc agtcccacc 840
 ctaggtggga gtgaccaggg ggctctggcc caggattccc aaccctggaa ggcagctcca 900
 aggcaggaag agaattgggc atcgggtacg aacctggcag ctcaggagtc ggggtccac 960
 tcaccccaca caaaaagatg aaaaagcgca aagagctcaa tgcattgatt ggtttggctg 1020
 gggacagccg gagaaagaag cccaagaaag gcccaagcag tcaccgctg cttcgactg 1080
 agcctcccga ctcagactct gagtccagct ccgaagagga agaggaattc ggtgtggtg 1140
 gaaatcgctc tcgctttgcc aaggagact atttacgatg ctgcaagatc tgttatccgc 1200
 tctgtggttt tgtcatcctt gctgcctgtg ttgtggcctg tgttggttg gtgtggatgc 1260
 aggttgctct caaggaggat ctggatgcc tcaaggaaaa atttogaaca atggaatcta 1320
 atcagaaaag ctcatccaa gaaatccca aacttaatga agaactactc agcaagcaaa 1380
 aacaacttga gaagattgaa tctggagaga tgggtttgaa caaagtctgg ataaacatca 1440
 cagaaatgaa taagcagatt tctctgttga cttctgcagt gaaccacctc aaagccaatg 1500
 ttaagtcagc tgcagacttg attagcctgc ctaccactgt agagggactt cagaagagtg 1560
 tagcttccat tggcaatact ttaaacagcg tccatcttgc tgtggaagca ctacagaaaa 1620
 ctgtggatga acacaagaaa acgatggaat tactgcagag tgatatgaat cagcacttct 1680
 tgaaggagac tcctggaagc aaccagatca ttccgtcacc ttcagccaca tcagaacttg 1740

acaataaaac	ccacagtga	aatttgaaac	aggatatcct	gtaccttcac	aactctttag	1800
aggaggtaaa	cagtgccta	gtgggtacc	agagacagaa	tgatcttaa	ctcgaggga	1860
tgaacgagac	agtcagta	cttaccaga	gagtcaacct	gatagaaagc	gatgtggtg	1920
ctatgagcaa	ggtagaaa	aaagcaaacc	tgctcttcag	catgatgggt	gatagatctg	1980
ccactctgaa	aagacagt	ttggatcaag	tcaccaacag	aacagataca	gtaaaaatcc	2040
aaagcataaa	gaaagaag	agttcaaatt	ctcaggtatc	caagctaaga	gagaaactcc	2100
agctgatcag	tgctcttaca	aacaaacctg	agagcaacag	gcctccagag	accgccgatg	2160
aagagcaagt	agagagtttc	acatcaaagc	catcagcatt	gccaaaat	tcacagtttc	2220
ttggagaccc	agttgagaaa	gctgccaac	taagacctat	ctccctacca	ggagtttcta	2280
gcactgaaga	tcttcaggat	ttattccgca	agactggcca	ggacgtggat	gggaagctga	2340
cctaccagga	aatctggacc	tccctaggtt	ctgctatgcc	agaaccagag	agcttgagag	2400
catttgattc	cgatggagat	ggaagatact	cattcctgga	gctaagggtg	gctttaggta	2460
tctagcttca	tcaggcatat	tttagaaatg	gactgcctaa	tatctattta	cctaacaaca	2520
aaacaaccct	tacttaccca	tcagtcctct	agtcctccaa	actactgtag	cagatacttt	2580
gccacctttt	aacttgtttg	aagaagctat	ataaaagtta	tttttttaa	gaagaagacc	2640
attttactta	tgatgttcag	aatctatga	tttctacaa	ccagtaagat	cttacat	2700
aaaattgcca	gaaaaaaaa	taaagccctc	ttttttctc	tttcttttt	ttgaggggag	2760
gagaccttat	cttttaaagc	tgggaaatgt	atatagagag	agaataagcc	acttttatat	2820
ttcacttaaa	tttgccctaa	attagctgca	ctttatagag	actcagaaaa	tgtcttttct	2880
ttaaaagata	ggccttttct	gtttgtaaat	atttaaata	aagaaagcat	tgtgcatatt	2940
gtgtggaaag	taggaagaat	ggttttgaac	aggatatgaa	caaatgactt	attaaaaatt	3000
gctgatctgg	tgtaggtggc	agctgaaact	acatccatgt	ctccataagg	catccctcaa	3060
aggcccaggc	gctgccaggg	ggtttgcct	ggtagctgga	ggaaccgatt	tcagggaata	3120
gacactggag	acaatactga	ctccaggcat	ggctcatgga	agtaggattc	tggttctttg	3180
ttcctattcc	ctcagcta	ccaacctgg	gaatcagaga	agtcttgggg	atttttctca	3240
tttttagtac	tatttcaggg	tttatgagca	taaaaagtta	tccattgggg	agctccattt	3300
tccctgctga	gtgagctaga	ttgccttccc	caccaccca	cttaagtctg	tcttaaagcc	3360
gtagctggct	cccaccacca	gtaccatctc	catttgaaatg	gcagggctaa	attccccag	3420
ccattatctc	acactgacca	cccagagctt	tagaagagag	ctgtgcttct	aattttgacc	3480
cagaaaacca	tacccttga	gattttacct	agaggcta	caagagccta	atatgtttct	3540
ctgggggatg	actaaagcca	aaaaggctgt	gagatgaaac	atgtgaaata	atattcagtt	3600
tccttaccat	taccagctca	gaagtagcta	gaggctttct	acccaaagga	tgccaaagta	3660

tagcagggca	ggcctggagc	tagggccttc	acatgggtgg	agcaagtttt	tcaaatactaa	3720
tacaatacaag	tacaatactt	cctttaaatg	cttctgtgga	cctggcatga	aagatcccta	3780
gattgaaagg	aataatacct	ccatgtctcc	tgtatgttga	gtctagaatt	gctgtgttgt	3840
tcttagaagc	agtctttggg	caacaacttg	aaaggggaaa	aaaaaactac	aaaaacttaa	3900
ctttggtata	ggccaagtca	gggagaaagt	agagaaagct	gtcatgccac	agacttcttt	3960
agtggagatc	atttcctttt	taactttgtt	caggttgccc	ttcaccatgg	atacagtccg	4020
gtacccttaa	acatttaagg	gctgtttttt	tttttcttta	tatgatgttc	agcttggtat	4080
taaccaaaact	taaatttttt	ttccagaagt	attaaaattt	agttaaagca	agatgaagtc	4140
ttttcccatg	aatgtgtccc	cttatctcat	tatagcttat	gctcagtttc	actttcttgg	4200
aaagggttaa	acaaattagc	ctggcacttt	aggtaacttg	aaaataactc	tcacccttct	4260
gactgccttc	catccactct	cacccacac	ctttttttta	tatatataca	cccttacaga	4320
ttttctaaca	accaaataaa	attctagcaa	taaattgaat	tatactgcta	tatttgtata	4380
tactgtacca	taaatagtat	gtctgtacct	ggaaggattt	tttttgagaa	ctgatttata	4440
tgaaatatac	ttgagggtaa	tgtagcttgt	cctttttagt	ttcaaattct	tattcatagg	4500
cagatacttt	gaagacacct	taactctttg	ttgtgtgtat	tttcccttgc	atcagataac	4560
tatacatgga	gaaatccatt	ttgttttttg	gtttatcttc	tgttcttctc	accttgtgct	4620
tgtctgacta	gtatcacccc	tgagccagta	tacagtacca	tcctctcccc	ctaccaagtt	4680
tggcagaggt	gtttatactg	cattctcaga	tctatgaata	tttttgtcat	cctgcctgag	4740
aaagtacatc	ttccaggcaa	aagtagggta	tcttgaactc	ctttcttctg	agttgagcag	4800
actcactggc	ttaagcatcc	cataagccta	ttatttagtg	tgaatttggg	acttttatatt	4860
gtatgttgtc	ttaacttact	gagcctgtga	cttgtattta	gctgtagtca	agcctcacca	4920
gccatatctc	atctctgaga	ggaagactgg	atagtgaagt	tggttttggg	tttggtttta	4980
aaacctttta	ccttttcagt	ctggcagctc	ctactgctgc	tttcctaagt	tactcaaagc	5040
actgcttctt	ggtccttggt	ttcttttgcc	ttgtactatc	ttgtataatg	tttactaaaa	5100
gagcctcccc	ataaatcaga	gaggcaaaaa	ggtagcatat	aaaaagaaga	acaaaaagaa	5160
gagaaaacct	gacttctaga	tgactgtggg	gtcatctaaa	aggcagaggt	gcctgaccct	5220
caggaatgtg	tatgatgcaa	cagagctgaa	gaatgggttc	ccttcaccta	ctgctggcca	5280
ttgtcccttc	tgttgagat	tcctttcttt	atgggtgctg	ttcataacct	cttcaacacc	5340
cccccttget	ggtcctgccg	gtcacagttg	aggattttgg	ctgtgatggg	ctcactactca	5400
taatggcctg	ttgagttggt	tttagtgaca	ggttttgttt	gctgcacacc	tcattggcct	5460
gcttaccac	accagctaga	agtggctcct	ctttcatatc	caaaccagaa	cacatttggg	5520
tattcctgag	actttataga	gcgtgtattg	tttgttatac	gaacctaac	taccttgttg	5580

ttttctcata ttaagaaatt taaatctact ttgttttagt gaagctatatt tggtaatgta 5640
 taagcaatttt tggttctgtt tagggagaat tcagggtttc catgacttga tttgagcttt 5700
 gcatattttca aataagtaac acagccattt taatttagct aaataaaaaa atttcttttt 5760
 tctctc 5766

<210> 10
 <211> 495
 <212> PRT
 <213> Homo sapiens

<400> 10

Met Lys Lys Arg Lys Glu Leu Asn Ala Leu Ile Gly Leu Ala Gly Asp
 1 5 10 15
 Ser Arg Arg Lys Lys Pro Lys Lys Gly Pro Ser Ser His Arg Leu Leu
 20 25 30
 Arg Thr Glu Pro Pro Asp Ser Asp Ser Glu Ser Ser Ser Glu Glu Glu
 35 40 45
 Glu Glu Phe Gly Val Val Gly Asn Arg Ser Arg Phe Ala Lys Gly Asp
 50 55 60
 Tyr Leu Arg Cys Cys Lys Ile Cys Tyr Pro Leu Cys Gly Phe Val Ile
 65 70 75 80
 Leu Ala Ala Cys Val Val Ala Cys Val Gly Leu Val Trp Met Gln Val
 85 90 95
 Ala Leu Lys Glu Asp Leu Asp Ala Leu Lys Glu Lys Phe Arg Thr Met
 100 105 110
 Glu Ser Asn Gln Lys Ser Ser Phe Gln Glu Ile Pro Lys Leu Asn Glu
 115 120 125
 Glu Leu Leu Ser Lys Gln Lys Gln Leu Glu Lys Ile Glu Ser Gly Glu
 130 135 140
 Met Gly Leu Asn Lys Val Trp Ile Asn Ile Thr Glu Met Asn Lys Gln
 145 150 155 160
 Ile Ser Leu Leu Thr Ser Ala Val Asn His Leu Lys Ala Asn Val Lys
 165 170 175
 Ser Ala Ala Asp Leu Ile Ser Leu Pro Thr Thr Val Glu Gly Leu Gln
 180 185 190

Lys Ser Val Ala Ser Ile Gly Asn Thr Leu Asn Ser Val His Leu Ala
 195 200 205

Val Glu Ala Leu Gln Lys Thr Val Asp Glu His Lys Lys Thr Met Glu
 210 215 220

Leu Leu Gln Ser Asp Met Asn Gln His Phe Leu Lys Glu Thr Pro Gly
 225 230 235 240

Ser Asn Gln Ile Ile Pro Ser Pro Ser Ala Thr Ser Glu Leu Asp Asn
 245 250 255

Lys Thr His Ser Glu Asn Leu Lys Gln Asp Ile Leu Tyr Leu His Asn
 260 265 270

Ser Leu Glu Glu Val Asn Ser Ala Leu Val Gly Tyr Gln Arg Gln Asn
 275 280 285

Asp Leu Lys Leu Glu Gly Met Asn Glu Thr Val Ser Asn Leu Thr Gln
 290 295 300

Arg Val Asn Leu Ile Glu Ser Asp Val Val Ala Met Ser Lys Val Glu
 305 310 315 320

Lys Lys Ala Asn Leu Ser Phe Ser Met Met Gly Asp Arg Ser Ala Thr
 325 330 335

Leu Lys Arg Gln Ser Leu Asp Gln Val Thr Asn Arg Thr Asp Thr Val
 340 345 350

Lys Ile Gln Ser Ile Lys Lys Glu Asp Ser Ser Asn Ser Gln Val Ser
 355 360 365

Lys Leu Arg Glu Lys Leu Gln Leu Ile Ser Ala Leu Thr Asn Lys Pro
 370 375 380

Glu Ser Asn Arg Pro Pro Glu Thr Ala Asp Glu Glu Gln Val Glu Ser
 385 390 395 400

Phe Thr Ser Lys Pro Ser Ala Leu Pro Lys Phe Ser Gln Phe Leu Gly
 405 410 415

Asp Pro Val Glu Lys Ala Ala Gln Leu Arg Pro Ile Ser Leu Pro Gly
 420 425 430

Val Ser Ser Thr Glu Asp Leu Gln Asp Leu Phe Arg Lys Thr Gly Gln
 435 440 445

Asp Val Asp Gly Lys Leu Thr Tyr Gln Glu Ile Trp Thr Ser Leu Gly
 450 455 460

Ser Ala Met Pro Glu Pro Glu Ser Leu Arg Ala Phe Asp Ser Asp Gly
 465 470 475 480

Asp Gly Arg Tyr Ser Phe Leu Glu Leu Arg Val Ala Leu Gly Ile
 485 490 495

<210> 11
 <211> 6550
 <212> DNA
 <213> Homo sapiens

<400> 11
 tcccggaagg caaatggctg gcgtggaagc acaaccgct ttcactcttc gaatttgtgc 60
 ttagctcttt tcttgtagct tgcgactcgt gaccaacatg ctgtgatgtg tgccgagggg 120
 ggaattgggtc agctacacaa cctggatctt accacagttt ggatatgact gaggtctctc 180
 aatggggccag atatcactgg cgacggctga tcagaggtgc aaccagggat gatgattcag 240
 ggccatacaa ctattcctcg ttgctcgctt gtgggcgcaa gtcctctcag acccctaaac 300
 tgtcaggaag gcaccggatt gttgttcccc acatccagcc cttcaaggat gagtatgaga 360
 agttctccgg agcctatgtg aacaatcgaa tacgaacaac aaagtacaca cttctgaatt 420
 ttgtgccaaag aaatttatatt gaacaatttc acagagctgc caatttatat ttctgttcc 480
 tagttgtcct gaactgggta cctttggtag aagccttcca aaaggaaatc accatgttgc 540
 ctctgggtggt ggtccttaca attatcgcaa ttaaagatgg cctggaagggt tatcggaat 600
 acaaaattga caaacagatc aataatttaa taactaaagt ttatagtagg aaagagaaaa 660
 aatacattga cagatgctgg aaagacgtta ctgttgggga ctttattcgc ctctcctgca 720
 acgaggtcat ccctgcagac atggtactac tcttttccac tgatccagat ggaatctgtc 780
 acattgagac ttctgggtctt gatggagaga gcaatttaaa acagaggcag gtggttcggg 840
 gatatgcaga acaggactct gaagttgatc ctgagaagtt ttccagtagg atagaatgtg 900
 aaagcccaaa caatgacctc agcagattcc gaggttctct agaacattcc aacaaagaac 960
 gcgtgggtct cagtaaagaa aatttggtgc ttagaggatg caccattaga aacacagagg 1020
 ctgttggtgg cattgtggtt tatgcaggcc atgaaaccaa agcaatgctg aacaacagt 1080
 ggccacggta taagcgcagc aaattagaaa gaagagcaaa cacagatgtc ctctgggtgtg 1140
 tcatgcttct ggtcataatg tgcttaactg gcgcagtagg tcatggaatc tggctgagca 1200
 ggtatgaaaa gatgcatttt ttcaatgttc ccgagcctga tggacatatc atatcaccac 1260
 tggtggcagg attttatatg ttttggacca tgatcatttt gttacaggtc ttgattccta 1320
 tttctctcta tgtttccatc gaaattgtga agcttggaca aatatatttc attcaaagt 1380

atgtggattt	ctacaatgaa	aaaatggatt	ctattgttca	gtgccgagcc	ctgaacatcg	1440
ccgaggatct	gggacagatt	cagtacctct	tttccgataa	gacaggaacc	ctcactgaga	1500
ataagatggg	ttttcgaaga	tgtagtgtgg	caggatttga	ttactgccat	gaagaaaatg	1560
ccaggagggt	ggagtcctat	caggaagctg	tctctgaaga	tgaagatttt	atagacacag	1620
tcagtgggtc	cctcagcaat	atggcaaaac	cgagagcccc	cagctgcagg	acagttcata	1680
atgggccttt	gggaaataag	ccctcaaata	atcttgctgg	gagctctttt	actctaggaa	1740
gtggagaagg	agccagtga	gtgcctcatt	ccagacaggc	tgctttcagt	agccccattg	1800
aaacagacgt	ggtaccagac	accaggcttt	tagacaaatt	tagtcagatt	acacctcggc	1860
tctttatgcc	actagatgag	accatccaaa	atccaccaat	ggaaactttg	tacattatcg	1920
actttttcat	tgcattggca	atttgcaaca	cagtagtggg	ttctgctcct	aaccaacccc	1980
gacaaaagat	cagacaccct	tactgggggg	ggttgcccat	taagtctttg	gaagagatta	2040
aaagtctttt	ccagagatgg	tctgtccgaa	gatcaagttc	tccatcgctt	aacagtggga	2100
aagagccatc	ttctggaggt	ccaaacgcct	ttgtgagcag	actccctctc	tttagtcgaa	2160
tgaaaccagc	ttcacctgtg	gaggaagagg	tctcccagggt	gtgtgagagc	ccccagtgtc	2220
ccagtagctc	agcttgctgc	acagaaacag	agaaacaaca	cggtgatgca	ggcctcctga	2280
atggcaaggc	agagtccttc	cctggacagc	cattggcctg	caacctgtgt	tatgaggccg	2340
agagcccaga	cgaagcggcc	ttagtgtatg	ccgccagggc	ttaccaatgc	actttacggg	2400
ctcggacacc	agagcaggtc	atggtggact	ttgctgcttt	gggaccatta	acatttcaac	2460
tcttacacat	cctgcccttt	gactcagtaa	gaaaaagaat	gtctgttgtg	gtccgacacc	2520
ctctttccaa	tcaagttgtg	gtgtatacga	aaggcgctga	ttctgtgatc	atggagttac	2580
tgtcgggtgg	ttccccagat	ggagcaagtc	tggagaaaca	acagatgata	gtaagggaga	2640
aaaccagaa	gcacttggat	gactatgcca	aacaaggcct	tctacttta	tgtatagcaa	2700
agaaggatcat	gagtgcact	gaatatgcag	agtggctgag	gaatcatttt	ttagctgaaa	2760
ccagcattga	caacagggaa	gaattactac	ttgaatctgc	catgagggtg	gagaacaaac	2820
ttacattact	tggtgctact	ggcattgaag	accgtctgca	ggagggagtc	cctgaatcta	2880
tagaagctct	tcacaaagcg	ggcatcaaga	tctggatgct	gacaggggac	aagcaggaga	2940
cagctgtcaa	catagcttat	gcatgcaaac	tactggagcc	agatgacaag	ctttttatcc	3000
tcaataccca	aagtaaagat	gcctgtggga	tgctgatgag	cacaattttg	aaagaacttc	3060
agaagaaaac	tcaagccctg	ccagagcaag	tgtcattaag	tgaagattta	cttcagcctc	3120
ctgtcccccg	ggactcaggg	ttacgagctg	gactcattat	cactgggagg	accctggagt	3180
ttgccctgca	agaaagtctg	caaaagcagt	tcttgggaact	gacatcttgg	tgtcaagctg	3240
tggtctgctg	ccgagccaca	ccgctgcaga	aaagtgaagt	ggtgaaattg	gtccgcagcc	3300

atctccaggt	gatgaccctt	gctattggtg	atggtgccaa	tgatgttagc	atgatacaag	3360
tggcagacat	tgggataggg	gtctcaggtc	aagaaggcat	gcaggctgtg	atggccagtg	3420
actttgccgt	ttatcagttc	aaacatctca	gcaagctcct	tcttgtccat	ggacactggg	3480
gttatacacg	gctttccaac	atgattctct	atTTTTtcta	taagaatgtg	gcctatgtga	3540
acctcctttt	ctggtaccag	ttcttttgtg	gatttttcagg	gacatccatg	actgattact	3600
gggttttgat	cttcttcaac	ctcctcttca	catctgcccc	tcctgtcatt	tatgggtgtt	3660
tggagaaaaga	tgtgtctgca	gagaccctca	tgcaactgcc	tgaactttac	agaagtgggc	3720
agaaatcaga	ggcatactta	ccccatacct	tctggatcac	cttattggat	gctttttatc	3780
aaagcctggg	ctgcttcttt	gtgccttatt	ttacctacca	gggctcagat	actgacatct	3840
ttgcatttgg	aaacccccctg	aacacagccg	ctctgttcat	cgttctcctc	catctgggtca	3900
ttgaaagcaa	gagtttgact	tggattcact	tgctgggtcat	cattggtagc	atcttgtctt	3960
atTTTTtatt	tgccatagtt	tttggagcca	tgtgtgtaac	ttgcaacca	ccatccaacc	4020
cttactggat	tatgcaggag	cacatgctgg	atccagtatt	ctacttagtt	tgtatcctca	4080
cgacgtccat	tgctcttctg	cccaggtttg	tatacagagt	tcttcaggga	tccctgtttc	4140
catctccaat	tctgagagct	aagcactttg	acagactaac	tccagaggag	aggactaaag	4200
ctctcaagaa	gtggagaggg	gctggaaaaga	tgaatcaagt	gacatcaaag	tatgctaacc	4260
aatcagctgg	caagtcagga	agaagacca	tgcttggccc	ttctgctgta	tttgcaatga	4320
agtcagcaag	ttctgtgct	attgagcaag	gaaacttatc	tctgtgtgag	actgctttag	4380
atcaaggcta	ctctgaaact	aaggcctttg	agatggctgg	accctccaaa	ggtaaagaaa	4440
gctagatacc	ctccttggag	ttgcaagtat	tctttcaagg	ttggaagagg	gattttgaag	4500
aggtatctct	ccaagcaaga	atgacttggt	tttcataaag	ggacatgagc	atTTtactag	4560
gcttggaaga	gctgacatga	tgagcattat	tgtatgtttg	tatatacatt	tgtgatagag	4620
ggctagagtt	tgacctagag	agagtttaag	gaagtgaaat	atttaattca	gaaccaaagt	4680
cttttgtaaa	actttttgga	ttttgtaaaa	gcattttcat	tctcttagaa	attcaagtat	4740
tttcaagggg	agtcatttga	gatatattta	ttttactagg	agatcttata	ttctagggaa	4800
atgcttttaa	tggtcaggct	ccagtcggaa	tttttttaag	aaaaaagtag	tttttaatac	4860
attgggttagg	actcagagga	aatacggaaa	aaacattgta	gatgggttaat	ttacagataa	4920
aatcccaaga	gcctttttaa	caacaaggta	cctaaaatag	gggtataatta	tactgcttaa	4980
aatacagtta	gtgcctatta	atagcttttt	atttccatag	ggaagatgct	tttgggtctc	5040
tggctgaaga	tgtaggcata	cctctcactc	atttcaatgt	tttctgagg	tggagccttc	5100
attggaaaagg	ggaaagaggg	attctagggt	ttcatcaggg	accaggaatg	cattcctctg	5160
tcagggttcca	atcaagagaa	gaccttttat	gagatctgcc	tctgtataga	tgttgtcagt	5220

```

taggaaactg aagccatagg tcaggcagac atcagctcag cctgtggccc attgggtgat 5280
ttcctgtatt ttaaaactga cagtagcctg atcaaagtga tacaatcaat ttcaaaacaa 5340
tcttccagag accacttgaa ggttcatagt ttttacaata ccctgagact tttcaggtgt 5400
tggagcctct aaaatatgag atataaacag aaactaatac aagttgttct ctggagggtt 5460
ctatgaggtt cttagaaaaa tttggtttta aaatcatttg aggacaggaa tgtctatagc 5520
aagtttactc ctattgcgaa tcatgtatgc tggcttttagt tgtaacaaac gattttattc 5580
taagtaaggc caggtgctac tataaaatca tatattcctg ttgaagttct tttgaagtcc 5640
tatctctatt tatttatattt gaaagttgtc agccaccagt catccagaat ttccttcctg 5700
aatctccatg ctcatatgca atgtctacat caaggctctt ttaatgacta ttattctcag 5760
ggtttagttt tctaccttct gcctactatt ttggtctgac attttttagt ccttctgtta 5820
ttattggaaa tagtctctta cataagctga tttcgagaac tttcaaaatc tcacatagct 5880
aatggaagtt gctttctgct ttcttatgac tgtttttata aataaaactgt ttcataaata 5940
atgtatactg gaaaattgtc atcactcttg taaacatttg ccaaatacaa ctgtaattag 6000
taccagaact tagaaggaaa gtcacaaagc atgcctttta ccagaatatt gctttgcctt 6060
ttcttatatg gaaaatcact gatgtcatat cattttattat cctcacagca gctggcattc 6120
attccctgaa atcgtcgtct cctgataaac aataatttag agatatgttc acttgtaaatt 6180
aaaaaggtta gaatctaagg acaaagaatg atccaaaaat actaaggaaa aagaacaatt 6240
ttcaattttc agaaatgaat tatctgtaca tatgaataga ggaaaatact gagtgctaatt 6300
gttacagggc cacaatctca tatccacaat tccaaaatct ccaaacctct gaaaactgaa 6360
atatttttca taaattaggc gttaaattcta tgccctgaac aaatgtaaag ctatttatgt 6420
actttgctta atgtaatcat tcatatactt tgctacaaaa atattaatat atttcattac 6480
tgaatgctaa actgtatttc tttttaaaat ttggaaaatt ttttaattctg aaacacaatc 6540
tagaccttag 6550

```

```

<210> 12
<211> 1426
<212> PRT
<213> Homo sapiens

```

```

<400> 12

```

```

Met Thr Glu Ala Leu Gln Trp Ala Arg Tyr His Trp Arg Arg Leu Ile
1           5           10           15

```

```

Arg Gly Ala Thr Arg Asp Asp Asp Ser Gly Pro Tyr Asn Tyr Ser Ser
          20           25           30

```

```

Leu Leu Ala Cys Gly Arg Lys Ser Ser Gln Thr Pro Lys Leu Ser Gly

```

35					40					45					
Arg	His	Arg	Ile	Val	Val	Pro	His	Ile	Gln	Pro	Phe	Lys	Asp	Glu	Tyr
50					55					60					
Glu	Lys	Phe	Ser	Gly	Ala	Tyr	Val	Asn	Asn	Arg	Ile	Arg	Thr	Thr	Lys
65					70					75					80
Tyr	Thr	Leu	Leu	Asn	Phe	Val	Pro	Arg	Asn	Leu	Phe	Glu	Gln	Phe	His
				85					90					95	
Arg	Ala	Ala	Asn	Leu	Tyr	Phe	Leu	Phe	Leu	Val	Val	Leu	Asn	Trp	Val
			100					105					110		
Pro	Leu	Val	Glu	Ala	Phe	Gln	Lys	Glu	Ile	Thr	Met	Leu	Pro	Leu	Val
		115					120					125			
Val	Val	Leu	Thr	Ile	Ile	Ala	Ile	Lys	Asp	Gly	Leu	Glu	Gly	Tyr	Arg
	130					135					140				
Lys	Tyr	Lys	Ile	Asp	Lys	Gln	Ile	Asn	Asn	Leu	Ile	Thr	Lys	Val	Tyr
145					150					155					160
Ser	Arg	Lys	Glu	Lys	Lys	Tyr	Ile	Asp	Arg	Cys	Trp	Lys	Asp	Val	Thr
				165					170					175	
Val	Gly	Asp	Phe	Ile	Arg	Leu	Ser	Cys	Asn	Glu	Val	Ile	Pro	Ala	Asp
			180					185					190		
Met	Val	Leu	Leu	Phe	Ser	Thr	Asp	Pro	Asp	Gly	Ile	Cys	His	Ile	Glu
		195					200					205			
Thr	Ser	Gly	Leu	Asp	Gly	Glu	Ser	Asn	Leu	Lys	Gln	Arg	Gln	Val	Val
	210					215					220				
Arg	Gly	Tyr	Ala	Glu	Gln	Asp	Ser	Glu	Val	Asp	Pro	Glu	Lys	Phe	Ser
225					230					235					240
Ser	Arg	Ile	Glu	Cys	Glu	Ser	Pro	Asn	Asn	Asp	Leu	Ser	Arg	Phe	Arg
				245					250					255	
Gly	Phe	Leu	Glu	His	Ser	Asn	Lys	Glu	Arg	Val	Gly	Leu	Ser	Lys	Glu
			260					265				270			
Asn	Leu	Leu	Leu	Arg	Gly	Cys	Thr	Ile	Arg	Asn	Thr	Glu	Ala	Val	Val
		275					280					285			

Gly Ile Val Val Tyr Ala Gly His Glu Thr Lys Ala Met Leu Asn Asn
 290 295 300

Ser Gly Pro Arg Tyr Lys Arg Ser Lys Leu Glu Arg Arg Ala Asn Thr
 305 310 315 320

Asp Val Leu Trp Cys Val Met Leu Leu Val Ile Met Cys Leu Thr Gly
 325 330 335

Ala Val Gly His Gly Ile Trp Leu Ser Arg Tyr Glu Lys Met His Phe
 340 345 350

Phe Asn Val Pro Glu Pro Asp Gly His Ile Ile Ser Pro Leu Leu Ala
 355 360 365

Gly Phe Tyr Met Phe Trp Thr Met Ile Ile Leu Leu Gln Val Leu Ile
 370 375 380

Pro Ile Ser Leu Tyr Val Ser Ile Glu Ile Val Lys Leu Gly Gln Ile
 385 390 395 400

Tyr Phe Ile Gln Ser Asp Val Asp Phe Tyr Asn Glu Lys Met Asp Ser
 405 410 415

Ile Val Gln Cys Arg Ala Leu Asn Ile Ala Glu Asp Leu Gly Gln Ile
 420 425 430

Gln Tyr Leu Phe Ser Asp Lys Thr Gly Thr Leu Thr Glu Asn Lys Met
 435 440 445

Val Phe Arg Arg Cys Ser Val Ala Gly Phe Asp Tyr Cys His Glu Glu
 450 455 460

Asn Ala Arg Arg Leu Glu Ser Tyr Gln Glu Ala Val Ser Glu Asp Glu
 465 470 475 480

Asp Phe Ile Asp Thr Val Ser Gly Ser Leu Ser Asn Met Ala Lys Pro
 485 490 495

Arg Ala Pro Ser Cys Arg Thr Val His Asn Gly Pro Leu Gly Asn Lys
 500 505 510

Pro Ser Asn His Leu Ala Gly Ser Ser Phe Thr Leu Gly Ser Gly Glu
 515 520 525

Gly Ala Ser Glu Val Pro His Ser Arg Gln Ala Ala Phe Ser Ser Pro
 530 535 540

Ile Glu Thr Asp Val Val Pro Asp Thr Arg Leu Leu Asp Lys Phe Ser
545 550 555 560

Gln Ile Thr Pro Arg Leu Phe Met Pro Leu Asp Glu Thr Ile Gln Asn
565 570 575

Pro Pro Met Glu Thr Leu Tyr Ile Ile Asp Phe Phe Ile Ala Leu Ala
580 585 590

Ile Cys Asn Thr Val Val Val Ser Ala Pro Asn Gln Pro Arg Gln Lys
595 600 605

Ile Arg His Pro Ser Leu Gly Gly Leu Pro Ile Lys Ser Leu Glu Glu
610 615 620

Ile Lys Ser Leu Phe Gln Arg Trp Ser Val Arg Arg Ser Ser Ser Pro
625 630 635 640

Ser Leu Asn Ser Gly Lys Glu Pro Ser Ser Gly Val Pro Asn Ala Phe
645 650 655

Val Ser Arg Leu Pro Leu Phe Ser Arg Met Lys Pro Ala Ser Pro Val
660 665 670

Glu Glu Glu Val Ser Gln Val Cys Glu Ser Pro Gln Cys Ser Ser Ser
675 680 685

Ser Ala Cys Cys Thr Glu Thr Glu Lys Gln His Gly Asp Ala Gly Leu
690 695 700

Leu Asn Gly Lys Ala Glu Ser Leu Pro Gly Gln Pro Leu Ala Cys Asn
705 710 715 720

Leu Cys Tyr Glu Ala Glu Ser Pro Asp Glu Ala Ala Leu Val Tyr Ala
725 730 735

Ala Arg Ala Tyr Gln Cys Thr Leu Arg Ser Arg Thr Pro Glu Gln Val
740 745 750

Met Val Asp Phe Ala Ala Leu Gly Pro Leu Thr Phe Gln Leu Leu His
755 760 765

Ile Leu Pro Phe Asp Ser Val Arg Lys Arg Met Ser Val Val Val Arg
770 775 780

His Pro Leu Ser Asn Gln Val Val Val Tyr Thr Lys Gly Ala Asp Ser
785 790 795 800

Val Ile Met Glu Leu Leu Ser Val Ala Ser Pro Asp Gly Ala Ser Leu
 805 810 815

Glu Lys Gln Gln Met Ile Val Arg Glu Lys Thr Gln Lys His Leu Asp
 820 825 830

Asp Tyr Ala Lys Gln Gly Leu Arg Thr Leu Cys Ile Ala Lys Lys Val
 835 840 845

Met Ser Asp Thr Glu Tyr Ala Glu Trp Leu Arg Asn His Phe Leu Ala
 850 855 860

Glu Thr Ser Ile Asp Asn Arg Glu Glu Leu Leu Leu Glu Ser Ala Met
 865 870 875 880

Arg Leu Glu Asn Lys Leu Thr Leu Leu Gly Ala Thr Gly Ile Glu Asp
 885 890 895

Arg Leu Gln Glu Gly Val Pro Glu Ser Ile Glu Ala Leu His Lys Ala
 900 905 910

Gly Ile Lys Ile Trp Met Leu Thr Gly Asp Lys Gln Glu Thr Ala Val
 915 920 925

Asn Ile Ala Tyr Ala Cys Lys Leu Leu Glu Pro Asp Asp Lys Leu Phe
 930 935 940

Ile Leu Asn Thr Gln Ser Lys Asp Ala Cys Gly Met Leu Met Ser Thr
 945 950 955 960

Ile Leu Lys Glu Leu Gln Lys Lys Thr Gln Ala Leu Pro Glu Gln Val
 965 970 975

Ser Leu Ser Glu Asp Leu Leu Gln Pro Pro Val Pro Arg Asp Ser Gly
 980 985 990

Leu Arg Ala Gly Leu Ile Ile Thr Gly Arg Thr Leu Glu Phe Ala Leu
 995 1000 1005

Gln Glu Ser Leu Gln Lys Gln Phe Leu Glu Leu Thr Ser Trp Cys
 1010 1015 1020

Gln Ala Val Val Cys Cys Arg Ala Thr Pro Leu Gln Lys Ser Glu
 1025 1030 1035

Val Val Lys Leu Val Arg Ser His Leu Gln Val Met Thr Leu Ala
 1040 1045 1050

Ile Gly 1055	Asp Gly	Ala Asn	Asp	Val Ser Met	Ile Gln 1065	Val Ala Asp
Ile Gly 1070	Ile Gly Val	Ser Gly 1075	Gln Glu Gly Met	Gln 1080	Ala Val Met	
Ala Ser 1085	Asp Phe Ala Val	Tyr 1090	Gln Phe Lys His	Leu 1095	Ser Lys Leu	
Leu Leu 1100	Val His Gly His	Trp 1105	Cys Tyr Thr Arg	Leu 1110	Ser Asn Met	
Ile Leu 1115	Tyr Phe Phe Tyr	Lys 1120	Asn Val Ala Tyr	Val 1125	Asn Leu Leu	
Phe Trp 1130	Tyr Gln Phe Phe	Cys 1135	Gly Phe Ser Gly	Thr 1140	Ser Met Thr	
Asp Tyr 1145	Trp Val Leu Ile	Phe 1150	Phe Asn Leu Leu	Phe 1155	Thr Ser Ala	
Pro Pro 1160	Val Ile Tyr Gly	Val 1165	Leu Glu Lys Asp	Val 1170	Ser Ala Glu	
Thr Leu 1175	Met Gln Leu Pro	Glu 1180	Leu Tyr Arg Ser	Gly 1185	Gln Lys Ser	
Glu Ala 1190	Tyr Leu Pro His	Thr 1195	Phe Trp Ile Thr	Leu 1200	Leu Asp Ala	
Phe Tyr 1205	Gln Ser Leu Val	Cys 1210	Phe Phe Val Pro	Tyr 1215	Phe Thr Tyr	
Gln Gly 1220	Ser Asp Thr Asp	Ile 1225	Phe Ala Phe Gly	Asn 1230	Pro Leu Asn	
Thr Ala 1235	Ala Leu Phe Ile	Val 1240	Leu Leu His Leu	Val 1245	Ile Glu Ser	
Lys Ser 1250	Leu Thr Trp Ile	His 1255	Leu Leu Val Ile	Ile 1260	Gly Ser Ile	
Leu Ser 1265	Tyr Phe Leu Phe	Ala 1270	Ile Val Phe Gly	Ala 1275	Met Cys Val	
Thr Cys 1280	Asn Pro Pro Ser	Asn 1285	Pro Tyr Trp Ile	Met 1290	Gln Glu His	

Met Leu Asp Pro Val Phe Tyr Leu Val Cys Ile Leu Thr Thr Ser
1295 1300 1305

Ile Ala Leu Leu Pro Arg Phe Val Tyr Arg Val Leu Gln Gly Ser
1310 1315 1320

Leu Phe Pro Ser Pro Ile Leu Arg Ala Lys His Phe Asp Arg Leu
1325 1330 1335

Thr Pro Glu Glu Arg Thr Lys Ala Leu Lys Lys Trp Arg Gly Ala
1340 1345 1350

Gly Lys Met Asn Gln Val Thr Ser Lys Tyr Ala Asn Gln Ser Ala
1355 1360 1365

Gly Lys Ser Gly Arg Arg Pro Met Pro Gly Pro Ser Ala Val Phe
1370 1375 1380

Ala Met Lys Ser Ala Ser Ser Cys Ala Ile Glu Gln Gly Asn Leu
1385 1390 1395

Ser Leu Cys Glu Thr Ala Leu Asp Gln Gly Tyr Ser Glu Thr Lys
1400 1405 1410

Ala Phe Glu Met Ala Gly Pro Ser Lys Gly Lys Glu Ser
1415 1420 1425

<210> 13
<211> 8083
<212> DNA
<213> Homo sapiens

<400> 13
ctcgggcagc ccagtccttg ccaccccttg ccagccggtg tgggtgcttgt gtgtcacagc 60
cttgtagccg ggagtcgctg ccgagtgggc gctcagtttt cgggtcgtca tggctggcta 120
cgaatacgtg agcccgagc agctggctgg ctttgataag tacaagtaca gtgctgtgga 180
taccaatcca ctttctctgt atgtcatgca tccattctgg aacactatag taaaggattt 240
tcctacttgg ctggcgccca atctgataac ttttctggc tttctgctgg tcgtattcaa 300
tttctgcta atggcatact ttgatcctga cttttatgcc tcagcaccag gtcacaagca 360
cgtgcctgac tgggtttgga ttgtagtggg catcctcaac ttcgtagcct aactctaga 420
tgggtgtggac ggaaagcaag ctgcgagaac caattctagc actcccttag gggagctttt 480
tgatcatggc ctggatagtt ggtcatgtgt ttactttgtt gtgactgttt attccatctt 540
tggaagagga tcaactggtg tcagtgtttt tgttctttat ctctgctat gggtagtttt 600
gttttctttc atcctgtccc actgggaaaa gtataacaca gggattcttt tcctgccatg 660

gggatatgac	attagccagg	tgactatttc	ttttgtctac	atagtgactg	cagttgtggg	720
agttgaggcc	tggtatgaac	ctttcctggt	taattttotta	tatagagacc	tattcactgc	780
aatgattatt	ggttgtgcat	tatgtgtgac	tcttccaatg	agtttattaa	actttttcag	840
aagctataaa	aataacacct	tgaaactcaa	ttcagtctat	gaagctatgg	ttcccttatt	900
ttctccatgc	ttgctgttca	ttttgtctac	agcgtggatc	ctttggtcac	cttcagatat	960
tttagagcta	catcctagag	tattctactt	tatggttgga	acagcttttg	ccaacagtac	1020
atgtcagctg	attgtttgcc	aatgagtag	taccogggtg	ccaactttga	attggttgct	1080
ggttcctctc	ttcttggttg	tcttagtggt	aaacctagga	gtagcctctt	acgttgagag	1140
cattctcctg	tatacattaa	caactgcttt	tactctggcc	cacatccatt	atggagtacg	1200
agtggtaaag	cagctgagca	gccattttca	gattttacccc	ttctcattga	ggaaaccaa	1260
ctcagattga	ctaggaatgg	aagaaaagaa	tattggcctg	taataatctt	tctttgggca	1320
caaagaagta	ctgtaaataa	atgcttgtaa	atatttctctc	catcaccatt	gaactagact	1380
gatctgcttg	acagacgtgg	gatctcagta	tggtacttgg	acagcaggaa	tgatacatat	1440
aatctgaact	tgggaaattt	tggacctact	aactctaagc	ctattttatt	ttttataaaa	1500
ctatgtgaca	ttttggttga	gcagaatgta	cgttagacca	gcaaaatggt	cctaattggt	1560
ctaacttcgt	gagtttacia	tgttgtgatt	catgcagggt	taaagatgct	ttgtttttat	1620
tttttaagta	ccaaaattgg	tttcagaaca	ctgataaacac	tcagaaaacc	acagtgtggt	1680
ttcatatttg	gaactttgta	atagcgggag	tagcagtagt	ccaaacctag	tatagggaaa	1740
ggataaaaat	aagtcacctt	caccaagaga	tgccaatgat	taccaaacaa	cagacagttg	1800
ccaaatactg	gtttctcttt	cccctgaaaa	tggcttttgt	tctcaaataga	taagagagct	1860
aatacattta	gctaataattc	tagctctctt	tattatggaa	cagatcttga	tagatggttt	1920
aattttctcc	taaagagaaa	taatcagttg	agaatttgag	aatgggttgt	aattatcgct	1980
caccatttgg	gatggttcat	tgtttaata	tggcattttc	cccccttcag	ctgcaggttc	2040
ctgagatttg	gtgcctgtga	gctctgattg	taggaatgca	tgtgacagtc	ccagtcctat	2100
ggtaatgact	taggaggaat	gcagataaaa	gtaccttgta	agataaatat	aaattggagt	2160
taggaatttc	atgaacctca	ctatgaccaa	aattaatttt	ttgattcagt	ttgtctgtct	2220
gtctgtcctt	cccctctctt	cttttttcag	ggtgaggtgc	tgtgtttctt	atttcatacg	2280
agataaaaaca	gagagaagtt	ctctcttctc	cagcttgtcc	atttccccac	ttgaagaaaa	2340
cttttgatat	atatgcctta	ctgagtacat	gcccccttta	atgttaacat	gacttggagt	2400
aattttctgag	gtttactgac	aaacataaaa	atccccctta	ttgtagtgta	gttgttctat	2460
aaaccatatt	ttttcatgat	gtggatattt	tcttctattt	ctttgtcttc	atttaatttg	2520
gtgggtggtga	actttacttg	ctgattttct	tttatttttc	actgaatgaa	gtttgtgctt	2580

gaatgaagag	tgtatcttaa	acccccctttt	tttggacagg	ctgcacttgg	ataaaaatagg	2640
caccactgtg	ttgatatgta	attaaattca	taactattta	tgaatgtgac	cattgttaaa	2700
tttaattata	gtgtatttct	ttcttgaatt	tcagtctgct	gctttgtaaa	aaataactgt	2760
aattgatatt	tcatttttta	aattttataaa	ttccaatcat	tgtcattcaa	gttattagaa	2820
atgagtcacg	gcactaactc	atggttctat	actcatttta	agaaattgta	aagggtgagg	2880
tgtatttggg	gactttctct	gagacacaga	tgaacgttgg	aaatagatga	gaaatggatg	2940
cagaataatg	cagatagtga	tcccatgttt	gttaattgcc	atcatggaag	tcagagaatt	3000
aaccaattaa	gtgacggtaa	ctattttttg	tataccttgt	tctttgtgat	ctataaaatt	3060
aaactggata	aaatgttaat	tgggggtagg	gaaaaagctt	tgttttaaga	tttctaagct	3120
aattatttta	cogttttgtt	gacctgttac	acaagaatga	gattaatctg	tattatataa	3180
gaaaagaact	gataaatatt	tataaaatga	caaacttgga	ccacaggctt	tcttgacctt	3240
gtacagtata	gtaaaaatac	cctttacttc	tgacatttta	aggtcacgtc	ctatagtttg	3300
ggggcccttt	tacttcaaag	gtgtgtttct	gccctctttt	ggttgatatca	aatttatttg	3360
aaaattgttt	tgtgctgtac	tttatgactt	gtttccctat	ggtggtaaag	aatgatgcca	3420
tagtcagttt	tattgacatt	gtagatgaa	agaaaagtag	ctgtcaacaa	ggggaaagtg	3480
aaattgagca	attcactgtt	tcaggatact	tgaggatctt	tagtgtactg	ctgactcttt	3540
ttcttgtttt	gtctgccatt	cttgaaagtg	catcttgaga	gccccatttt	gttgtgtaga	3600
gatatagcag	ctgaagcagg	acttcattcc	agatagctgt	gtgcatttat	gtcatgccaa	3660
ttactataac	atttccaaag	tatgaaaggt	ctttcaggct	tatgcatgat	attgtgggtg	3720
acttcatgcc	ccagcctttc	tccccagcat	acagtgagct	gccttttggg	gagcaatgcc	3780
atgataagga	gcatacgtgc	atgttttccg	tactttcaga	atagagtgcc	ttagctaggc	3840
cagaggctcg	tattgtgttt	ctgctgacca	aagttgcata	cacctgtaag	aagtgtcttt	3900
ttaggttatg	gtattaaaaa	agtcgtttca	ggcttatcag	ataataattc	tcctatatatt	3960
gaggggggca	tgggataaaa	agatttaata	atacctggat	agagaatgct	aatgctgcct	4020
tttgtgatcc	tttaatatata	tccatgcttt	tataaagtgc	agtacacatt	ttgacattgc	4080
cacagtttac	agtgtcattc	ttccacgaat	aggtaattga	tagagtttat	gggtcttagt	4140
cttaagggtg	acagagtga	taccttccaa	ataaaaagct	aaatattcct	ccttgtcatt	4200
ttaatgtcat	tattgaaata	cttttaattc	taattttcaa	aagaaagcta	aatgctgatt	4260
gtgggttagg	atgttacagt	actaatacca	tcattgttatt	tattctaata	tagaggcatt	4320
gagaggaggg	gagtgtatgt	ctagagttac	cccaattcta	aataggtaacc	aaaagagtat	4380
tgattatgac	ctggagcctg	agctttggga	gaatatacaa	aggaaaaagc	aggcatagtt	4440
tccactttta	agggaagaag	ggactttaac	atcctatgaa	tttgaaacat	cctatgaatt	4500

```

tgaaagtttc ctaagtgttg ttttcaaaga actggtttat tttatagtca gtcaaaatgc 4560
at ttggaatg taccttgctt ccataacaga ctttgactct atgacctatg aagaataaaa 4620
agataatctg aagggtgttg agtcagaggg tacttttccc ctttatattt tatatagaag 4680
cctatttgta attcttcttc ttctgtcttc acctgggttt cctcttatct acccaccacc 4740
taggacagag gctgccact cagggtcttg cttgttgccc ctagatgttg ccacattcac 4800
atctccttga tggccttaga tctcttgact agtcaaggac tgggaacgtg ccagggggaa 4860
caaagtaagc ttgatgggct ggaacttttg actttgggtc cccaaacagc caaaaacaga 4920
tacagaaatc tttggtgttg tggatgcctg tgttcagagt ggccttatct tgacctcagt 4980
tctggattta agtactctat agaaaatttg tgaagtcaga ccagaatttt ctttgtccca 5040
tgagggtgtc ttgtaagtca gcatgctctt gcttgaacac cctgtgggtg aataacttta 5100
gtaagttagc tgccagcatt agaaccttcc aggaagaatt ttaactagta tgctacattt 5160
gaagcactct gacatgtaca tttaaagtga cagttttaaa agtaaaatgg tgttaaactc 5220
ctgaaacttg aacgtatgat gctctggaca ttttaggaag ctttaagtaa tttctgattt 5280
tcctttattg attatgtaac ccacgtgatt tgcatatttt agtttttatg tttgacagtg 5340
gtacaaatga ctgacattgg caggcagtta ttgtagggta tatggttaat cacgattgtc 5400
taacttttcc agagaaagtg gtgatatgct tcttccttgc atggagctgt gggaattgga 5460
tttgttggct cttcatttcc agtgtttcta cagtgcagaa taaatgtaac tttttaagcc 5520
tgcatgctag taatatgctt tgctctttag taaacaaata tgcaatagat agtaatctat 5580
ctactttttt taaagcaggg atctgctccc tatctagaca gatcttttaa tgcattaata 5640
tttagagtag caaaagaagt tttgccattt gtagctctct cacaggtagc ttaagcgagg 5700
gggttggtgt gcatgtgctt agaaaggcca ggatcccaga attggaagggt gccacatata 5760
cttttagaat attttaacaa aggcttgtct ctaggaaaat ccacttggac tgcagcagga 5820
ttgttctggt cttccacaga ccattcacat ggtctttctc tccagagaat cattcttaga 5880
aagccaggct aaatcagagt agtgctgttt ttaaagcaag tatattggat gatggctata 5940
gggtacctta atgctgtgtt gaaattggaa gcagactgct cttcagcagc acatatactg 6000
caactgaagt tagagacagt actgaagacg agggagtagc taaatgctgt ggatatttca 6060
tgtctgtttg gctaattgtc tctcttgagg agtttctatg ggtagcaatt aaaattagaa 6120
tttattttaa ggtttttatt aaaatatata caatttgcca taaattctag attctgatat 6180
tcttgacaag cactttttga atatctcaag agtaagttct tgacctggaa catatatagt 6240
tagtacttga tgcactaac caagttgtag aaggagcctg ttataattct gcaagatctg 6300
tgaatagcat tataaatctg gggcaggtgc ctgtaatccc agcactttgg gaagccaagg 6360
cagggtggatt gcttgagccc aggagttcga caccaacctg agcaacatgg cgaaaccccg 6420

```

```

tctgtaataa aaatacaaaa aaattagctg ggcattggtgg catgtgcctg taatcccagt 6480
tactggggag gctgaggcag gagaatcact tgaacctggg agatggagat tgcagtgagc 6540
agagattgtg cactgcact ccagccagag tgacagagca agactccatc tcaaaaaata 6600
ataataaaaa taaacaaatc tgttttaaatt actgttgctt ttcagcatta gactttgagt 6660
ttaataacta caaattgaga ctgctcacia tattaacttt tttttaggtt tattttgttg 6720
tttagataac ttgcctccct agaggagcat tcaggagata aagacctagc tacatgtaat 6780
gatatgatca tttcaaaaat gtgccaaaga agcaaaattc attattgaac tctaaatttc 6840
tgggtgtttt ttttttttaa gtagcatttt ctctgggtaa agggaaaggc aatgaatgat 6900
tgccaacatg taaactccct gctgcccgcc ttccccagt cccctccatc taacataata 6960
cagtaaatth gtagccagtt gtagaaaaag aaattgatat ctttctgagt aaggtttcat 7020
gccctgtgac taagaatagg tggaggaatc atggccaaat caaatatgaa ttgttttagct 7080
tggctcctgt gcagttggct ttctgtagtg ttctgaaaca agaggacgat tccattcctt 7140
ctcaggaacc tagaaacaac accgcttgag caactggaat atgtttgctg caagcagaat 7200
atthttggga gaggaagagt agtttaattc aagtagttta attgaacata ttagtcattg 7260
gtctgtctgg gacgtgcagt gttcatagta gcaatgtatg taccatttat tttatctgg 7320
tgtgtgtgat gtgtgtgtat gcctatttaa tatgtacaca tatattcact acatatgtat 7380
gtatgatata ttcatatata catgcagtct gcttgattat cagcaaaatg gtcagccttt 7440
atcagatagt ttcttcatgt ggagttcatc tgcattgtgg ccttactctg aagcctcttc 7500
ctgatctgga gccacagtct gtctgtcttc cagttcatct cagtcctcga gaaaggccct 7560
ttaaatatgt cactttccca ttttctttta accatgggtt gtgtgagcca gaaagagctt 7620
tgagaaagat ggctgcttcc accaggggtg aggttcttag gtctgcatga tgatggggcc 7680
cgthttctgg cagaggggtg ctctgggagc agttgtgctg cgggcttgct gggggagaac 7740
tctaactgtt gcagaaacag agcttcatgg cttgcttaaa ttacttagct ggaatatttt 7800
aaagtgtcag ataattgtat gtacaaagag agtatgccga tgcatttcat tgtttttgca 7860
ttatctgtat cagtgtttcc cattgctagc ccgaataatc aagaccagac tgtttcttgt 7920
cactggattc tggaagcctt gccctttcaa ctggatttat gtttgthttc gctgctgtga 7980
aaccctaaag taatgcagta ggttttaatt gaaaattgtt taattctatt attgataatt 8040
atthattgta ataaaagtaa aagagtaatt tttaaatcct ttt 8083

```

<210> 14
<211> 386
<212> PRT
<213> Homo sapiens

<400> 14

Met Ala Gly Tyr Glu Tyr Val Ser Pro Glu Gln Leu Ala Gly Phe Asp
 1 5 10 15
 Lys Tyr Lys Tyr Ser Ala Val Asp Thr Asn Pro Leu Ser Leu Tyr Val
 20 25 30
 Met His Pro Phe Trp Asn Thr Ile Val Lys Val Phe Pro Thr Trp Leu
 35 40 45
 Ala Pro Asn Leu Ile Thr Phe Ser Gly Phe Leu Leu Val Val Phe Asn
 50 55 60
 Phe Leu Leu Met Ala Tyr Phe Asp Pro Asp Phe Tyr Ala Ser Ala Pro
 65 70 75 80
 Gly His Lys His Val Pro Asp Trp Val Trp Ile Val Val Gly Ile Leu
 85 90 95
 Asn Phe Val Ala Tyr Thr Leu Asp Gly Val Asp Gly Lys Gln Ala Arg
 100 105 110
 Arg Thr Asn Ser Ser Thr Pro Leu Gly Glu Leu Phe Asp His Gly Leu
 115 120 125
 Asp Ser Trp Ser Cys Val Tyr Phe Val Val Thr Val Tyr Ser Ile Phe
 130 135 140
 Gly Arg Gly Ser Thr Gly Val Ser Val Phe Val Leu Tyr Leu Leu Leu
 145 150 155 160
 Trp Val Val Leu Phe Ser Phe Ile Leu Ser His Trp Glu Lys Tyr Asn
 165 170 175
 Thr Gly Ile Leu Phe Leu Pro Trp Gly Tyr Asp Ile Ser Gln Val Thr
 180 185 190
 Ile Ser Phe Val Tyr Ile Val Thr Ala Val Val Gly Val Glu Ala Trp
 195 200 205
 Tyr Glu Pro Phe Leu Phe Asn Phe Leu Tyr Arg Asp Leu Phe Thr Ala
 210 215 220
 Met Ile Ile Gly Cys Ala Leu Cys Val Thr Leu Pro Met Ser Leu Leu
 225 230 235 240
 Asn Phe Phe Arg Ser Tyr Lys Asn Asn Thr Leu Lys Leu Asn Ser Val
 245 250 255

Tyr Glu Ala Met Val Pro Leu Phe Ser Pro Cys Leu Leu Phe Ile Leu
 260 265 270

Ser Thr Ala Trp Ile Leu Trp Ser Pro Ser Asp Ile Leu Glu Leu His
 275 280 285

Pro Arg Val Phe Tyr Phe Met Val Gly Thr Ala Phe Ala Asn Ser Thr
 290 295 300

Cys Gln Leu Ile Val Cys Gln Met Ser Ser Thr Arg Cys Pro Thr Leu
 305 310 315 320

Asn Trp Leu Leu Val Pro Leu Phe Leu Val Val Leu Val Val Asn Leu
 325 330 335

Gly Val Ala Ser Tyr Val Glu Ser Ile Leu Leu Tyr Thr Leu Thr Thr
 340 345 350

Ala Phe Thr Leu Ala His Ile His Tyr Gly Val Arg Val Val Lys Gln
 355 360 365

Leu Ser Ser His Phe Gln Ile Tyr Pro Phe Ser Leu Arg Lys Pro Asn
 370 375 380

Ser Asp
 385

<210> 15
 <211> 2788
 <212> DNA
 <213> Homo sapiens

<400> 15
 atagaaggaa gaggagcgaa catggcagcg cgttggcggt tttggtgtgt ctctgtgacc 60
 atggtggtgg cgctgctcat cgtttgcgac gttccctcag cctctgccaa aagaaagaag 120
 gagatggtgt tatctgaaaa ggtagtcag ctgatggaat ggactaaca aagacctgta 180
 ataagaatga atggagacaa gttcgcgc cttgtgaaag cccaccgag aaattactcc 240
 gttatcgta tgttactgc tctccaactg catagacagt gtgtcgttg caagcaagct 300
 gatgaagaat tccagatcct ggcaaactcc tggcgatact ccagtgcatt caccaacagg 360
 atatTTTTTg ccatgggtgga ttttgatgaa ggctctgatg tatttcagat gctaaacatg 420
 aattcagctc caactttcat caactttcct gcaaaagga aacccaaacg gggatgataca 480
 tatgagttac aggtgcgggg tttttcagct gagcagattg cccgggtggat cgccgacaga 540
 actgatgtca atattagagt gattagaccc ccaaattatg ctggtcccct tatgttggga 600
 ttgcttttgg ctgttattgg tggacttgtg tatcttcgaa gaagtaatat ggaatttctc 660

tttaataaaaa	ctggatgggc	ttttgcagct	ttgtgttttg	tgcttgctat	gacatctggg	720
caaagtgtga	accatataag	aggaccacca	tatgcccata	agaatcccca	cacgggacat	780
gtgaattata	tccatggaag	cagtcaagcc	cagttttag	ctgaaacaca	cattgttctt	840
ctgtttaatg	gtggagttac	cttaggaatg	gtgcttttat	gtgaagctgc	tacctctgac	900
atggatattg	gaaagcgaaa	gataatgtgt	gtggctggta	ttggacttgt	tgtattattc	960
ttcagttgga	tgctctctat	ttttagatct	aaatatcatg	gctaccata	cagctttctg	1020
atgagttaaa	aagggtcccag	agatatatag	acactggagt	actggaaatt	gaaaaacgaa	1080
aatcgtgtgt	gtttgaaaag	aagaatgcaa	cttgtatatt	ttgtattacc	tctttttttc	1140
aagtgattta	aatagttaat	catttaacca	aagaagatgt	gtagtgcctt	aacaagcaat	1200
cctctgtcaa	aatctgaggt	at ttgaaaat	aattatcctc	ttaaccttct	cttcccagtg	1260
aactttatgg	aacatttaat	ttagtacaat	taagtatatt	ataaaaattg	taaaactact	1320
actttgtttt	agttagaaca	aagctcaaaa	ctactttagt	taacttggtc	atctgatttt	1380
atattgcctt	atccaaagat	ggggaaagta	agtcctgacc	agggtgtccc	acatatgcct	1440
gttacagata	actacattag	gaattcattc	ttagcttctt	catctttgtg	tggatgtgta	1500
tactttacgc	atcttttctt	ttgagtagag	aaattatgtg	tgtcatgtgg	tcttctgaaa	1560
atggaacacc	attcttcaga	gcacacgtct	agccctcagc	aagacagttg	tttctcctcc	1620
tccttgcata	tttctactg	aaatacagtg	ctgtctatga	ttgtttttgt	tttgttgttt	1680
ttttgagacg	gtctcgctgt	gtcacacagg	ctggagtgca	gtggcgtgag	ctcggctgac	1740
tgcaaactct	gcctcccagg	tttaagcgat	tctcctgtca	cagcttccca	agtagctggg	1800
at ttacaggt	gtgcaccgcc	atgccaggct	aatttttgtg	tttttagtag	agacagggtt	1860
tcgccaaagt	gtccaggctg	gtcttgaact	cctgggctca	agtgatccgc	ccgcctcagt	1920
ctcccaaagt	gcgaggatga	catgtgtgag	ctaccacacc	agcaatgtct	atgcttctcg	1980
atagctgtga	acatgaaaag	acatctattg	ggagtccgag	gcagggtggat	tgcttgaggc	2040
caggagttag	agaccagcct	ggccaacaag	gcaaaacccc	gtctctacta	aaaatatgaa	2100
aattagctgg	gcttggtggc	tcatgcctat	aatcctagct	acttgggagg	ctgaggcacg	2160
agacttgctt	aatacctggg	aggcggagat	tgcagtgagc	cgagatcacg	ctactgcgct	2220
ccagcctgag	tgatagagtg	agactctgtc	tcaaaaaaaaa	gtatctctaa	atacaggatt	2280
ataatttctg	cttgagtatg	gtgttaacta	ccttgtattt	agaaagattt	cagattcatt	2340
ccatctcctt	agttttcttt	taaggtagacc	catctgtgat	aaaaatatag	cttagtgcta	2400
aaatcagtg	aacttataca	tggcctaata	tgtttctaca	aattagagtt	tgtcacttat	2460
tccatttgta	cctaagagaa	aaataggctc	agttagaaaa	ggactccctg	gccaggcgca	2520
gtgacttacg	cctgtaatct	cagcactttg	ggaggccaag	gcaggcagat	cacgagggtca	2580

ggagttcgag accatcctgg ccaacatggt gaaaccccgt ctctactaaa aatataaaaa 2640
 ttagctgggt gtggtggcag gagcctgtaa tcccagctac acaggagggt gaggcacgag 2700
 aatcacttga actcaggaga tggaggtttc agtgagccaa gatcacacca ctgcactcca 2760
 gcctgggaac agagcgagac tccatctc 2788

<210> 16
 <211> 335
 <212> PRT
 <213> Homo sapiens

<400> 16

Met Ala Ala Arg Trp Arg Phe Trp Cys Val Ser Val Thr Met Val Val
 1 5 10 15

Ala Leu Leu Ile Val Cys Asp Val Pro Ser Ala Ser Ala Lys Arg Lys
 20 25 30

Lys Glu Met Val Leu Ser Glu Lys Val Ser Gln Leu Met Glu Trp Thr
 35 40 45

Asn Lys Arg Pro Val Ile Arg Met Asn Gly Asp Lys Phe Arg Arg Leu
 50 55 60

Val Lys Ala Pro Pro Arg Asn Tyr Ser Val Ile Val Met Phe Thr Ala
 65 70 75 80

Leu Gln Leu His Arg Gln Cys Val Val Cys Lys Gln Ala Asp Glu Glu
 85 90 95

Phe Gln Ile Leu Ala Asn Ser Trp Arg Tyr Ser Ser Ala Phe Thr Asn
 100 105 110

Arg Ile Phe Phe Ala Met Val Asp Phe Asp Glu Gly Ser Asp Val Phe
 115 120 125

Gln Met Leu Asn Met Asn Ser Ala Pro Thr Phe Ile Asn Phe Pro Ala
 130 135 140

Lys Gly Lys Pro Lys Arg Gly Asp Thr Tyr Glu Leu Gln Val Arg Gly
 145 150 155 160

Phe Ser Ala Glu Gln Ile Ala Arg Trp Ile Ala Asp Arg Thr Asp Val
 165 170 175

Asn Ile Arg Val Ile Arg Pro Pro Asn Tyr Ala Gly Pro Leu Met Leu
 180 185 190

Gly Leu Leu Leu Ala Val Ile Gly Gly Leu Val Tyr Leu Arg Arg Ser
 195 200 205

Asn Met Glu Phe Leu Phe Asn Lys Thr Gly Trp Ala Phe Ala Ala Leu
 210 215 220

Cys Phe Val Leu Ala Met Thr Ser Gly Gln Met Trp Asn His Ile Arg
 225 230 235 240

Gly Pro Pro Tyr Ala His Lys Asn Pro His Thr Gly His Val Asn Tyr
 245 250 255

Ile His Gly Ser Ser Gln Ala Gln Phe Val Ala Glu Thr His Ile Val
 260 265 270

Leu Leu Phe Asn Gly Gly Val Thr Leu Gly Met Val Leu Leu Cys Glu
 275 280 285

Ala Ala Thr Ser Asp Met Asp Ile Gly Lys Arg Lys Ile Met Cys Val
 290 295 300

Ala Gly Ile Gly Leu Val Val Leu Phe Phe Ser Trp Met Leu Ser Ile
 305 310 315 320

Phe Arg Ser Lys Tyr His Gly Tyr Pro Tyr Ser Phe Leu Met Ser
 325 330 335

<210> 17
 <211> 3239
 <212> DNA
 <213> Homo sapiens

<400> 17
 gcttggggaa cgaggagcag gacgcggcct cggtggggcc cgggccgaac ggctgctggac 60
 acctgggctgc cgaggagccg agcgccgccc tctccggcat ggatcagtgc gtgacgggtg 120
 agcgcgagct ggagaaggtg ctgcacaagt tctcaggcta cgggcagctg tgcgagcgcg 180
 gcctggagga gctcatcgac tacaccggcg gcctcaagca cgagatcctg cagagccacg 240
 gccaaagtgc tgaattatca gggacacttt cacttggttt gacacagtgc tgtaaaagaa 300
 taaaggatac tgttcaaaaa ttggcctccg accacaaaga catccacagc agtgtttctc 360
 gggttggaaa agccattgat aagaattttg attctgacat tagcagtgtg ggaatagatg 420
 gctgctggca ggcagacagc caaaggcttc tcaatgaagt gatggtggag cacttctttc 480
 gacaaggaat gctggatgtg gctgaggagc tctgtcagga atctggtctt tctgtagacc 540
 caagtcagaa ggaaccattt gtggagttaa atagaatatt agaggcatta aaggtcagag 600
 ttctgagacc tgctctggag tgggcagtgt caaacgggga aatgcttata gctcaaaaca 660

gttccttgga atttaagcta cacagactgt attttattag cttgttaatg ggtggaacca	720
caaatcagcg agaggcatta caatatgcta aaaattttca gccatttgcc ctaaatacatc	780
aaaaagacat tcagggtttg atgggaagcc ttgtgtacct gagacaaggg attgagaact	840
caccatatgt tcacctactt gatgcaaacc agtgggctga tatctgtgac atctttacac	900
gggatgcttg tgccctcctg gggctctccg tggagtcccc tctcagtgtc agtttctcag	960
cagggtgtgt ggcgctgcca gctttaatta acatcaaagc cgtgattgaa cagaggcagt	1020
gtactggagt ttggaaccag aaagatgaat tacctattga agtggacctt ggtaaaaagt	1080
gctggtatca ctctatattt gcttgcccca ttcttcgtca gcaaacaaca gataacaatc	1140
cacctatgaa attgggtctgt ggtcatatta tatcaagaga tgccctgaat aaaatgttta	1200
atggtagcaa attaaaatgt ccctactgtc caatggaaca aagtccagga gatgccaac	1260
agatattttt ctgaagagat aaottttagtt tgcaatttgt aagtgaact gaatcgtggg	1320
tgcatttcag aagagaacgt tccatataat gcagctaacc aaggactcct gtgtttctat	1380
aagctaatgc tccagaaact ttgccaacct gttagtgtac acacactgag gggagtgtc	1440
ccggtgaata ttatcatagg gctttattat attcttggtc ttcatttctg atcaagtaaa	1500
tacaccagca gttgtcattc aatgcagggt tttgtactta attatatggt gattttttta	1560
ctttttaaga gcagaaacgg aaattgacct ccccgccatg tgtttaatat tccctctgct	1620
tttacttttg tcattttctt gataatcgta agccttgaga gtgtttgtga aaaagtttta	1680
tttctgttta tgtatacata attaaatgaa aattcttcag aaaaagtttg ataaattgaa	1740
ttgtggttat gaaactaatt tgcattttta tttgcttaag aaagaaagct gtgatagatt	1800
ccagatatgc tttttgatgt tttcctctgc tccagctcca agaagtcagc acacctgcat	1860
tttagctctg catgcagccc cagcaggctg cgtgtttaag aatttcattg tttaactggc	1920
tggtgtgaga agtcttccgt tagcatagag tggaaggagt actattgttt gggtgggttt	1980
ttgtttgttt gttttttgtt tttgctttta ttgccaagag gtgcttgttt taaaagtatg	2040
tttaataaaa tgaaattcta aagttagaag tgttcttaag ttgatattta ctctcttggt	2100
cttggtagcc ctattatatc attgcagaca cagtgttgtg catgtgtgta ttgtatatgt	2160
gcaccagcat caaacattgt gtatctggaa ggaaagcagc atgttccagt tctaaaatgc	2220
agttaccaga cccatcactt taaatcctta aagttagaag ctagcaaatt ttctgtagtg	2280
cagaacgttt attgctgttt ttgtgtttga atagtataat gtttgatgcc tctcttctgc	2340
aaagcgtatc atctcattga ctgtgaatct gtatattatt ctgatggata cagataatga	2400
tcttttctct tgtgaggtat cttcatttat gcactgtcca aaaatagcca tgtgtaagag	2460
tctttctgta tgacgaacta catggaaaag acttctgtgg acataattct gaccgaaacc	2520
catgaagtta cttcagtata agaagaacgt tacacggaaa tcaccaaata ttttgcaact	2580

ttattttcttc tgacatggag tgaacatcaa taggaataact ttcaaagaaa atgaaaacac 2640
 agaagcaaag agaaatgtgg cacttcacat tttaaactac agatggactt ggtttgaggg 2700
 agggggaatc acagatttgg tgctaagtta attagaaact ggcagcgttt tacagtagta 2760
 caccagcctg gatgtttttt ctaaaatgtt tacctgggag agctgggggtt tgtttgtgag 2820
 gagaaagagt actgtggaaa acctctgctt gagtaccatg tggccaggcc tatgtggatg 2880
 gctactccgt gctgtgcggc ttcaccagcg gttgggattg gccagcttg gagtgcctgt 2940
 gtgggtccaac ctcatctgg ccccatagtg acttttgccc catgattctg cttcactgtt 3000
 ggaatcctct ttgaagttcc ccctctcttt gctaaagcag tgaaggaaga gaacagagac 3060
 aaactctttg gactgtgaaa gagaaggtag agaattccag gcaacagtct gaccaaggg 3120
 gtaaaccagt ttattatata tatatttttc ctttgaatta aaacctatga gtgactacaa 3180
 aaaggattag actgagccga ataaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 3239

<210> 18
 <211> 391
 <212> PRT
 <213> Homo sapiens

<400> 18

Met Asp Gln Cys Val Thr Val Glu Arg Glu Leu Glu Lys Val Leu His
1 5 10 15

Lys Phe Ser Gly Tyr Gly Gln Leu Cys Glu Arg Gly Leu Glu Glu Leu
20 25 30

Ile Asp Tyr Thr Gly Gly Leu Lys His Glu Ile Leu Gln Ser His Gly
35 40 45

Gln Asp Ala Glu Leu Ser Gly Thr Leu Ser Leu Val Leu Thr Gln Cys
50 55 60

Cys Lys Arg Ile Lys Asp Thr Val Gln Lys Leu Ala Ser Asp His Lys
65 70 75 80

Asp Ile His Ser Ser Val Ser Arg Val Gly Lys Ala Ile Asp Lys Asn
85 90 95

Phe Asp Ser Asp Ile Ser Ser Val Gly Ile Asp Gly Cys Trp Gln Ala
100 105 110

Asp Ser Gln Arg Leu Leu Asn Glu Val Met Val Glu His Phe Phe Arg
115 120 125

Gln Gly Met Leu Asp Val Ala Glu Glu Leu Cys Gln Glu Ser Gly Leu

130		135		140
Ser Val Asp Pro Ser Gln Lys Glu Pro Phe Val Glu Leu Asn Arg Ile				
145		150		155 160
Leu Glu Ala Leu Lys Val Arg Val Leu Arg Pro Ala Leu Glu Trp Ala				
	165		170	175
Val Ser Asn Arg Glu Met Leu Ile Ala Gln Asn Ser Ser Leu Glu Phe				
	180		185	190
Lys Leu His Arg Leu Tyr Phe Ile Ser Leu Leu Met Gly Gly Thr Thr				
	195		200	205
Asn Gln Arg Glu Ala Leu Gln Tyr Ala Lys Asn Phe Gln Pro Phe Ala				
	210		215	220
Leu Asn His Gln Lys Asp Ile Gln Val Leu Met Gly Ser Leu Val Tyr				
225		230		235 240
Leu Arg Gln Gly Ile Glu Asn Ser Pro Tyr Val His Leu Leu Asp Ala				
	245		250	255
Asn Gln Trp Ala Asp Ile Cys Asp Ile Phe Thr Arg Asp Ala Cys Ala				
	260		265	270
Leu Leu Gly Leu Ser Val Glu Ser Pro Leu Ser Val Ser Phe Ser Ala				
	275		280	285
Gly Cys Val Ala Leu Pro Ala Leu Ile Asn Ile Lys Ala Val Ile Glu				
	290		295	300
Gln Arg Gln Cys Thr Gly Val Trp Asn Gln Lys Asp Glu Leu Pro Ile				
305		310		315 320
Glu Val Asp Leu Gly Lys Lys Cys Trp Tyr His Ser Ile Phe Ala Cys				
	325		330	335
Pro Ile Leu Arg Gln Gln Thr Thr Asp Asn Asn Pro Pro Met Lys Leu				
	340		345	350
Val Cys Gly His Ile Ile Ser Arg Asp Ala Leu Asn Lys Met Phe Asn				
	355		360	365
Gly Ser Lys Leu Lys Cys Pro Tyr Cys Pro Met Glu Gln Ser Pro Gly				
	370		375	380

Asp Ala Lys Gln Ile Phe Phe
385 390

<210> 19
<211> 5565
<212> DNA
<213> Homo sapiens

<400> 19
gcccctcgct gttctgcctt ttgtcttcca cgctgcccc ttcaactcct cttgccctgc 60
aaaccctggg tgcgcacctc cacatccctt ccacttctac ctgctgcct tggcctgcat 120
tttattttca accaagggtg aaatggtagc cccacgcaaa atccgatggg gtgtggagga 180
actgccagct cattgcagtc tgcacgcctt cttaagtgcc agatgcgagc aaggccgacc 240
aggaactgcc gatgggcttc acctgtcaga ctggactgcc acctgccaca cattcattga 300
actttcaata agcactgccg ctgtgcttgg cgttttgctg ggctccagaa ataaaaaatt 360
cagagaggcc aaagtctctc agggatgcag acaggtgata gaaccatcat ttattgagca 420
cttgctaagt gctaaatact cttacctgcg ttatcttgct taatcctcac aaccacactc 480
aacgacggcc tattgaatgc tctctaggtg tctggcagtg ttccaaacag tttatgtgta 540
ttactttggt tgatcttccc aaccaccaat gagcggatat tcaactcctt tgtttacact 600
gagaaaaatc caaatggaga aagtgaagta cagcagctag ccaaaaaaat ccgagagaag 660
ttcaaccgtt acttggaatg ggtcaatcgg aacaagcaag ttgtagaagc atcctatacg 720
gctcacctaa cctctcccct aactgcaatt caagactgct gtactatccc accttccatg 780
atggaattcg atgggaactt taataccaat gtgtctagaa caattagttg tgatcgactt 840
tctactactg ttaatagccg ggcttccaat ccaggacgag acttaaattc agttcttgca 900
gacaacctga aatccaaccc tggaattaag tggcaatatt tcagttcaga agaaggaatt 960
ttcactgttt tcccagcaca caagttccgg tgtaagggca gctacgaaca ccgcagtaga 1020
tttctgtgtt aactgtggca gataccgtcc ggacttgctc actagaccag tgctataaga 1080
ccttctgtgc tccagccacc agtgagacaa aaaggaaaat gtccaccttt gttagcagcg 1140
tgaagtcttc agacagtcct acccagcacg cagtgggatt ccaaaaggca tttcagctga. 1200
ttcgaagtac aaacaataac acaaagttcc aagcaaatag agacatggct atcatttacc 1260
tgtcagctgg cattacatca aaggactctt cggaagaaga taaaaaagcg actctccaag 1320
tcatcaatga agaaaatagc tttctaaaca actctgtaat gattctcacc tatgccctca 1380
tgaacgatgg ggtgactggg ttgaaagagc tggcttttct gagggatcta gctgaacaga 1440
attcagggaa gtacgggtgtg ccagaccgga cggccttgcc tgtgattaag ggcagcatga 1500
tggtgctgaa tcagttgagc aacctggaga ccacagtggg caggttctac acaaaccttc 1560
ccaaccgat gattgatgaa gccgtcttca gcctgccctt ctctgatgag atgggagatg 1620

gtttgataat	gactgtgagt	aaaccctgtt	attttgga	cctacttctg	ggaattgtag	1680
gtgtggacgt	gaatctggct	tacattcttg	aagacgtgac	gtattaccaa	gactctttgg	1740
cttcctatac	ttttctcata	gacgacaaag	gatatacact	tatgcaccca	tctcttacca	1800
ggccatattt	attgtcagag	ccccacttc	atactgacat	catacattat	gaaaatattc	1860
caaaatttga	attagttcgg	caaaatatcc	taagcctccc	tctgggcagc	cagattatcg	1920
cagtcctgt	gaactcatcc	ctgtcttggc	acataaaca	gctgagagaa	actggaaagg	1980
aagcctacaa	tgtagctat	gcctggaaga	tggtacaaga	cacttccttt	attctgtgta	2040
ttgtggtgat	acaaccagaa	atacctgtga	aacaactgaa	gaacctcaac	actgttccca	2100
gcagcaagct	gctgtaccac	cggctggatc	tccttggcca	gccagtgct	tgccctccact	2160
tcaaacagct	ggcaacccta	gaaagtccca	ccatcatgct	gtctgctggc	agcttttccct	2220
ccccctatga	gcacctcagc	cagccagaga	caaagcgcat	ggtagagcac	tacaccgcct	2280
atctcagcga	caacacccgc	ctcattgcta	acccgggcct	caaattctct	gtcagaaatg	2340
aagtaatggc	taccagccac	gtcacagatg	aatggatgac	acaaatggaa	atgagtagcc	2400
tgaacactta	cattgtccgc	cgttacatag	caacacccaa	tggcgctcctc	agaatttatc	2460
ctggttccct	catggacaaa	gcatttgatc	ccactaggag	acaatgggat	ctccatgcag	2520
tagctaatac	agggttgatt	tctttgactg	gtccttactt	agatgttggg	ggagccgggt	2580
atgttgtgac	aatcagtcac	acaattcatt	catccagtac	acagctgtct	tctgggcaca	2640
ctgtggctgt	gatgggcatt	gacttcacac	tcagatactt	ctacaaagtt	ctgatggacc	2700
tattacctgt	ctgtaaccaa	gatgggtggc	acaaaataag	gtgcttcata	atggaggaca	2760
ggggttatct	ggtggcgcac	ccgactctca	tcgaccccaa	aggacatgca	cctgtggagc	2820
agcagcacat	caccacacaag	gagcccctgg	tagcaaata	tatcctcaac	caccccaact	2880
ttgtaaagaa	aaacctgtgc	aacagcttca	gtgacagaac	ggtccagagg	ttttataaat	2940
tcaacaccag	ccttgcgggg	gatttgacga	accttgtgca	tggcagccac	tgttccaaat	3000
acagattagc	aaggatccca	ggaaccaacg	cgtttgttgg	cattgtcaac	gaaacctgcg	3060
actctcttgc	cttctgtgcc	tgacagatgg	tggaccgact	ctgtctcaac	tgtcaccgaa	3120
tggaacaaaa	tgaatgtgaa	tgctcttggt	agtgcctctc	agaggtcaat	gagtgcactg	3180
gcaacctcac	caatgcagag	aaccgaaacc	ccagctgcga	ggtccacctg	gagccggtga	3240
catacacagc	tattgaccct	ggcctgcaag	atgctcttca	ccagtgtgtc	aacagcaggt	3300
gcagtcagag	gctggaaaagt	ggggactgtt	ttgggggtgct	ggattgtgaa	tggtgcatgg	3360
tggacagtga	tggaaaagact	cacctggaca	aacctactg	tgccccccag	aaagaatgct	3420
tcgggggggat	tgtgggagcc	aaaagtcctc	acgttgatga	catgggagca	ataggtgatg	3480
aggtgatcac	attaaacatg	attaaaagcg	cccctgtggg	tcctgtggct	ggaggggatca	3540

tgggatgcat catggtcttg gtcctggcgg tgtatgccta ccgccaccag attcatcgcc	3600
ggagccatca gcatatgtct cctcttgctg cccaagaaat gtcagtgcgt atgtccaacc	3660
tggagaatga cagagatgaa agggacgacg acagccacga agacagaggc atcatcagca	3720
acactcgggt tatagctgcg gtcatcgaac gacatgcaca cagtccagaa agaaggcgcc	3780
gctactgggg tcgatcagga acagaaagtg atcatgggta cagcaccatg agcccacagg	3840
aggacagtga aaatcctcca tgcaacaatg accccttgct agccgggggtc gatgtgggaa	3900
accatgatga ggacttagac ctggataccc cccctcagac tgctgcccta ctaagtcaca	3960
agttccacca ctaccggtca caccacccta cacttcatca tagccaccac ttacaggcgg	4020
ccgtcacggt acacactgtc gatgcagaat gctaacaatc tcctcacctc cagccaaga	4080
tgagatctgg gagccacaga atgttctgga aagaaaaaga accggcttaa aaccacagc	4140
aagagacctc ccttggtgtt gtgctttgtg cagagttgtt tgagtcattt cctgcctgtc	4200
gacatgggta aaaacgagag aaacaacaac acagtcacat ttgtgaagat gtgaggctgg	4260
ttctgaaatg gaggggaaat aagcctgatg aacagacctg ccataacact aatggaaggt	4320
aacagaaggc gaacctccaa acacagagac ggaacctgca agtgaagctg agccagagga	4380
atgttccaaa gagccagaag cattcagctc tccttaactg gaagagagaa aaatctgctc	4440
accagagac tggaatgtgg cacatgcaga tacaatgtg tgcattgaag atttcgcttt	4500
gtttcttagc ggtacctgga taccacagtt gctgtatgga actcatgtta tgcctaaac	4560
gatgcactc agaatttcta agtaaaggat tatttttcta ctatttattg aactttcaaa	4620
cattctcaaa ctttggggaa aaggaaagga aacacaggag aagttttcag cagttgcccc	4680
gagctgtttt gtgtgtaatg aagtggttct ttgattaagg agctctattt cttatttaac	4740
tgatatccca ctgccccact ccacaaaata ggaaaatgaa gaaatctttc tctctgactt	4800
gtttacatca tttcacggaa acacatcttt gtttgtaatg cagtattctt tctctgtgtt	4860
tgacagagat ggggaggggc agaggaattt aagaggtttt aaaagaaatg ttatgtttct	4920
tatgacttgt ttccactcct cgtacaatgc tattcttagg tttctacgaa acctaatgtt	4980
agaaccgcat cctttcagct aaggagggt tggatttatt ttccttgttt tagagactac	5040
aaatttttaa atatccatt ttgactgaga atattgacat ataagggaag aagttttcta	5100
aattgtgaaa gtctggttct taattaaaga attttttttt taatatcacg gttaaaagct	5160
gctgccagtt agccaagaca ttatccacca aattgctttg tgatttatac agggattaat	5220
caaatctggc tactataaca tggggcattg taactttaaa gtagtgtttt aattacagtg	5280
atgtatttta gactcacatt ttgtgattca aatatgttat aaaggcatte ttgcaccatg	5340
gtaaagaatg tgtgtggtaa atctccgttt atatgtagtt ggaaaaaatt cactgaataa	5400
tgttttaatg atagggtatt atgatacaat gtaaaaaaca attggttctt cagcagtaca	5460

gaaagtaaac tatatatgtg ctatcaggaa accccttcac actgtgtata aaattgcaat 5520
 ctagtgaaat aaactgtatg caatggaaaa aaaaaaaaaa aaaca 5565

<210> 20
 <211> 978
 <212> PRT
 <213> Homo sapiens
 <400> 20

Met Ser Thr Phe Val Ser Ser Val Lys Ser Ser Asp Ser Pro Thr Gln
 1 5 10 15

His Ala Val Gly Phe Gln Lys Ala Phe Gln Leu Ile Arg Ser Thr Asn
 20 25 30

Asn Asn Thr Lys Phe Gln Ala Asn Thr Asp Met Val Ile Ile Tyr Leu
 35 40 45

Ser Ala Gly Ile Thr Ser Lys Asp Ser Ser Glu Glu Asp Lys Lys Ala
 50 55 60

Thr Leu Gln Val Ile Asn Glu Glu Asn Ser Phe Leu Asn Asn Ser Val
 65 70 75 80

Met Ile Leu Thr Tyr Ala Leu Met Asn Asp Gly Val Thr Gly Leu Lys
 85 90 95

Glu Leu Ala Phe Leu Arg Asp Leu Ala Glu Gln Asn Ser Gly Lys Tyr
 100 105 110

Gly Val Pro Asp Arg Thr Ala Leu Pro Val Ile Lys Gly Ser Met Met
 115 120 125

Val Leu Asn Gln Leu Ser Asn Leu Glu Thr Thr Val Gly Arg Phe Tyr
 130 135 140

Thr Asn Leu Pro Asn Arg Met Ile Asp Glu Ala Val Phe Ser Leu Pro
 145 150 155 160

Phe Ser Asp Glu Met Gly Asp Gly Leu Ile Met Thr Val Ser Lys Pro
 165 170 175

Cys Tyr Phe Gly Asn Leu Leu Leu Gly Ile Val Gly Val Asp Val Asn
 180 185 190

Leu Ala Tyr Ile Leu Glu Asp Val Thr Tyr Tyr Gln Asp Ser Leu Ala
 195 200 205

Ser Tyr Thr Phe Leu Ile Asp Asp Lys Gly Tyr Thr Leu Met His Pro
 210 215 220

Ser Leu Thr Arg Pro Tyr Leu Leu Ser Glu Pro Pro Leu His Thr Asp
 225 230 235 240

Ile Ile His Tyr Glu Asn Ile Pro Lys Phe Glu Leu Val Arg Gln Asn
 245 250 255

Ile Leu Ser Leu Pro Leu Gly Ser Gln Ile Ile Ala Val Pro Val Asn
 260 265 270

Ser Ser Leu Ser Trp His Ile Asn Lys Leu Arg Glu Thr Gly Lys Glu
 275 280 285

Ala Tyr Asn Val Ser Tyr Ala Trp Lys Met Val Gln Asp Thr Ser Phe
 290 295 300

Ile Leu Cys Ile Val Val Ile Gln Pro Glu Ile Pro Val Lys Gln Leu
 305 310 315 320

Lys Asn Leu Asn Thr Val Pro Ser Ser Lys Leu Leu Tyr His Arg Leu
 325 330 335

Asp Leu Leu Gly Gln Pro Ser Ala Cys Leu His Phe Lys Gln Leu Ala
 340 345 350

Thr Leu Glu Ser Pro Thr Ile Met Leu Ser Ala Gly Ser Phe Ser Ser
 355 360 365

Pro Tyr Glu His Leu Ser Gln Pro Glu Thr Lys Arg Met Val Glu His
 370 375 380

Tyr Thr Ala Tyr Leu Ser Asp Asn Thr Arg Leu Ile Ala Asn Pro Gly
 385 390 395 400

Leu Lys Phe Ser Val Arg Asn Glu Val Met Ala Thr Ser His Val Thr
 405 410 415

Asp Glu Trp Met Thr Gln Met Glu Met Ser Ser Leu Asn Thr Tyr Ile
 420 425 430

Val Arg Arg Tyr Ile Ala Thr Pro Asn Gly Val Leu Arg Ile Tyr Pro
 435 440 445

Gly Ser Leu Met Asp Lys Ala Phe Asp Pro Thr Arg Arg Gln Trp Tyr
 450 455 460

Leu His Ala Val Ala Asn Pro Gly Leu Ile Ser Leu Thr Gly Pro Tyr
 465 470 475 480

Leu Asp Val Gly Gly Ala Gly Tyr Val Val Thr Ile Ser His Thr Ile
 485 490 495

His Ser Ser Ser Thr Gln Leu Ser Ser Gly His Thr Val Ala Val Met
 500 505 510

Gly Ile Asp Phe Thr Leu Arg Tyr Phe Tyr Lys Val Leu Met Asp Leu
 515 520 525

Leu Pro Val Cys Asn Gln Asp Gly Gly Asn Lys Ile Arg Cys Phe Ile
 530 535 540

Met Glu Asp Arg Gly Tyr Leu Val Ala His Pro Thr Leu Ile Asp Pro
 545 550 555 560

Lys Gly His Ala Pro Val Glu Gln Gln His Ile Thr His Lys Glu Pro
 565 570 575

Leu Val Ala Asn Asp Ile Leu Asn His Pro Asn Phe Val Lys Lys Asn
 580 585 590

Leu Cys Asn Ser Phe Ser Asp Arg Thr Val Gln Arg Phe Tyr Lys Phe
 595 600 605

Asn Thr Ser Leu Ala Gly Asp Leu Thr Asn Leu Val His Gly Ser His
 610 615 620

Cys Ser Lys Tyr Arg Leu Ala Arg Ile Pro Gly Thr Asn Ala Phe Val
 625 630 635 640

Gly Ile Val Asn Glu Thr Cys Asp Ser Leu Ala Phe Cys Ala Cys Ser
 645 650 655

Met Val Asp Arg Leu Cys Leu Asn Cys His Arg Met Glu Gln Asn Glu
 660 665 670

Cys Glu Cys Pro Cys Glu Cys Pro Leu Glu Val Asn Glu Cys Thr Gly
 675 680 685

Asn Leu Thr Asn Ala Glu Asn Arg Asn Pro Ser Cys Glu Val His Leu
 690 695 700

Glu Pro Val Thr Tyr Thr Ala Ile Asp Pro Gly Leu Gln Asp Ala Leu
 705 710 715 720

His Gln Cys Val Asn Ser Arg Cys Ser Gln Arg Leu Glu Ser Gly Asp
 725 730 735

Cys Phe Gly Val Leu Asp Cys Glu Trp Cys Met Val Asp Ser Asp Gly
 740 745 750

Lys Thr His Leu Asp Lys Pro Tyr Cys Ala Pro Gln Lys Glu Cys Phe
 755 760 765

Gly Gly Ile Val Gly Ala Lys Ser Pro Tyr Val Asp Asp Met Gly Ala
 770 775 780

Ile Gly Asp Glu Val Ile Thr Leu Asn Met Ile Lys Ser Ala Pro Val
 785 790 795 800

Gly Pro Val Ala Gly Gly Ile Met Gly Cys Ile Met Val Leu Val Leu
 805 810 815

Ala Val Tyr Ala Tyr Arg His Gln Ile His Arg Arg Ser His Gln His
 820 825 830

Met Ser Pro Leu Ala Ala Gln Glu Met Ser Val Arg Met Ser Asn Leu
 835 840 845

Glu Asn Asp Arg Asp Glu Arg Asp Asp Asp Ser His Glu Asp Arg Gly
 850 855 860

Ile Ile Ser Asn Thr Arg Phe Ile Ala Ala Val Ile Glu Arg His Ala
 865 870 875 880

His Ser Pro Glu Arg Arg Arg Arg Tyr Trp Gly Arg Ser Gly Thr Glu
 885 890 895

Ser Asp His Gly Tyr Ser Thr Met Ser Pro Gln Glu Asp Ser Glu Asn
 900 905 910

Pro Pro Cys Asn Asn Asp Pro Leu Ser Ala Gly Val Asp Val Gly Asn
 915 920 925

His Asp Glu Asp Leu Asp Leu Asp Thr Pro Pro Gln Thr Ala Ala Leu
 930 935 940

Leu Ser His Lys Phe His His Tyr Arg Ser His His Pro Thr Leu His
 945 950 955 960

His Ser His His Leu Gln Ala Ala Val Thr Val His Thr Val Asp Ala
 965 970 975

Glu Cys

<210> 21
 <211> 3352
 <212> DNA
 <213> Homo sapiens

<400> 21
 agcgggcagg gaccagatcc gcgagcccggt cagcctgcgc catgggctgc gacggccgcg 60
 tgtcggggct gctccgccgc aacctgcagc ccacgctcac ctactggagc gtcttcttca 120
 gcttcggcct gtgcatcgcc ttctggggc ccacgctgct ggacctgcgc tgtcagacgc 180
 acagctcgct gccccagatc tcctgggtct tcttctcgca gcagctctgc ctctgctgg 240
 gcagcgccct cgggggcgtc ttcaaaagga ccctggccca gtcactatgg gccctgttca 300
 cctcctctct ggccatctcc ctgggtgttg ccgtcatccc cttctgccgc gacgtgaagg 360
 tgctggcctc agtcatggcg ctggcgggct tggccatggg ctgcatcgac accgtggcca 420
 acatgcagct ggtaaggatg taccagaagg actcggccgt ctctctccag gtgtccatt 480
 tcttcgtggg ctttggtgct ctgctgagcc cccttattgc tgaccctttc ctgtctgagg 540
 ccaactgctt gcctgccaat agcacggcca acaccacctc ccgaggccac ctgttccatg 600
 tctccagggt gctgggccag caccacgtag atgccaagcc ttggtccaac cagacgttcc 660
 cagggtgac tccaaaggac ggggcaggga cccgagtgtc ctatgccttc tggatcatgg 720
 ccatcatcaa tcttccagtg cccatggctg tgctgatgct gctgtccaag gagcggctgc 780
 tgacctgctg tccccagagg aggccctgc ttctgtctgc tgatgagctt gccttgaga 840
 cacagcctcc tgagaaggaa gatgcctcct cactgcccc aaagtctcag tcacacctag 900
 ggcatgagga cctgttcagc tgctgccaaa ggaagaacct cagaggagcc cttattcct 960
 tctttgccat ccacatcacg gccgcctgg tcctgttcat gacggatggg ttgacgggtg 1020
 cctattccgc cttcgtgtac agctatgctg tggagaagcc cctgtctgtg ggacacaagg 1080
 tggctggcta cctccccagc ctcttctggg gcttcatcac actgggccgg ctctctcca 1140
 ttcccatatc ctcaagaatg aagccggcca ccatggtttt catcaacgtg gttggcgtgg 1200
 tggtgacgtt cctgggtgctg cttattttct cctacaacgt cgtcttcctg ttcgtgggga 1260
 cggcaagcct gggcctgttt ctcagcagca ctttccag catgctggcc tacacggagg 1320
 actcgtgca gtacaaaggc tgtgcaacca cagtgtggt gacaggggca ggagttggcg 1380
 agatggtgct gcagatgctg gttggttoga tattccaggc tcagggcagc tatagtttcc 1440
 tggctctgtg cgtgatcttt ggttgtctgg cttttacctt ctatatcttg ctctgtttt 1500
 tccacaggat gcaccctgga ctcccatcag ttctaccca agacagatca attggaatgg 1560
 aaaactctga gtgctaccag aggtaaaact gggtaagaa ggcaagagaa gactttcagc 1620

```

ctcttgatca ccagcacgac catactgttt cagaaagctg ggtggtggtg gaggcgctat 1680
ctcaatggct attcaagtct tctccactaa aacttggttg ggtagaggaa attaaattga 1740
gtcctggtac ctggtcaaaa tcattagaag tttacctggc ttctcaagtt atcttcttcc 1800
ctggttcaga ctggttggtaa gagctgtcca gatacccaga tgggaaggaa ggagacagcc 1860
gcgcgcttca ctccatttgt cacctcatgc atggaccata ctctgggttt gagatcattc 1920
ttcattgaag tttgtaaaaa taggttgaaa ttgtaaagct ccatgatcat tgctatgtag 1980
atatatttca atttaagcaa aacaagctgc aagttattcc ctggcatgct caaaggattt 2040
tcgtgctttt cacttaatatg tccaaagtct cttaaattcc tgctgcagac atcaatagct 2100
tatctatatt ctcaaacacc aaaaggaaaa gttgaatctt gctctctttg gtataactaat 2160
gtagtggtat gctaagctgg ctcataccaa cttagaaaag ctgattgtaa aattttcatt 2220
ttgacagctg gttattaaat gcagccatta ttaaaaatca aatcatacaa acttataatt 2280
aatcaatta catttaaaac aaaggtaata aatattcaaa gcatatcact tcctaatttg 2340
atcttgatgc tcttgaggta atttacgtcc atggtacctg tgtggtggaa ttactatata 2400
tgatggtgtg ctactgtgca ccttgtctca actccactct tcgtgatagc atggtggtag 2460
cttgaaatca gcctggtggg agtattacca tggacactgg caaaagctac agatcccgga 2520
gagccagtgg ttaaacattt accagcatat cactgctagt aatcaaggct aactggtcca 2580
gaaatcgccc aggagaatga aatggatgtt ccattttttt ctactgacat tgactagcat 2640
ataaaaggta tagaaacagc actaagactt tctgaaaata cctaataaaa attttacatc 2700
tttttttggt tttgtttttg tttttgagac agagtctcac tctgttgcca ggctggaatg 2760
cagtggcgtc atctcggtc actgcaacct ccgcctccca gtttcaagcg attctcctgc 2820
ctcagcctcc caagtagctg tgactacagg cgcatgccat cacgcacagc taattttttg 2880
tatttttagt agagacggga tttcacctg ttagccagga tggctcttgat ctctgacct 2940
catgatctgc ccgtctcggc ctcccaaagt gctgggatta caggcgtgag cactgcacc 3000
cagccaaaaa ttttacatct tttatagagg gaaaaaaact ctttatacca tggcaaggcc 3060
ttttctttca caaaaagctg ggcctactga acaattcaag ctgtgcagta gtagactgaa 3120
agcaggattt gttgaggagt tacagctcct gtccagagca aatcctgtag tgatacaagg 3180
agaatgtaaa cttgccagct tagacaggga tcagtcctga gactgctggc agtagcaaat 3240
ggctattaga gtaactgtat aatggttttg cctgcacttt ctctatgtat atacaaatgt 3300
acatgtataa atataaaaat taagtgtatc tgaaagaaaa aaaaaaaaaa aa 3352

```

```

<210> 22
<211> 403
<212> PRT
<213> Homo sapiens

```

<400> 22

Met Ala Leu Ala Gly Leu Ala Met Gly Cys Ile Asp Thr Val Ala Asn
 1 5 10 15

Met Gln Leu Val Arg Met Tyr Gln Lys Asp Ser Ala Val Phe Leu Gln
 20 25 30

Val Leu His Phe Phe Val Gly Phe Gly Ala Leu Leu Ser Pro Leu Ile
 35 40 45

Ala Asp Pro Phe Leu Ser Glu Ala Asn Cys Leu Pro Ala Asn Ser Thr
 50 55 60

Ala Asn Thr Thr Ser Arg Gly His Leu Phe His Val Ser Arg Val Leu
 65 70 75 80

Gly Gln His His Val Asp Ala Lys Pro Trp Ser Asn Gln Thr Phe Pro
 85 90 95

Gly Leu Thr Pro Lys Asp Gly Ala Gly Thr Arg Val Ser Tyr Ala Phe
 100 105 110

Trp Ile Met Ala Ile Ile Asn Leu Pro Val Pro Met Ala Val Leu Met
 115 120 125

Leu Leu Ser Lys Glu Arg Leu Leu Thr Cys Cys Pro Gln Arg Arg Pro
 130 135 140

Leu Leu Leu Ser Ala Asp Glu Leu Ala Leu Glu Thr Gln Pro Pro Glu
 145 150 155 160

Lys Glu Asp Ala Ser Ser Leu Pro Pro Lys Phe Gln Ser His Leu Gly
 165 170 175

His Glu Asp Leu Phe Ser Cys Cys Gln Arg Lys Asn Leu Arg Gly Ala
 180 185 190

Pro Tyr Ser Phe Phe Ala Ile His Ile Thr Ala Ala Leu Val Leu Phe
 195 200 205

Met Thr Asp Gly Leu Thr Gly Ala Tyr Ser Ala Phe Val Tyr Ser Tyr
 210 215 220

Ala Val Glu Lys Pro Leu Ser Val Gly His Lys Val Ala Gly Tyr Leu
 225 230 235 240

Pro Ser Leu Phe Trp Gly Phe Ile Thr Leu Gly Arg Leu Leu Ser Ile

```
<210> 23
<211> 1893
<212> DNA
<213> Homo sapiens
```

<400>	23	gtgctgttcc	tgttctgtgt	ggccctatac	caggtgcact	atgaatgggtg	gaggaaaaatt	60
		ctaaaaatatt	tttggatgtc	agtggttatt	tacactatgc	tggtgcttat	ctttatatac	120
		acatatcagt	ttgaggactt	cccaggcctg	tggcaaaaata	tgactggatt	gaaaaaagaa	180
		aagcttgagg	atcttggctt	aaagcagttt	actgtggctg	aactattcac	tcgcatattc	240
		atcccaacct	cctttctgct	ggtgtgcatt	ttacacctgc	actacttcca	tgaccgggttc	300
		cttgaactca	cagacctcaa	gtccattccc	agcaaagaag	acaacaccat	ctacagccat	360
		gccaaagtca	atggtcgcgt	atatctaata	ataaatagca	tcaagaaaaa	attaccaatc	420

```

caccaaaatg aactggccca cccggaagga agcctcccgg acctcaccat gatgcatctg 480
actgccagcc tggagaagcc ggaggtgagg aagttggctg agcctgggga ggagaagctt 540
gagggctact ctgaaaaagc ccagaagggg gatcttggga aagacagcga ggagtcagag 600
gaggacggag aggaagagga ggaatccgag gaggaggaag aaacatcaga ctttaaggaac 660
aatggcacc tgggtgattga ccgcctcact gtgctcttct taaaattcct ggagtatttt 720
cacaagctgc aggtgttcat gtggtggatt ttggagttgc acatcatcaa aatcgtttcc 780
tcttacatta tctgggtttc tgtgaaagag gtgtctctgt tcaactatgt atttttgatt 840
tcttgggctt ttgctctgcc gtacgccaaag ctgcgcctgc tggcttcaag tgtctgcacg 900
gtctggacgt gtgtgatcat cgtctgcaaa atgtcgtacc agcttcaaac cattaagcct 960
gagaacttct ctgttaactg ttccttgcca aatgaaaatc aaacaaacat cccctttaat 1020
gagttgaaca agtctctgct ctacagcgct cctatcgatc ctacagagtg ggtcggcctg 1080
cggaagtctt cgcctctgct agtctacctg aggaataacc tcctgatgct ggctatcctg 1140
gcctttgaag tcaccattta ccgccatcag gaataactatc gaggtcgaaa taacctgacg 1200
gcccctgtgt ctagaactat ctttcatgac attacaagac tacatctaga tgatggactt 1260
attaattgtg ccaaataattt cattaattac ttcttttaca agtttggctt ggagacctgt 1320
ttcctaattg cagttaacgt cattggccag cgaatggatt tctatgccat gatccacgcc 1380
tgctggctga tcgctgtctt atatagacgc agaaggaaag ccatcgcaga gatctggccc 1440
aagtactgct gcttcttggc atgcatcatc accttccagt atttcatctg cattggcatc 1500
ccacctgctc cttgccgaga ttacctgtgg agattcaagg gtgccagctt caatgacaac 1560
atcataaagt ggctgtactt cccagatttc attgtgcggc ccaaccctgt gtttctcgtc 1620
tatgacttca tgctgcttct gtgtgcctcc ttacaacggc agatttttga ggatgagaac 1680
aaggctgcag tgcgaaatcat ggcaggtgac aatgtcgaga tctgcatgaa tcttgatgcy 1740
gcctccttca gccaacataa ccctgggtgcc accttgcaa cttttgtatt cacaggcatc 1800
ccatctccat cacagataaa atcttaggag ataaacacat tcaaaaagga atgagataaa 1860
aagaataagg caataaatgt tgattggaac etc 1893

```

```

<210> 24
<211> 583
<212> PRT
<213> Homo sapiens

```

```

<400> 24

```

```

Met Ser Val Val Ile Tyr Thr Met Leu Val Leu Ile Phe Ile Tyr Thr
1           5           10          15

```

```

Tyr Gln Phe Glu Asp Phe Pro Gly Leu Trp Gln Asn Met Thr Gly Leu

```


20	25	30	
Lys Lys Glu Lys Leu Glu Asp	Leu Gly Leu Lys Gln Phe Thr Val Ala		
35	40	45	
Glu Leu Phe Thr Arg Ile Phe Ile Pro Thr Ser Phe Leu Leu Val Cys			
50	55	60	
Ile Leu His Leu His Tyr Phe His Asp Arg Phe Leu Glu Leu Thr Asp			
65	70	75	80
Leu Lys Ser Ile Pro Ser Lys Glu Asp Asn Thr Ile Tyr Ser His Ala			
85	90	95	
Lys Val Asn Gly Arg Val Tyr Leu Ile Ile Asn Ser Ile Lys Lys Lys			
100	105	110	
Leu Pro Ile His Gln Asn Glu Leu Ala His Pro Glu Gly Ser Leu Pro			
115	120	125	
Asp Leu Thr Met Met His Leu Thr Ala Ser Leu Glu Lys Pro Glu Val			
130	135	140	
Arg Lys Leu Ala Glu Pro Gly Glu Glu Lys Leu Glu Gly Tyr Ser Glu			
145	150	155	160
Lys Ala Gln Lys Gly Asp Leu Gly Lys Asp Ser Glu Glu Ser Glu Glu			
165	170	175	
Asp Gly Glu Glu Glu Glu Glu Ser Glu Glu Glu Glu Glu Thr Ser Asp			
180	185	190	
Leu Arg Asn Lys Trp His Leu Val Ile Asp Arg Leu Thr Val Leu Phe			
195	200	205	
Leu Lys Phe Leu Glu Tyr Phe His Lys Leu Gln Val Phe Met Trp Trp			
210	215	220	
Ile Leu Glu Leu His Ile Ile Lys Ile Val Ser Ser Tyr Ile Ile Trp			
225	230	235	240
Val Ser Val Lys Glu Val Ser Leu Phe Asn Tyr Val Phe Leu Ile Ser			
245	250	255	
Trp Ala Phe Ala Leu Pro Tyr Ala Lys Leu Arg Arg Leu Ala Ser Ser			
260	265	270	

Val Cys Thr Val Trp Thr Cys Val Ile Ile Val Cys Lys Met Ser Tyr
 275 280 285

Gln Leu Gln Thr Ile Lys Pro Glu Asn Phe Ser Val Asn Cys Ser Leu
 290 295 300

Pro Asn Glu Asn Gln Thr Asn Ile Pro Phe Asn Glu Leu Asn Lys Ser
 305 310 315 320

Leu Leu Tyr Ser Ala Pro Ile Asp Pro Thr Glu Trp Val Gly Leu Arg
 325 330 335

Lys Ser Ser Pro Leu Leu Val Tyr Leu Arg Asn Asn Leu Leu Met Leu
 340 345 350

Ala Ile Leu Ala Phe Glu Val Thr Ile Tyr Arg His Gln Glu Tyr Tyr
 355 360 365

Arg Gly Arg Asn Asn Leu Thr Ala Pro Val Ser Arg Thr Ile Phe His
 370 375 380

Asp Ile Thr Arg Leu His Leu Asp Asp Gly Leu Ile Asn Cys Ala Lys
 385 390 395 400

Tyr Phe Ile Asn Tyr Phe Phe Tyr Lys Phe Gly Leu Glu Thr Cys Phe
 405 410 415

Leu Met Ser Val Asn Val Ile Gly Gln Arg Met Asp Phe Tyr Ala Met
 420 425 430

Ile His Ala Cys Trp Leu Ile Ala Val Leu Tyr Arg Arg Arg Arg Lys
 435 440 445

Ala Ile Ala Glu Ile Trp Pro Lys Tyr Cys Cys Phe Leu Ala Cys Ile
 450 455 460

Ile Thr Phe Gln Tyr Phe Ile Cys Ile Gly Ile Pro Pro Ala Pro Cys
 465 470 475 480

Arg Asp Tyr Pro Trp Arg Phe Lys Gly Ala Ser Phe Asn Asp Asn Ile
 485 490 495

Ile Lys Trp Leu Tyr Phe Pro Asp Phe Ile Val Arg Pro Asn Pro Val
 500 505 510

Phe Leu Val Tyr Asp Phe Met Leu Leu Leu Cys Ala Ser Leu Gln Arg
 515 520 525

Gln Ile Phe Glu Asp Glu Asn Lys Ala Ala Val Arg Ile Met Ala Gly
 530 535 540

Asp Asn Val Glu Ile Cys Met Asn Leu Asp Ala Ala Ser Phe Ser Gln
 545 550 555 560

His Asn Pro Gly Ala Thr Thr Ala Thr Phe Val Phe Thr Gly Ile Pro
 565 570 575

Ser Pro Ser Gln Ile Lys Ser
 580

<210> 25

<211> 2763

<212> DNA

<213> Homo sapiens

<400> 25

```

agccgggcag ctgcagcggg gccgcgggagc gggcgggcgagg gccagggctg tgcgcttggg      60
gagcgcgggaa tgtgaggctt ggcgggccgc agcacgctcg gacggggccag gggcgggcgac      120
ccctcgcgga cgcggggctg cgcggcgggc cgggggacttg cccttgcacg ctccctgcgc      180
cctccagctc gccggcgggg ccatgaagaa gttctctcgg atgcccaagt cggagggcgagg      240
cagcggcgggc ggagcggcgg gtggcgggggc tggcgggggcc ggggcccgggg ccggctgcgg      300
ctccggcgggc tegtccgtgg gggtcggggg gttcgcggtc ggccgccacc aggtcacctt      360
ggaagagtcg ctggccgaag gtggattctc cacagttttc ctctgctgta ctacgggtgg      420
aatccgatgt gcattgaagc gaatgtatgt caataacatg ccagacctca atgtttgtaa      480
aagggaaatt acaattatga aagagctatc tggtcacaaa aatattgtgg gctatttgga      540
ctgtgctgtt aattcaatta gtgataatgt atgggaagtc cttatcttaa tggaatattg      600
tcgagctgga caggtagtga atcaaagaa taagaagcta cagacgggtt ttacagaacc      660
agaagtgtta cagatattct gtgatacctg tgaagctgtt gcaagggtgc atcagtgtaa      720
gactccaata attcaccggg atctgaaggt agaaaatatt ttgttgaatg atgggtggaa      780
ctatgtactt tgtgactttg gcagtgccac taataaattt cttaatcttc aaaaagatgg      840
agttaatgta gtagaagaag aaattaaaaa gtatacaact ctgtcataca gagcccctga      900
aatgatcaac ctttatggag ggaaacccat caccaccaag gctgatattt gggcactggg      960
atgtctactc tataaacttt gtttcttcac tcttcctttt ggtgagagtc aggttgctat      1020
ctgtgatggc aacttcacca tcccagacaa ttctcggttac tcccgtaca tacattgctt      1080
aataagggtc atgcttgaac cagatccgga acatagacct gatataattt aagtgtcata      1140
ttttgcattt aaatttgcca aaaaggattg tccagtctcc aacatcaata agtggtgtaa      1200
acaattactg agacacggag cactgttaac tgaaattctt ctattccttc agctcttctt      1260

```

```

gaaccgatga ctgctagtga agcagctgct agggaaaagcc aaataaaaagc cagaataaca 1320
gataccattg gaccaacaga aacctcaatt gcaccaagac aaagacccaaa ggccaactct 1380
gctactactg ccactcccag tgtgctgacc attcaaagtt cagcaacacc tgttaaagtc 1440
cttgctcctg gtgaattcgg taaccataga ccaaaagggg cactaagacc tggaaatggc 1500
cctgaaatth tattgggtca gggacctcct cagcagccgc cacagcagca tagagtactc 1560
cagcaactac agcagggaga ttggagatta cagcaactcc atttacagca tcgtcatcct 1620
caccagcagc agcagcagca gcagcagcaa cagcaacagc agcagcagca acagcaacag 1680
cagcagcagc agcagcagca gcagcaccac caccaccacc accaccacct acttcaagat 1740
gcttatatgc agcagtatca acatgcaaca cagcagcaac agatgcttca acaacaatth 1800
ttaatgcatt cggatatatca accacaacct tctgcatcac agtatcctac aatgatgccg 1860
cagtatcagc aggcctttctt tcaacagcag atgctagctc aacatcagcc gtctcaacaa 1920
caggcatcac ctgaatatct tacctccctt caagagttct caccagcctt agtttcttac 1980
acttcatcac ttccagctca ggttggaacc ataatggact cctcctatag tgccaatagg 2040
tcagttgctg ataaagaggc cattgcaaat ttcacaaatc agaagaacat cagcaatcca 2100
cctgatatgt caggggtgga tccttttgga gaggataatt tctctaagtt aacagaagag 2160
gaactattgg acagagaatt tgaccttcta agatcaagta agggacactt gaaggcttat 2220
tttgcttcac agtaaaataa cagctctatt attattcagc aaggccaaag acttttgaga 2280
atgtgtatgg aaaattcttt ttgtgcattt gagggcaaaa ttcaggccat cttcttatac 2340
atatactatc aaattatggt gtgtgcatta gaaatcagtt gcttgatagt agctattaaa 2400
cccaatattg ctgatagtat gctaatatcc taaaacttaa atattgcata tctatgaatg 2460
ttaaattcag aatatctcta aaacatggaa aattgatggt tcaataaaaaa gggagacatt 2520
ttattattht gccttactaa tgattttgca gctctgttht tctgcacact caatagaaaa 2580
tattgtgggt ctgagatgcc ctttgaaaat gcctgaaaga aaacatgggc tactacatta 2640
tggttaatggt ttgtaatgtc cttttaatga gtggtgacat aaaagggctg ctttgttgtc 2700
ctgatatggc aaaaattaaa taaattcttt ttcactthaa aaaaaaaaaa aaaaaaaaaa 2760
aaa 2763

```

<210> 26
 <211> 355
 <212> PRT
 <213> Homo sapiens

<400> 26

Met Lys Lys Phe Ser Arg Met Pro Lys Ser Glu Gly Gly Ser Gly Gly
 1 5 10 15

Gly Ala Ala Gly Gly Gly Ala Gly Gly Ala Gly Ala Gly Ala Gly Cys
 20 25 30

Gly Ser Gly Gly Ser Ser Val Gly Val Arg Val Phe Ala Val Gly Arg
 35 40 45

His Gln Val Thr Leu Glu Glu Ser Leu Ala Glu Gly Gly Phe Ser Thr
 50 55 60

Val Phe Leu Val Arg Thr His Gly Gly Ile Arg Cys Ala Leu Lys Arg
 65 70 75 80

Met Tyr Val Asn Asn Met Pro Asp Leu Asn Val Cys Lys Arg Glu Ile
 85 90 95

Thr Ile Met Lys Glu Leu Ser Gly His Lys Asn Ile Val Gly Tyr Leu
 100 105 110

Asp Cys Ala Val Asn Ser Ile Ser Asp Asn Val Trp Glu Val Leu Ile
 115 120 125

Leu Met Glu Tyr Cys Arg Ala Gly Gln Val Val Asn Gln Met Asn Lys
 130 135 140

Lys Leu Gln Thr Gly Phe Thr Glu Pro Glu Val Leu Gln Ile Phe Cys
 145 150 155 160

Asp Thr Cys Glu Ala Val Ala Arg Leu His Gln Cys Lys Thr Pro Ile
 165 170 175

Ile His Arg Asp Leu Lys Val Glu Asn Ile Leu Leu Asn Asp Gly Gly
 180 185 190

Asn Tyr Val Leu Cys Asp Phe Gly Ser Ala Thr Asn Lys Phe Leu Asn
 195 200 205

Pro Gln Lys Asp Gly Val Asn Val Val Glu Glu Glu Ile Lys Lys Tyr
 210 215 220

Thr Thr Leu Ser Tyr Arg Ala Pro Glu Met Ile Asn Leu Tyr Gly Gly
 225 230 235 240

Lys Pro Ile Thr Thr Lys Ala Asp Ile Trp Ala Leu Gly Cys Leu Leu
 245 250 255

Tyr Lys Leu Cys Phe Phe Thr Leu Pro Phe Gly Glu Ser Gln Val Ala
 260 265 270

Ile Cys Asp Gly Asn Phe Thr Ile Pro Asp Asn Ser Arg Tyr Ser Arg
 275 280 285

Asn Ile His Cys Leu Ile Arg Phe Met Leu Glu Pro Asp Pro Glu His
 290 295 300

Arg Pro Asp Ile Phe Gln Val Ser Tyr Phe Ala Phe Lys Phe Ala Lys
 305 310 315 320

Lys Asp Cys Pro Val Ser Asn Ile Asn Lys Cys Cys Lys Gln Leu Leu
 325 330 335

Arg His Gly Ala Leu Leu Thr Glu Ile Leu Leu Phe Leu Gln Leu Phe
 340 345 350

Leu Asn Arg
 355

<210> 27
 <211> 4894
 <212> DNA
 <213> Homo sapiens

<400> 27
 cgaaaacgga gaaacccccg gtccggcgag aggggctgtg acagtcggag tcccaagctg . 60
 cggttcggct gctgccgaga actgcaaggt gtggaatatt tctggcttct agtccaatgc 120
 caagtgtgtg acctgtggct acatgattcc ctgaaagata agaacaatgt tatgttgggg 180
 atattggtct ctgggccaac ctggtatcag caccaacctg caggggaattg tggctgagcc 240
 ccaggtgtgt gggttcatat ctgacagaag tgtcaaggaa gtggcctgtg ggggaaacca 300
 ctctgtgttc ctgctggaag atggggaagt ttacacatgt ggtttgaaca ccaaggggca 360
 actgggccat gagagggaag gaaacaagcc agaacaaatt ggagctctgg cagatcagca 420
 tatcattcat gtggcatgtg gcgagtccca cagtctggcc ctcaagtacc gaggccagct 480
 gttttcttgg ggtgcaggga gtgatgttca gctaggactc atgactactg aggattctgt 540
 ggcagtgtcc aggttaatac aaaagctgaa ccagcaaaca atattacaag tttcctgtgg. 600
 caactggcat tgcttggctc ttgcggctga tggccagttc ttcacctggg gaaagaacag 660
 ccatgggcag cttggcttag ggaaggagtt cccctcccaa gccagccac agagggtgag 720
 gtccctggag gggatccac tggctcaggt ggctgccgga ggggctcaca gctttgccct 780
 gtctctctca ggagctgttt ttggctgggg gatgaataat gccgggcagc tagggctcag 840
 tgatgaaaaa gatcgagaat ctccatgcca tgtaaaactc ttacgcacgc aaaaagttgt 900
 ctatattagt tgtggagaag aacacacagc agttctcaca aagagtggag gtgtgtttac 960
 ctttggcgct ggctcctgtg ggcaacttgg acacgactcc atgaatgatg aggttaaccc 1020

tagaagagtt	ctagagctga	tgggtagtga	agtaactcaa	attgcttgtg	gcagacaaca	1080
taccctagcc	ttcgtgcctt	cttctggact	catctatgca	tttggttgtg	gagcaagagg	1140
tcaattagga	actgggcaca	cttgtaatgt	taagtgccca	tctcctgtca	agggttactg	1200
ggctgcccac	agtggccagc	tttcagcccg	agctgatcgc	tttaaataac	atatcgtaa	1260
gcagatcttc	tctggaggag	accagacttt	tgtactttgc	tccaaatacg	agaattattc	1320
tctgtctgtt	gacttcagga	ctatgaacca	agcacattat	accagtttaa	taaagtatga	1380
aaccatagca	gtttggagac	aaaaactctc	agaacacaac	aatgcaaata	caatcaatgg	1440
tgttggtcag	atattatctt	ctgcagcctg	ttggaatgga	agttttcttg	aaaaaaaaat	1500
tgatgaacat	tttaaaacga	gtcccaaaat	ccctgggatt	gacctgaact	caactagggg	1560
gttatttgag	aagttaatga	actctcagca	ctccatgatt	ctagaacaga	ttttgaacag	1620
ttttgaaagt	tgtctgattc	cccagttgtc	aagctcacca	ccagatgttg	aagccatgag	1680
aatctattta	atactacctg	agtttcccct	actccaggat	tccaagtatt	atataacatt	1740
gactattccc	ttggctatgg	ccattcttcg	gctggataca	aaccccagca	aagtactaga	1800
taactggtgg	tctcaggtat	gcccgaata	tttcatgaag	ctggtaaacc	tctataaagg	1860
tgcagtcctt	tatctactga	ggggaagaaa	gacattctta	attcccgtac	tgtttaacaa	1920
ttatatcaca	gcagctctca	aactcttgga	gaagttatat	aaggtaaata	ttaaagtga	1980
gcattgtgga	tatgatacat	tttacattcc	tgagatttcc	aatctcgtgg	acattcagga	2040
agactacctc	atgtggttct	tgcattcaagc	agggatgaag	gctagaccat	caataatata	2100
ggatactgta	acactttggt	cctacccttt	catctttgat	gcccagcca	agaccaaata	2160
gttacagaca	gatgctgaac	tacagatgca	ggtaggcagc	aatggagcca	acctgcagaa	2220
tgtcttcatg	cttctcacc	tggagcctct	gctggccaga	agcccttcc	tggctcttca	2280
cgttcgcagg	aacaaccttg	ttggagatgc	cctaagagag	ctgagcattc	attctgatat	2340
tgatttgaaa	aagcctctca	aagtaatctt	tgatggtgaa	gaagcagtg	atgccggtgg	2400
tgttacaaag	gaattttttc	ttttgctgtt	aaaagaactt	ttgaatocca	tctatggaat	2460
gtttacctac	tatcaagatt	caaactctct	gtggttttca	gacacgtgtt	ttgtagagca	2520
caactgggtt	cacttgattg	gtataacctg	tggactagct	atctacaact	ccactgtggt	2580
cgatctccac	ttcccattgg	ctctctacaa	gaagttactc	aatgtaaagc	ctggcttgga	2640
agacttaaag	gagttgtcac	ccactgaagg	aaggagtctc	caagagcttt	tagattaccc	2700
cggggaggat	gtggaggaga	ctttctgcct	caacttcacg	atctgccgag	aaagctatgg	2760
agtgattgaa	cagaagaagc	tgataacctg	gggagataat	gtaactgtgt	gcaaggataa	2820
caggcaggaa	tttgtggatg	cttatgtgaa	ttatgtcttc	caaactctcag	ttcatgaatg	2880
gtacacagcc	ttctctagt	gcttcctaaa	ggtgtgtggt	ggcaaagtac	ttgagctctt	2940

ccagccttca gaactgaggg ctatgatggt ggggaacagc aactacaact gggaagaact 3000
 ggaagagact gccatctaca agggagatta ctcggccaca catcccactg taaaactatt 3060
 ttgggaaaca tttcatgagt ttccattgga aaagaagaag aagtttctct tgttcctgac 3120
 aggcagcgat cggattccca tctacggcat ggccagtctg cagattgtca tccagtccac 3180
 agccagcggg gaggagtact tgccggtggc ccacacttgc tacaaccttc ttgacctccc 3240
 caagtacagc agcaaagaga ttctgagtgc ccggctgacc caggcccttg acaactatga 3300
 agggtttagt ttggcctgag gcttctcagc ttgtccagta tttcccttcg ttcctcagt 3360
 tccacattga ggcctataca gaaaatcatg gggagtgatt tctatTTTTT tattgtctaa 3420
 gtgggttggg actttttaaat actgagcctg gttgatgtgt ttctgggatt gtatagcagt 3480
 aaacaacctt tttgaaaaat tagaggttgg ggatgggggtg aaaaattggc ccttgtatgg 3540
 gaggtgtttt tgtttttgtt ttaaaccaaa ctaccagta ttccctgcac ttgtgaatgt 3600
 gttgcactct gctggatgaa atggcagtgg atttttaaac tttaatttcc caaatgtctc 3660
 tctcagccct gatgttttct cacagtgtt ccttgtcctt ctcttaactt ctcattcctc 3720
 tataagaatg atttagactg acctgtcctt ttttatctgc gcatgcgaga acatcacctt 3780
 cctctgtaca cttggaaatg cctctggcct gttgcagccc tcccttaacc caaaggagga 3840
 aaggactgct tcagaaactc ccaattccaa aaagctgagt ctgggtccat tattttggca 3900
 gaactcctaa gaatttatgg gagcctatat aaacatatct tgcttttaaa aagttcttga 3960
 gggaatagca actttcccat ggctgtgcct atttcctaga ccttttaaaa gatgtgcaga 4020
 gcagcttagc attcgttgca gctgagccta attttttctt gctcatcctt gtccctttga 4080
 caataagggt aattgataga cccaccacct cttgcactct cgcttttgga gcaagttgca 4140
 ttaactatTT tgagtctcta tattgtccaa gaaaagtaga aataataaat ttactttccc 4200
 tttttctatc accttatgtc ctctaccatt ttctccttcc tcccttccct tattttctcc 4260
 ttttctgacc ctgtgtcctc cctgattttc ctttctgttc ttctttatTT tatcccatc 4320
 tctgttactt gactcagtgc tcccttctc tccctcctt ctagtggatg catgcagcct 4380
 ttttttcaat tttatttaa attgcaaaat tttactcag attttttttc ctcttccta 4440
 attgctaaga ttaaggacg ttctttatta tgaaacttta tcacattcga aatgtttgtt 4500
 tacagtggga ttttaggggg gattgtgttt aaatcaaata tatgtatTTT aaaaataatg 4560
 acatgctcaa ccttcctcat catggagtaa gaaaattcta catgattaaa gaatccatgt 4620
 aagtctaatt ttaaattcct agtaactaga gaaaagactt atttatataa aatgaagtat 4680
 ttatgaactg tgataaagca tcaaatcttg atgaaggatt gtagatTTTt gctttttctt 4740
 tttgttttta aaacttatTC caattgctaa attggtagtt tttcagtctt tataaatata 4800
 ggattaaaaa tatatatata gttatatgaa atgtttatTT tctatgtgtg tgcatatagt 4860

tcaatattat gcaataaaatt tggtgtttta actt

4894

<210> 28
 <211> 1050
 <212> PRT
 <213> Homo sapiens

<400> 28

Met Leu Cys Trp Gly Tyr Trp Ser Leu Gly Gln Pro Gly Ile Ser Thr
 1 5 10 15

Asn Leu Gln Gly Ile Val Ala Glu Pro Gln Val Cys Gly Phe Ile Ser
 20 25 30

Asp Arg Ser Val Lys Glu Val Ala Cys Gly Gly Asn His Ser Val Phe
 35 40 45

Leu Leu Glu Asp Gly Glu Val Tyr Thr Cys Gly Leu Asn Thr Lys Gly
 50 55 60

Gln Leu Gly His Glu Arg Glu Gly Asn Lys Pro Glu Gln Ile Gly Ala
 65 70 75 80

Leu Ala Asp Gln His Ile Ile His Val Ala Cys Gly Glu Ser His Ser
 85 90 95

Leu Ala Leu Ser Asp Arg Gly Gln Leu Phe Ser Trp Gly Ala Gly Ser
 100 105 110

Asp Gly Gln Leu Gly Leu Met Thr Thr Glu Asp Ser Val Ala Val Pro
 115 120 125

Arg Leu Ile Gln Lys Leu Asn Gln Gln Thr Ile Leu Gln Val Ser Cys
 130 135 140

Gly Asn Trp His Cys Leu Ala Leu Ala Ala Asp Gly Gln Phe Phe Thr
 145 150 155 160

Trp Gly Lys Asn Ser His Gly Gln Leu Gly Leu Gly Lys Glu Phe Pro
 165 170 175

Ser Gln Ala Ser Pro Gln Arg Val Arg Ser Leu Glu Gly Ile Pro Leu
 180 185 190

Ala Gln Val Ala Ala Gly Gly Ala His Ser Phe Ala Leu Ser Leu Ser
 195 200 205

Gly Ala Val Phe Gly Trp Gly Met Asn Asn Ala Gly Gln Leu Gly Leu

210	215	220
Ser Asp Glu Lys Asp Arg Glu Ser Pro Cys His Val Lys Leu Leu Arg 225 230 235 240		
Thr Gln Lys Val Val Tyr Ile Ser Cys Gly Glu Glu His Thr Ala Val 245 250 255		
Leu Thr Lys Ser Gly Gly Val Phe Thr Phe Gly Ala Gly Ser Cys Gly 260 265 270		
Gln Leu Gly His Asp Ser Met Asn Asp Glu Val Asn Pro Arg Arg Val 275 280 285		
Leu Glu Leu Met Gly Ser Glu Val Thr Gln Ile Ala Cys Gly Arg Gln 290 295 300		
His Thr Leu Ala Phe Val Pro Ser Ser Gly Leu Ile Tyr Ala Phe Gly 305 310 315 320		
Cys Gly Ala Arg Gly Gln Leu Gly Thr Gly His Thr Cys Asn Val Lys 325 330 335		
Cys Pro Ser Pro Val Lys Gly Tyr Trp Ala Ala His Ser Gly Gln Leu 340 345 350		
Ser Ala Arg Ala Asp Arg Phe Lys Tyr His Ile Val Lys Gln Ile Phe 355 360 365		
Ser Gly Gly Asp Gln Thr Phe Val Leu Cys Ser Lys Tyr Glu Asn Tyr 370 375 380		
Ser Pro Ala Val Asp Phe Arg Thr Met Asn Gln Ala His Tyr Thr Ser 385 390 395 400		
Leu Ile Asn Asp Glu Thr Ile Ala Val Trp Arg Gln Lys Leu Ser Glu 405 410 415		
His Asn Asn Ala Asn Thr Ile Asn Gly Val Val Gln Ile Leu Ser Ser 420 425 430		
Ala Ala Cys Trp Asn Gly Ser Phe Leu Glu Lys Lys Ile Asp Glu His 435 440 445		
Phe Lys Thr Ser Pro Lys Ile Pro Gly Ile Asp Leu Asn Ser Thr Arg 450 455 460		

Val Leu Phe Glu Lys Leu Met Asn Ser Gln His Ser Met Ile Leu Glu
 465 470 475 480

Gln Ile Leu Asn Ser Phe Glu Ser Cys Leu Ile Pro Gln Leu Ser Ser
 485 490 495

Ser Pro Pro Asp Val Glu Ala Met Arg Ile Tyr Leu Ile Leu Pro Glu
 500 505 510

Phe Pro Leu Leu Gln Asp Ser Lys Tyr Tyr Ile Thr Leu Thr Ile Pro
 515 520 525

Leu Ala Met Ala Ile Leu Arg Leu Asp Thr Asn Pro Ser Lys Val Leu
 530 535 540

Asp Asn Trp Trp Ser Gln Val Cys Pro Lys Tyr Phe Met Lys Leu Val
 545 550 555 560

Asn Leu Tyr Lys Gly Ala Val Leu Tyr Leu Leu Arg Gly Arg Lys Thr
 565 570 575

Phe Leu Ile Pro Val Leu Phe Asn Asn Tyr Ile Thr Ala Ala Leu Lys
 580 585 590

Leu Leu Glu Lys Leu Tyr Lys Val Asn Leu Lys Val Lys His Val Glu
 595 600 605

Tyr Asp Thr Phe Tyr Ile Pro Glu Ile Ser Asn Leu Val Asp Ile Gln
 610 615 620

Glu Asp Tyr Leu Met Trp Phe Leu His Gln Ala Gly Met Lys Ala Arg
 625 630 635 640

Pro Ser Ile Ile Gln Asp Thr Val Thr Leu Cys Ser Tyr Pro Phe Ile
 645 650 655

Phe Asp Ala Gln Ala Lys Thr Lys Met Leu Gln Thr Asp Ala Glu Leu
 660 665 670

Gln Met Gln Val Ala Val Asn Gly Ala Asn Leu Gln Asn Val Phe Met
 675 680 685

Leu Leu Thr Leu Glu Pro Leu Leu Ala Arg Ser Pro Phe Leu Val Leu
 690 695 700

His Val Arg Arg Asn Asn Leu Val Gly Asp Ala Leu Arg Glu Leu Ser
 705 710 715 720

Ile His Ser Asp Ile Asp Leu Lys Lys Pro Leu Lys Val Ile Phe Asp
 725 730 735
 Gly Glu Glu Ala Val Asp Ala Gly Gly Val Thr Lys Glu Phe Phe Leu
 740 745 750
 Leu Leu Leu Lys Glu Leu Leu Asn Pro Ile Tyr Gly Met Phe Thr Tyr
 755 760 765
 Tyr Gln Asp Ser Asn Leu Leu Trp Phe Ser Asp Thr Cys Phe Val Glu
 770 775 780
 His Asn Trp Phe His Leu Ile Gly Ile Thr Cys Gly Leu Ala Ile Tyr
 785 790 795 800
 Asn Ser Thr Val Val Asp Leu His Phe Pro Leu Ala Leu Tyr Lys Lys
 805 810 815
 Leu Leu Asn Val Lys Pro Gly Leu Glu Asp Leu Lys Glu Leu Ser Pro
 820 825 830
 Thr Glu Gly Arg Ser Leu Gln Glu Leu Leu Asp Tyr Pro Gly Glu Asp
 835 840 845
 Val Glu Glu Thr Phe Cys Leu Asn Phe Thr Ile Cys Arg Glu Ser Tyr
 850 855 860
 Gly Val Ile Glu Gln Lys Lys Leu Ile Pro Gly Gly Asp Asn Val Thr
 865 870 875 880
 Val Cys Lys Asp Asn Arg Gln Glu Phe Val Asp Ala Tyr Val Asn Tyr
 885 890 895
 Val Phe Gln Ile Ser Val His Glu Trp Tyr Thr Ala Phe Ser Ser Gly
 900 905 910
 Phe Leu Lys Val Cys Gly Gly Lys Val Leu Glu Leu Phe Gln Pro Ser
 915 920 925
 Glu Leu Arg Ala Met Met Val Gly Asn Ser Asn Tyr Asn Trp Glu Glu
 930 935 940
 Leu Glu Glu Thr Ala Ile Tyr Lys Gly Asp Tyr Ser Ala Thr His Pro
 945 950 955 960
 Thr Val Lys Leu Phe Trp Glu Thr Phe His Glu Phe Pro Leu Glu Lys
 965 970 975

Lys Lys Lys Phe Leu Leu Phe Leu Thr Gly Ser Asp Arg Ile Pro Ile
 980 985 990

Tyr Gly Met Ala Ser Leu Gln Ile Val Ile Gln Ser Thr Ala Ser Gly
 995 1000 1005

Glu Glu Tyr Leu Pro Val Ala His Thr Cys Tyr Asn Leu Leu Asp
 1010 1015 1020

Leu Pro Lys Tyr Ser Ser Lys Glu Ile Leu Ser Ala Arg Leu Thr
 1025 1030 1035

Gln Ala Leu Asp Asn Tyr Glu Gly Phe Ser Leu Ala
 1040 1045 1050

<210> 29

<211> 4363

<212> DNA

<213> Homo sapiens

<400> 29

```

gagctggaat tcccttcccc cctggcgccg cccgggagag cccgcgcgag gacgcccgcc 60
gcgctccgcc ggcccttttt tggcgctgag ggaaaggaga agggcagggc cgccgcctcg 120
cgggagcggg ccccgaggag agggggccgc ggagaggccg ggcgagcgcg ggcggccctc 180
acctcgccgc tcctcccggg ccgccatccc tcggcgcccc gcccggaacc ggcgcgcgcg 240
taggggctgg gaggccggcg cgcagatctg gcggtgagcg ctgccgcccc ggggccccca 300
gcatgtcgg ccgaggagat ggtgcagatc cgcctggagg accgctgcta cccggtgagc 360
aagaggaagc tcatcgagca gagcgactac ttccgcgcc tctaccgctc cggcatgcgc 420
gaggccctga gccaggaggc cggcgggccc gaggtgcagc agctgcgcgg cctcagcgcg 480
ccgggcctgc ggctggtgct ggacttcata aacgcggcg gggcccgcga aggctggctc 540
ctgggccccg gcggggaaaa gggcggcggg gtggacgagg acgaggagat ggatgaggtg 600
agcctgctgt ccgagctggt ggaggcgcc tccttctctc aggtcacgtc cctgctgcag 660
ctgctgctgt cccaggtgcg gctcaataac tgccctggaga tgtaccgcct ggcgcaggtg 720
tacgggctgc ccgacctgca ggaggcctgc ctgcgcttca tggctgtcca cttccacgag 780
gtgctgtgca agccccagtt ccacctctg gggctctctc cccaagctcc aggggatgtc 840
agcctgaagc agaggctgag ggaggccccg atgactggga ctctgtcct cgtggccctc 900
ggggacttcc tggggggacc cctggcccc caccctacc agggggagcc cccgtccatg 960
ctcaggtacg aggagatgac tgagcgttgg ttcccgtgg ccaacaacct tcctcccagc 1020
ctggtcaatg tcaggggcta tgggtctgcc atcctggaca actaccttt catagtgggc 1080
gggtacagga tcactagcca ggagatctcc gctgcgcatt cctacaaccc cagcaccaac 1140

```

gagtggctcc	aggtggcctc	catgaaccag	aagaggtcta	acttcaaact	tgtggctggt	1200
aattcaaaac	tctatgccat	eggagggcag	gccgtttcta	acgttgagtg	ttacaacccc	1260
gagcaggatg	cgtggaattt	tgtggcgccc	ttaccaatc	ctctggctga	gttctctgcc	1320
tgtgagtgtg	agggaaaaat	ttatgtcatt	ggaggatata	ctaccagaga	ccggaacatg	1380
aacattttgc	agtactgccc	ctcttccgac	atgtggacgc	tctttgaaac	atgtgacgtc	1440
cacattcgca	agcagcagat	ggtgtctgtg	gaagagacca	tctacatcgt	gggggggtgt	1500
ctccacgagc	tgggggccaa	ccgcaggagc	agccagagcg	aggacatgct	caccgtgcag	1560
tcctacaaca	ccgtcacccg	ccagtggctc	tacctcaagg	agaacacgtc	caaatcgggt	1620
cttaacttga	cttgtgcgct	ccataacgac	ggcatctata	tcatgagcag	agacgtcacc	1680
ctgtcgacca	gcttgaaca	ccgagtgttc	ctcaagtata	acatcttttc	agatagttgg	1740
gaagcatttc	ggcgttttcc	agcttttggg	cataacttgc	tggtttcttc	tctttatctg	1800
cccaataaag	cagaaacatg	actgaattga	attggtagat	gaaaaaaaacc	tggttcaagt	1860
tttttttaaa	atgtggtgtc	ccattccaag	ggagaccaat	tcctaaaggg	taaagaaggg	1920
ttaaagtagg	tcacatatat	acagtagcag	ctgtaaataa	gcttacttga	actaattact	1980
tgaaaactgg	tggaaaaaag	agaccaactt	tgttattttt	taaattattg	atttttaaat	2040
tatgggagca	aatccatctc	caaacatgat	actcttgggc	caatgacgga	tcatcaaaat	2100
gtagattcat	tgtgtctgtg	tggttaaaggg	ccaaagtcaa	taattgggac	aaatggtaaa	2160
gatgatttta	aaatctgttg	tatttaggtc	ccttgacata	tcattagcac	cttaagttag	2220
aggttttctt	tttcttatca	gtgtccttga	caagcattca	tattaacata	ttcattctct	2280
cttctccatc	accatacccc	acacatacat	ctagtatggt	tagggttctg	tggtcagata	2340
ttttgtttgc	acactacttt	ttaggaaact	tataatgatg	gagccctaca	gtgaatcctt	2400
accagtgttc	agtcctcccc	aaaacttgca	taaaaatatt	cttagctatt	ttccagcct	2460
tttttgcagt	tacaaatgct	ggataaatag	actgataaaa	cctcttattt	tatttatgtg	2520
gcaggttgca	tatttatgtg	gacacatagg	ttgcaactgt	aggcattctg	tattgtttta	2580
aagaggatat	attattttaga	aaaatggaca	tatgaatatt	attaaaaatt	gagtcttaaa	2640
tttaaatggg	aagggaagaa	aacaccaaact	aagaaaaaaa	tttgccttta	agagttactt	2700
ttgttgaatg	tatttgactt	agttcttatt	ttaaggctaa	aaaaactttg	tcatatttaa	2760
gtcggcagct	ttctaattta	atcttagctt	tgtaattcat	gttttgcttt	tgtggactaa	2820
ggtaaggaca	gcattttaa	atgtgtgctg	gagtgagggg	aagaagctgt	tacagaagtg	2880
gaatggtttc	tggtggtatt	tctgaagtta	ggtaaagtgc	tttacttgcc	ttgggtgtta	2940
attgaaatag	aattcagctc	aagcagttgt	tttcagtgag	tgagttaaca	ataaaagctt	3000
ctgccaatca	tagatcaaca	ttcctcaaaa	atacatcac	tcatacttat	ggaaatttta	3060

```

ttttgattga aaacactctg aaattatctg atatttttga atggcatcag ttaaactgtg 3120
agctggattc catcttttctg cagtaacagc gggcagaacc atttcagtta gcatgtgcct 3180
cactttaaga aagtcctgta ggggaaaaaa aaaactgtca gagcgaattc atgcattcaa 3240
caaacactta tgagtgcctg gagtatgccg tccactgtgc agatgtagga caggggctaa 3300
tcacttagag ttcttcagtt tattagagga gtgactacag gttttcttta caaccatggc 3360
ctctagtata cttaaaataa gcgtccagtt ataaaaagcc tatttgcaca actttttggg 3420
aataacttcta ttaactaaaa tgaggtaagc cattattttg gacatagaaa cagatctttg 3480
tatcttgtca ccaaataatc ttacctctag tctacttaca actttgtaag ggtttctaag 3540
ctatgccccat ttattcccaa atattaagcc cttaaaaaaa taaaacctct gaacccaaat 3600
cttcccagga atagtactta atagatttta atgttaattt atattcattg tagaattgaa 3660
tttccagtgga cctgtaattt cattacatat attttaaaat ctatttattt tagatggtgt 3720
taacttttctg ttgctgaaga gcttaacttt ttaaaggctt tgtcctatac atttagaaga 3780
tgagtcagtt gaattagtac tggacaaaca gttggagatt agtttgcaaa taagcatatc 3840
atatcaaata caatattcca gcccaaccagt taattctctt cctgggttaca ttcattgggt 3900
ctattttgcct aatgttgaca attagattct tggaccaagt gaggggtgtac tctgatccac 3960
aaagctcttt tataaaccag gtttgagggt gtgtgtgtgt gtgggggtgt gtgtgtgtgt 4020
gtgtgtgtat gtttgcata tatagctctc tttaagacag acatctaaaa agattttggt 4080
tattctgggc tgccagcagt acgtgggaaa ggtggaatgt gcaataggaa cagatctgtg 4140
cattttgtga catcgtcctt cctgctgtgc acatcagctg tgatgacatt ttgagggtag 4200
ttttaattta gctcggttct cttgttggtg ccagtattac attgaagact acctatgata 4260
gaggcaatgt tgctttgaaa atctcccttc aggcggaggt tgcagtgagc cgagattgtg 4320
ccattgcact ccagcctggg cgacagtgca agactctgtc tcc 4363

```

<210> 30
 <211> 505
 <212> PRT
 <213> Homo sapiens

<400> 30

Met Ser Ala Glu Glu Met Val Gln Ile Arg Leu Glu Asp Arg Cys Tyr
 1 5 10 15

Pro Val Ser Lys Arg Lys Leu Ile Glu Gln Ser Asp Tyr Phe Arg Ala
 20 25 30

Leu Tyr Arg Ser Gly Met Arg Glu Ala Leu Ser Gln Glu Ala Gly Gly
 35 40 45

Pro Glu Val Gln Gln Leu Arg Gly Leu Ser Ala Pro Gly Leu Arg Leu
 50 55 60

Val Leu Asp Phe Ile Asn Ala Gly Gly Ala Arg Glu Gly Trp Leu Leu
 65 70 75 80

Gly Pro Arg Gly Glu Lys Gly Gly Gly Val Asp Glu Asp Glu Glu Met
 85 90 95

Asp Glu Val Ser Leu Leu Ser Glu Leu Val Glu Ala Ala Ser Phe Leu
 100 105 110

Gln Val Thr Ser Leu Leu Gln Leu Leu Leu Ser Gln Val Arg Leu Asn
 115 120 125

Asn Cys Leu Glu Met Tyr Arg Leu Ala Gln Val Tyr Gly Leu Pro Asp
 130 135 140

Leu Gln Glu Ala Cys Leu Arg Phe Met Val Val His Phe His Glu Val
 145 150 155 160

Leu Cys Lys Pro Gln Phe His Leu Leu Gly Ser Pro Pro Gln Ala Pro
 165 170 175

Gly Asp Val Ser Leu Lys Gln Arg Leu Arg Glu Ala Arg Met Thr Gly
 180 185 190

Thr Pro Val Leu Val Ala Leu Gly Asp Phe Leu Gly Gly Pro Leu Ala
 195 200 205

Pro His Pro Tyr Gln Gly Glu Pro Pro Ser Met Leu Arg Tyr Glu Glu
 210 215 220

Met Thr Glu Arg Trp Phe Pro Leu Ala Asn Asn Leu Pro Pro Asp Leu
 225 230 235 240

Val Asn Val Arg Gly Tyr Gly Ser Ala Ile Leu Asp Asn Tyr Leu Phe
 245 250 255

Ile Val Gly Gly Tyr Arg Ile Thr Ser Gln Glu Ile Ser Ala Ala His
 260 265 270

Ser Tyr Asn Pro Ser Thr Asn Glu Trp Leu Gln Val Ala Ser Met Asn
 275 280 285

Gln Lys Arg Ser Asn Phe Lys Leu Val Ala Val Asn Ser Lys Leu Tyr
 290 295 300

Ala Ile Gly Gly Gln Ala Val Ser Asn Val Glu Cys Tyr Asn Pro Glu
305 310 315 320

Gln Asp Ala Trp Asn Phe Val Ala Pro Leu Pro Asn Pro Leu Ala Glu
325 330 335

Phe Ser Ala Cys Glu Cys Lys Gly Lys Ile Tyr Val Ile Gly Gly Tyr
340 345 350

Thr Thr Arg Asp Arg Asn Met Asn Ile Leu Gln Tyr Cys Pro Ser Ser
355 360 365

Asp Met Trp Thr Leu Phe Glu Thr Cys Asp Val His Ile Arg Lys Gln
370 375 380

Gln Met Val Ser Val Glu Glu Thr Ile Tyr Ile Val Gly Gly Cys Leu
385 390 395 400

His Glu Leu Gly Pro Asn Arg Arg Ser Ser Gln Ser Glu Asp Met Leu
405 410 415

Thr Val Gln Ser Tyr Asn Thr Val Thr Arg Gln Trp Leu Tyr Leu Lys
420 425 430

Glu Asn Thr Ser Lys Ser Gly Leu Asn Leu Thr Cys Ala Leu His Asn
435 440 445

Asp Gly Ile Tyr Ile Met Ser Arg Asp Val Thr Leu Ser Thr Ser Leu
450 455 460

Glu His Arg Val Phe Leu Lys Tyr Asn Ile Phe Ser Asp Ser Trp Glu
465 470 475 480

Ala Phe Arg Arg Phe Pro Ala Phe Gly His Asn Leu Leu Val Ser Ser
485 490 495

Leu Tyr Leu Pro Asn Lys Ala Glu Thr
500 505

<210> 31

<211> 2501

<212> DNA

<213> Homo sapiens

<400> 31

gttgctggct ttttcaggag ctgctcgctc gcagccagag acgctgcttt ttttttccgg 60

gttcggagcc gttccggatg ctttaggctg ccggatgtct gatctccgaa taactgagggc 120

gtttctgtac atggattatc tgtgttttag agcactttgc tgcaagggac caccacctgc 180

acgaccagaa tatgacctgg ttgcatagg cctcacaggt tctggcaaaa ccagtctgtt	240
gtccaaactc tgcagtgaag gccccgataa cgtcgtgtcg accacagggt ttagtattaa	300
agcagtgccca ttccagaatg ccatcttgaa tgtaaaagaa cttggagggg ctgataacat	360
ccggaaatac tggagccgct actaccaagg atctcaagg gtaatatctt tattagacag	420
tgcctcttca gaggatgatt tagaagctgc tagaaatgag ctgcactcag ctcttcagca	480
tccacagtta tgcactttac cctttttaat attggccaat catcaagaca agccagcagc	540
tcgctcagta caagagatca aaaaatattt tgaacttgaa ccacttgac gtggaaaacg	600
ctggattcta cagccctgct cactggatga catggatgca ctgaaagaca gcttctctca	660
gctgattaat ttgttagaag aaaaagacca tgaagctgta agaattgtgaa atctggcaaa	720
gaaaacaggc tcccaaaagg ccttgaacct atcatgttag tttctcattt taattttatt	780
ttggtatcaa gactatattg gcttcagtat ctattttcct ctggtcattt cagactctca	840
tctctctatt aaagtgaata ttctgtagat cagatgaatt atcatcggca ttaacgtcaa	900
gtacacacta ctgagagaga atttattctc tgtttaccac taattatctt ccccgcatgt	960
ctcttccttc tgttgttcta agcatctctt ctatactctg tgctgaaacg gatggaattc	1020
atggccttatg gaaatcaaatt ccttttgaa ggactcagaa agactcaca gaccttgatc	1080
atcttctctg ggtttgggag tggagtatta tgtgactggg actccacaaa atatttatat	1140
tattttcaag acctattggg ggttttggtt tattttatct aattttttt ttttttggt	1200
atactgagct tatagactat gagagaaaag tgagcaccaa aaacttcatt aatcacctta	1260
tcaggaagga atatcctgcc atagtaaaat tggctctgaa gagcttagag gcaagttaca	1320
caatgttttc ttgattacat ctgaagtctc ttctctttta agagcaaaga cattgtttca	1380
gtttctgaga gtggtttagag gaaatttagg tacaatgcat gcatcacaga ctgtatttga	1440
ttgattcgag tcttcaaata ttgtttaatg gaatgatgag cctgcataaa gtgaagctgc	1500
aatgtaacat aggtatcttt ttctaaaaac ttcaagtgtt tgattaaaat agagaaaata	1560
atttttaaaa taatagataa aaatttttaa aacagccaga aaaagtataa tttatactgg	1620
ttagggttta agatggcatg gatttgacaa agataacaaa gacagaatca tctatatatt	1680
cattcattac gtgttactgt agtttattgg aatctgttaa tgccttctc tagtagttaa	1740
aggtttatca gtatttctat tataaacctc tattttcagg ggtttcaatg tgcccttaaa	1800
attgtgctgg ggctgcagca aataaataat attgggactc atattcagtc tttggaacta	1860
gttttttttc catttaattt aaatgacaaa attgcatagg gataaatgta cagttttact	1920
aaggtttagc acctttgatc ttttctaaat caggaaaaaa atctatgcat attgaggttc	1980
acagatgaaa tattcttttg catactttgt agactttgac cacttaaaat actaagtggc	2040
caaatgataa aaaaaactgc tccctgcaca tgagtgtaaa aacatgtttg taattttaag	2100

taaaatggac tgatacccaa ataccacaga ctcacactat attacagaat gatgtaatac 2160
 ctggtaattc tccagaagat agacttcaga cccttaatgc catttaaata tagtttttgt 2220
 ggtttaaatct tatgagtaga ttgtaatgaa atcattacct ttgaaacatc aaacacttct 2280
 ttaagctttt tttttttaat gtcatagaatc aggaaggata catagtgatt cgtatacaat 2340
 tttgtctgca gcattcttga atatttttaa cactgtgtaa tccctgagca gcacaaaact 2400
 tttattccga tgaccagtta cacaattcct gtttttattt tgtcctgttg tgctcatctt 2460
 ctatgaaaaa taaatgatta ttaggcaaaa aaaaaaaaaa a 2501

<210> 32
 <211> 204
 <212> PRT
 <213> Homo sapiens
 <400> 32

Met Ser Asp Leu Arg Ile Thr Glu Ala Phe Leu Tyr Met Asp Tyr Leu
 1 5 10 15

Cys Phe Arg Ala Leu Cys Cys Lys Gly Pro Pro Pro Ala Arg Pro Glu
 20 25 30

Tyr Asp Leu Val Cys Ile Gly Leu Thr Gly Ser Gly Lys Thr Ser Leu
 35 40 45

Leu Ser Lys Leu Cys Ser Glu Ser Pro Asp Asn Val Val Ser Thr Thr
 50 55 60

Gly Phe Ser Ile Lys Ala Val Pro Phe Gln Asn Ala Ile Leu Asn Val
 65 70 75 80

Lys Glu Leu Gly Gly Ala Asp Asn Ile Arg Lys Tyr Trp Ser Arg Tyr
 85 90 95

Tyr Gln Gly Ser Gln Gly Val Ile Phe Val Leu Asp Ser Ala Ser Ser
 100 105 110

Glu Asp Asp Leu Glu Ala Ala Arg Asn Glu Leu His Ser Ala Leu Gln
 115 120 125

His Pro Gln Leu Cys Thr Leu Pro Phe Leu Ile Leu Ala Asn His Gln
 130 135 140

Asp Lys Pro Ala Ala Arg Ser Val Gln Glu Ile Lys Lys Tyr Phe Glu
 145 150 155 160

Leu Glu Pro Leu Ala Arg Gly Lys Arg Trp Ile Leu Gln Pro Cys Ser

	165	170	175	
Leu Asp Asp Met Asp Ala Leu Lys Asp Ser Phe Ser Gln Leu Ile Asn				
	180	185	190	
Leu Leu Glu Glu Lys Asp His Glu Ala Val Arg Met				
	195	200		

<210> 33
 <211> 2334
 <212> DNA
 <213> Homo sapiens

<400> 33
 ctttatggcg tgggagaggc cacagcccg actccatcga ctccccggc tcttagacta 60
 aaatcatgcc caagttcaaa caacgaagac gaaagctaaa agccaaagcc gaaagattat 120
 tcaaaaaaaaa agaagcctct cactttcagt ccaagctaata tacacctcct cctccaccac 180
 cctcaccaga aagagtcggg atttcttcaa tagatatatc tcaaagcaga agctgggctaa 240
 catcatcctg gaacttcaat tttcctaaca tcagagatgc aataaaactt tggacaaata 300
 gagtatgggc tatatacagc tggtgccaga actgcataac ccagagttaa gaagtattga 360
 aagacaccat ctttccatct cgtatctgcc accgagaact ttacagtgtg aaacaacagt 420
 tttgcatttt ggaaagtaaa ttatgcaagc ttcaggaagc actgaagacc atctcagaaa 480
 gttcttctctg tccaagctgt ggtcaaacat gtcacatgag tggtaaactt acaaatgtgc 540
 ctgcctgcgt tctgatcacc cctggagact ccaaagctgt gcttctccc acactgccac 600
 agccagccag ccatttttct cctcctctc cacctccacc tctgccacct cctccaccac 660
 cactagcacc tgtgttgctc agaaaaccca gtctcgctaa agcacttcag gctggaccat 720
 taaaaaaaga tggacccatg cagataacag ttaaagatct gctgactgta aaattaaaga 780
 agacacagag tttagatgaa aagaggaagc ttataccatc gccgaaagca cggaatccac 840
 tagttaccgt ctctgacttg cagcatgtta ccctgaaacc taactccaaa gtgttatcga 900
 ctcgagttac aaacgtctta atcactcctg gaaaaagtca gatggatctg cggaaaactgc 960
 ttagaaaagt cgatgtagag aggaggccag gtggaacccc tcttaccaat aaggaaaata 1020
 tggaaacagg aacaggactg actccagtga tgacgcaggc ctttaaggaga aagtttcagc 1080
 tggctcacc tagaagcca actccaactc tgccactttc tacaagcagc tttgatgaac 1140
 aaaactgatg ccaactctgc ctactccat gagatgatcc ataacaata cagataaaaa 1200
 caccagctg ggtgcagtgg ctgcacctg taatcccagc actttgggag gctgaggcag 1260
 gtggatcacc tgaggtcggg agtttgagac cagcctggcc aacatgggtga aatcccctct 1320
 ctactgaaaa tacagaattg gccgggtgtg gtgtcacgcg cctgtggtcc cagttacttg 1380

```

aacctgggag gcggaggttg cagtgagccg agattgcaac actgccactc cagcctgggt 1440
gacagagcaa gagaccctgt ctcaaaaaaa aaaaaaaaaa aattaaatgg gtagtgacgt 1500
taagagatat atatcagctt ctagtaaaag tttttttttt taaacctgct agctacattt 1560
acattatgta aaaataaatg gaataatcac tgagaataaa gcagttgagt atttataaca 1620
ataatatatt atggggcgct tataatgttt ataatatgtt aaaccactgt gtactctatt 1680
catttaatgc taaatgactt gaccattctt gtgggataag agatcattaa aaaaatgcta 1740
gggccgggca ccatggctca cgctgtaat ccaggactt taggaggcca aggcaggtgg 1800
atgacttgag ctcaggagtt tgagaccagc ctgggcaaca tgatgaaaac tccgtctcta 1860
ccaaaaatca aaaaaattag ccaggtgtga tgttgtgcgc ctgtagtccc agctacttgt 1920
gaggctgagg tgggaggatg gtttgggccc cggaggtaga ggttgacgtg agctgagatt 1980
gtgccactgt attccatcct gggcaacaga gccagaccct gcctcaaaaa aaaaaaaaaa 2040
aaaaaaaaagg cctggcacag tggctcacgc ctgtaatccc agcactttgg gaggctgagg 2100
cgggcaaaac acttgaggtc agacgtttga gaccagcctg gccaacatgg tgaaaccccc 2160
tctctaccaa aaataaaaaat aaaaattatc caggcgtggt ggcatgtgcc tatagcccca 2220
gctactcagg aagctagagc acgagaatta tgtgcatcct ggaggtgaag gttgcagtga 2280
gccaatgca caccactcca ctccagtctg gacaacagag tgagaccctg tgtc 2334

```

```

<210> 34
<211> 360
<212> PRT
<213> Homo sapiens

```

```

<400> 34

```

```

Met Pro Lys Phe Lys Gln Arg Arg Arg Lys Leu Lys Ala Lys Ala Glu
1          5          10          15

```

```

Arg Leu Phe Lys Lys Lys Glu Ala Ser His Phe Gln Ser Lys Leu Ile
20          25          30

```

```

Thr Pro Pro Pro Pro Pro Pro Ser Pro Glu Arg Val Gly Ile Ser Ser
35          40          45

```

```

Ile Asp Ile Ser Gln Ser Arg Ser Trp Leu Thr Ser Ser Trp Asn Phe
50          55          60

```

```

Asn Phe Pro Asn Ile Arg Asp Ala Ile Lys Leu Trp Thr Asn Arg Val
65          70          75          80

```

```

Trp Ser Ile Tyr Ser Trp Cys Gln Asn Cys Ile Thr Gln Ser Leu Glu
85          90          95

```

Val Leu Lys Asp Thr Ile Phe Pro Ser Arg Ile Cys His Arg Glu Leu
 100 105 110

Tyr Ser Val Lys Gln Gln Phe Cys Ile Leu Glu Ser Lys Leu Cys Lys
 115 120 125

Leu Gln Glu Ala Leu Lys Thr Ile Ser Glu Ser Ser Ser Cys Pro Ser
 130 135 140

Cys Gly Gln Thr Cys His Met Ser Gly Lys Leu Thr Asn Val Pro Ala
 145 150 155 160

Cys Val Leu Ile Thr Pro Gly Asp Ser Lys Ala Val Leu Pro Pro Thr
 165 170 175

Leu Pro Gln Pro Ala Ser His Phe Pro Pro Pro Pro Pro Pro Pro Pro
 180 185 190

Leu Pro Pro Pro Pro Pro Pro Leu Ala Pro Val Leu Leu Arg Lys Pro
 195 200 205

Ser Leu Ala Lys Ala Leu Gln Ala Gly Pro Leu Lys Lys Asp Gly Pro
 210 215 220

Met Gln Ile Thr Val Lys Asp Leu Leu Thr Val Lys Leu Lys Lys Thr
 225 230 235 240

Gln Ser Leu Asp Glu Lys Arg Lys Leu Ile Pro Ser Pro Lys Ala Arg
 245 250 255

Asn Pro Leu Val Thr Val Ser Asp Leu Gln His Val Thr Leu Lys Pro
 260 265 270

Asn Ser Lys Val Leu Ser Thr Arg Val Thr Asn Val Leu Ile Thr Pro
 275 280 285

Gly Lys Ser Gln Met Asp Leu Arg Lys Leu Leu Arg Lys Val Asp Val
 290 295 300

Glu Arg Arg Pro Gly Gly Thr Pro Leu Thr Asn Lys Glu Asn Met Glu
 305 310 315 320

Thr Gly Thr Gly Leu Thr Pro Val Met Thr Gln Ala Leu Arg Arg Lys
 325 330 335

Phe Gln Leu Ala His Pro Arg Ser Pro Thr Pro Thr Leu Pro Leu Ser
 340 345 350

Thr Ser Ser Phe Asp Glu Gln Asn
 355 360

<210> 35
 <211> 6743
 <212> DNA
 <213> Homo sapiens

<400> 35
 cgacggcggc ggccgtcctc atcccggcgc ttgagaggac gcggggctgc gcaaatggct 60
 tgtcccgcctc tcggtctgga agctcttcag cccctgcagc ccgagccgcc ccccgagccc 120
 gccttctccg aggcgagaa gtggattgag caagtaactg gcagaagttt tggtagataaa 180
 gattttcgga caggtttaga aaatggaatc ctccctctgcg agttgctgaa tgctataaag 240
 ccaggacttg ttaaaaagat caatagattg cctaccccca ttgcaggact ggacaatatt 300
 atcttattct tgagagggtg taaagagctc ggccttaaag aatctcaact ttttgacccg 360
 agtgacctcc aggatacatc caacagagta acagtcaaga gccttgatta tagtaggaag 420
 ctgaaaaatg tattagttac catttactgg ctgggaaaag cagcaaacag ctgcacatcc 480
 tacagcggaa cgacactaaa cctgaaggag tttgaaggat tggtggctca gatgcgaaag 540
 gacactgatg acattgaaag tcctaaacgc agtatccgag acagtggcta catcgactgc 600
 tgggattccg agcgagcga ctccctctct cctcctcgcc acggcagaga tgattccttc 660
 gacagcctgg attcctttgg ctctcgtctc cggcagacgc cttcaccaga tgtagtcctc 720
 aggggaagca gcgatgggag aggaagcgac tctgaatccg acttgccctca tcggaagctg 780
 ccagatgtga agaaggatga catgtctgca cggcggactt cccatggtga gccgaaatca 840
 gcagtgcctt ttaaccagta cctcccgaa aaaagcaatc agacggccta cgtccccgcg 900
 cctctgagaa agaagaaagc agagagagag gaataccgca agagctggag taccgccacc 960
 tccccgctgg gtggggagag gcccttcagc tttcctgaaa cgatagagga agaggggagt 1020
 gaagtggggg ctgctgggtga agacaacca gctggccaaa tgaatcctgg ttggaagcct 1080
 tccgatggtg ggtgtgaatt acctgacggc agtggttaagg agcacccttc ttcagacggg 1140
 gctgtggtgg caccagctcc caagtctgaa gaaaaagatg ctgctgagat ccaaaagcgc 1200
 aaaaggctag agcaagctgg aatcaaggct atgccagcag cacagcgctt tgccagtcaa 1260
 aaacaattaa gtgaagagaa agaagctatt cgtgatattg tccttcgcaa agaaaactct 1320
 ttctgacccc accaaccatgg caacgattca gaggcagaag gtgaagttgt gtgtcgactg 1380
 cctgatcttg agaaggatga ctttgctgca aggagagcaa ggatgaacca aaccaagcca 1440
 atggtgccat taaatcaact cctctatggc ccttatccaa agaaaggggc agagaaatca 1500
 gatggcagca aacagctctc aaagggaatc tcaaaaaaaaa gaagtctaga atataaaaga 1560
 aaccaggggc acacagaaga ggtgaagttg atagtgcct gtaacatgag ggctcaggaa 1620

agtgaacctg	tggaaggagg	actcaggaag	gtgccagatc	ttcacaagga	tgacctggcc	1680
cagcagagaa	ttcagggcag	ccttgcccct	caccgcgagc	ccccgagctt	cattacgctc	1740
tccaacataa	cagaagctga	cttggagacg	tgggagagac	tcaaagtgtc	agaaaaggcg	1800
agggatggag	atgttcagca	catctgtgct	tctgagcctt	ccccagaaat	taaagcagaa	1860
actgccattc	gtgatgactt	tgccaaccgc	aaagcaaggg	cctctaagaa	agcttccagc	1920
cccaggcaaa	agtttgtgca	ctttggggcca	gtgacggagc	tagatcagca	gaaatggaag	1980
cggttagca	ttggcaaggc	tgggcctaga	gaggatgaag	aagaagtcac	ctgtcatggc	2040
agcaagattc	aaatggactc	tgtgtctcct	gtctcagcgg	ccacttccag	cttaaagggc	2100
caccaaataat	ttaatcgaca	aaatgactgc	agaacaatga	attgtggccg	aggtgactat	2160
tgcagaaggg	cctcgtggct	ggctcctgtg	cggagctctc	aggaagagtg	ggtctgcagt	2220
ttgggcgagt	gcccaggggg	gacagaggaa	gttacatcca	aacagctgcc	acaggatggc	2280
aaagaagaaa	cagaaagcgc	tccaagagat	tctgagaggc	tgtcaaaggc	agaaagatca	2340
gaggacagca	gccagccact	ggtgtgtcct	ctggcctctg	agtgtgaggc	ttcagggaca	2400
gaagagaagt	tggagaagat	gacagctcct	gcctggagtg	gcagtggact	gaaaggccag	2460
cgaaagttag	atgattcacg	aaaagatgac	atgatggcca	ggagaactgg	gatgtccctt	2520
agacacactg	gatccaaccc	aaaccagttc	cttccagttc	catttgcaaa	gcaacaggat	2580
gtggaagaat	cttctaaagg	tctacccatg	aaagatcaga	gatacggctc	gagaactcct	2640
gtgtctgatg	acgcagagag	caccagcatg	tttgacatgc	ggtgtgagga	ggaggccgcg	2700
gtgcagccgc	acagcagggc	cgcagaggag	cagctgcagc	tgataaataa	ccagctgagg	2760
gaagaggacg	acaaatggca	agatgacctg	gctcgttgga	agagtcgtag	aagaagtgtt	2820
tctcaggact	taatcaagaa	agaggaagaa	aggaaaaaaa	tggagaagtt	actggctgga	2880
gaagatggga	caagtgaacg	aaggaaaagc	atcaaaacct	acagagaaat	tgttcaagaa	2940
aaagagcggg	gagagagaga	gctgcatgaa	gcatataaga	acgctcggtc	ccaggaggag	3000
gcagagggga	tccttcaaca	gtacattgag	aggttcacca	tcagtgaggc	tgttctcgaa	3060
cgcttggaga	tgccaaaaat	tctggaaaga	agccattcaa	cagagccaaa	tttatcctcc	3120
ttcctgaatg	accccaatcc	catgaaatac	ctgcggcaac	agtcactgcc	tccacccaaa	3180
ttcactgcca	ctggtgaaac	caccattgct	cgtgccagtg	ttctggatac	cagcatgtca	3240
gcaggcagtg	ggtctccaag	caaaactgtc	actcccaaag	cagtgcctat	gctgacaccc	3300
aagccttact	cccagcccaa	aaattctcaa	gatgttctga	agacctttaa	ggtagacggg	3360
aaagtcatgt	tgaatggaga	gacggttcat	agagaggagg	agaaggaaaag	agagtgtccc	3420
acggtggcac	ctgcccactc	cttaacccaa	tcccagatgt	ttgaagggtg	ggccagagtg	3480
cacgggtctc	cactggagct	gaaacaagac	aacggtagca	tcgagatcaa	cataaagaag	3540

ccaaactctg	ttccccaaga	gctcgcagca	accactgaga	aaacggaacc	gaatagtcaa	3600
gaggacaaga	atgatggtgg	aaaatcaaga	aaagggaata	tagaacttgc	ctcatcagaa	3660
ccacagcatt	ttacaacaac	tgtgactcga	tgcagcccga	ccgtggcctt	tgtggaatth	3720
ccctccagcc	cccagctgaa	gaatgatgtg	tcggaagaaa	aagaccagaa	gaaaccagaa	3780
aatgaaatga	gtggaaaggt	ggagttggtg	ctgtcacaaa	aggtggtaaa	gccaaaatct	3840
ccagaacccg	aagcaacgct	gacattttcca	tttctggaca	aatgcctga	agccaaccaa	3900
ctacatttgc	caaattctcaa	ttctcaagtg	gattctccaa	gcagtgagaa	gtcacctgth	3960
atgacacctt	ttaagttctg	ggcatgggac	ccagaagagg	agcgcaggcg	acaggaaaaa	4020
tggcaacagg	aacaggaacg	tttgctccag	gagagatacc	agaaggagca	ggacaagctg	4080
aaagaagagt	gggaaaaggc	ccaaaaggag	gtggaagagg	aagaacgcag	atactatgag	4140
gaggagcgta	agataattga	agacactgtg	gttccattta	ctgtttcttc	aagttccgct	4200
gaccagctgt	ctacctcttc	ctccatgact	gaaggcagtg	ggacaatgaa	taagatagac	4260
ctgggaaact	gtcaagatga	aaaacaagac	agaagatgga	agaaatcatt	ccaggggagat	4320
gacagtgact	tattgctgaa	gactagggga	agtgatcgac	tggaggagaa	gggcagccta	4380
actgaagggg	ccttggtctca	ttctgggaac	cctgtatcaa	aaggagtcca	tgaagacat	4440
cagctggata	ccgaggctgg	ggccccacac	tgtggaacaa	accacagct	tgctcaggat	4500
ccatcccaga	atcagcagac	atcaaatcca	acgcacagtt	cagaagatgt	gaagccaaaa	4560
accctcccg	tggataaaaag	cattaacat	cagatcgagt	ctcccagtga	aaggcggaag	4620
aaaagtcccc	gagagcactt	ccaggctggg	cctttctctc	cctgttctcc	cacccctccc	4680
ggtcagtcac	caaacaggtc	tataagtgga	aagaagctgt	gctcttctg	tgggcttctt	4740
ttgggtaaag	gagctgcaat	gatcatcgag	accctcaatc	tctattttca	catccagtgt	4800
ttcagggtgtg	gaatttgtaa	aggccagctt	ggagatgcag	tgagtgggac	ggatgttagg	4860
attcgaaatg	gtctcctgaa	ctgtaatgat	tgctacatgc	gatccagaag	tgccgggcag	4920
cctacaacat	tgtgacacgg	ctttcaagct	tccggatcac	tcaccatttc	tttactgaga	4980
gtgtcccctg	gcaactgctt	aacaaaatcc	caagctcagg	ggcttctcag	cattttaccta	5040
atttctgaaa	ggctcttctg	aaagggtgga	tctgttcttt	cgtagcacag	tgthttatgtt	5100
tttctgtth	attgtthtgg	tttttttttt	ttttttgcat	ttgcacagta	tacacaaaag	5160
aatatgggg	tgtaatgatc	ctgaatagct	caaaaaaggt	tttagcatgg	tcaaacaggc	5220
ttatggthta	aaatgtgthta	ttctcttctt	tgggaattag	ctaaatgatg	caataaacct	5280
gtthttgtth	agaatgtcta	ggaattaaac	actthtatgtt	tacagaattg	agctgcagaa	5340
agtgcaagac	atgccaatth	gagacacacg	gtcttctaag	actgaaggat	aaatthtaatg	5400
cattthcagaa	actaaacatc	acagcaagct	ctatctctga	gctataatth	gtththtaatg	5460

```

caaagacact agtttgataa tatatactgt aatcctgaaa catttggtgtt acttaccttt 5520
ggaggtagaa attataccaa taaattattg caccgttagt attagattct gtgtaccttg 5580
gaagttatgt cattaatata ggctgggttca tcaaataaag caaaccttg caatatcagc 5640
tagatttaca ctccgggacg ttgccccaaag gtaggaagaa agcagaggga aatatttcag 5700
tcatcatttc caaagtcatt atcaaaatct gtgaggaagt ttaatcttcc aaagagtcaa 5760
tgtcagacat caggcctctg ttgcctgctt ctctcgaggc actagattag gagtcttcaa 5820
taagagactt aacatgaggt atatggaaga tgaggcaccg agataagttc atcattaggt 5880
gtgagcactg ctccaccctg ctggcaagtt ctccctaagg gcctgaagca caggtgtcca 5940
aagaaaagcg ttaagtccat cttaatagaa tctatgtggt atatgatgtg gtcagccct 6000
ggtctgtgat cagcaagaac ctacagcaca gattatgcc tgcccacttc aatgaatacc 6060
tactctctc cattctccat cacttttttt gctatcaaga actccggacc ttgcccatgg 6120
agaagtttag agaggaactc ttgtggagag ctggtttatt ttctgccctg tgcgacgagt 6180
ttcagctggc caagaaagga gtcaagttat taaaaagcat cacaatgtag atctccaggc 6240
tggttttttg ttttttggtg ttaagactgg ggaaaggggg actatttatt ctgccttaaa 6300
tcaatggcaa ataagtcaag atgacatttt gtgaatgtag actatggata cactccta 6360
agattgatgt agtcataaaa ggggggtcaag tagatgtttt tctgttatgt aagcaataat 6420
ttttccgtgt cttattgagt atggctagcg attatttatt acatgctaga tgggttcttt 6480
gcatgtgggt tccatatagg tgcagaaatt tcctcagcca ctggagggat ttcgaccata 6540
tttgtcattt ggatgagctg ttattagatt gaaatctaca catcatttca ttaaaaattg 6600
tgccttagaa aacgcaaagc tgttgcacat ggcgataaat tatggatgca gtacattgaa 6660
gagagatgaa gtcacttcca agtttccaag acttctcatg gaggtgtttg ctgttttaca 6720
ggaaaaaata aaaataaaaa aag 6743

```

<210> 36

<211> 1467

<212> PRT

<213> Homo sapiens

<400> 36

```

Met Arg Lys Asp Thr Asp Asp Ile Glu Ser Pro Lys Arg Ser Ile Arg
1           5           10           15

```

```

Asp Ser Gly Tyr Ile Asp Cys Trp Asp Ser Glu Arg Ser Asp Ser Leu
20           25           30

```

```

Ser Pro Pro Arg His Gly Arg Asp Asp Ser Phe Asp Ser Leu Asp Ser
35           40           45

```

Phe Gly Ser Arg Ser Arg Gln Thr Pro Ser Pro Asp Val Val Leu Arg
 50 55 60

Gly Ser Ser Asp Gly Arg Gly Ser Asp Ser Glu Ser Asp Leu Pro His
 65 70 75 80

Arg Lys Leu Pro Asp Val Lys Lys Asp Asp Met Ser Ala Arg Arg Thr
 85 90 95

Ser His Gly Glu Pro Lys Ser Ala Val Pro Phe Asn Gln Tyr Leu Pro
 100 105 110

Asn Lys Ser Asn Gln Thr Ala Tyr Val Pro Ala Pro Leu Arg Lys Lys
 115 120 125

Lys Ala Glu Arg Glu Glu Tyr Arg Lys Ser Trp Ser Thr Ala Thr Ser
 130 135 140

Pro Leu Gly Gly Glu Arg Pro Phe Ser Phe Pro Glu Thr Ile Glu Glu
 145 150 155 160

Glu Gly Ser Glu Val Gly Ser Ala Gly Glu Asp Asn Pro Ala Gly Gln
 165 170 175

Met Asn Pro Gly Trp Lys Pro Ser Asp Gly Gly Cys Glu Leu Pro Asp
 180 185 190

Gly Ser Gly Lys Glu His Pro Ser Ser Asp Gly Ala Val Val Ala Pro
 195 200 205

Ala Pro Lys Ser Glu Glu Lys Asp Ala Ala Glu Ile Gln Lys Arg Lys
 210 215 220

Arg Leu Glu Gln Ala Gly Ile Lys Val Met Pro Ala Ala Gln Arg Phe
 225 230 235 240

Ala Ser Gln Lys Gln Leu Ser Glu Glu Lys Glu Ala Ile Arg Asp Ile
 245 250 255

Val Leu Arg Lys Glu Asn Ser Phe Leu Thr His Gln His Gly Asn Asp
 260 265 270

Ser Glu Ala Glu Gly Glu Val Val Cys Arg Leu Pro Asp Leu Glu Lys
 275 280 285

Asp Asp Phe Ala Ala Arg Arg Ala Arg Met Asn Gln Thr Lys Pro Met
 290 295 300

Val Pro Leu Asn Gln Leu Leu Tyr Gly Pro Tyr Pro Lys Lys Gly Ala
 305 310 315 320

Glu Lys Ser Asp Gly Ser Lys Gln Leu Ser Lys Gly Ile Ser Lys Lys
 325 330 335

Arg Ser Leu Glu Tyr Lys Arg Asn Gln Gly His Thr Glu Glu Val Lys
 340 345 350

Leu Ile Val Thr Cys Asn Met Arg Ala Gln Glu Ser Glu Pro Val Glu
 355 360 365

Gly Gly Leu Arg Lys Val Pro Asp Leu His Lys Asp Asp Leu Ala Gln
 370 375 380

Gln Arg Ile Gln Gly Ser Leu Ala Pro His Arg Glu Pro Pro Ser Phe
 385 390 395 400

Ile Thr Leu Ser Asn Ile Thr Glu Ala Asp Leu Glu Thr Trp Glu Arg
 405 410 415

Leu Lys Val Ser Glu Lys Ala Arg Asp Gly Asp Val Gln His Ile Cys
 420 425 430

Ala Ser Glu Pro Ser Pro Glu Ile Lys Ala Glu Thr Ala Ile Arg Asp
 435 440 445

Asp Phe Ala Asn Arg Lys Ala Arg Ala Ser Lys Lys Ala Ser Ser Pro
 450 455 460

Arg Gln Lys Phe Val His Phe Gly Pro Val Thr Glu Leu Asp Gln Gln
 465 470 475 480

Lys Trp Lys Arg Leu Ser Ile Gly Lys Ala Gly Pro Arg Glu Asp Glu
 485 490 495

Glu Glu Val Ile Cys His Gly Ser Lys Ile Gln Met Asp Ser Val Ser
 500 505 510

Pro Val Ser Ala Ala Thr Ser Ser Leu Lys Gly His Gln Ile Phe Asn
 515 520 525

Arg Gln Asn Asp Cys Arg Thr Met Asn Cys Gly Arg Gly Asp Tyr Cys
 530 535 540

Arg Arg Ala Ser Trp Leu Ala Pro Val Pro Glu Ser Gln Glu Glu Trp
 545 550 555 560

Val Cys Ser Leu Gly Glu Cys Pro Arg Gly Thr Glu Glu Val Thr Ser
 565 570 575

Lys Gln Leu Pro Gln Asp Gly Lys Glu Glu Thr Glu Ser Ala Pro Arg
 580 585 590

Asp Ser Glu Arg Leu Ser Lys Ala Glu Arg Ser Glu Asp Ser Ser Gln
 595 600 605

Pro Leu Val Cys Pro Leu Ala Ser Glu Cys Glu Ala Ser Gly Thr Glu
 610 615 620

Glu Lys Leu Glu Lys Met Thr Ala Pro Ala Trp Ser Gly Ser Gly Leu
 625 630 635 640

Lys Gly Gln Arg Lys Leu Asp Asp Ser Arg Lys Asp Asp Met Met Ala
 645 650 655

Arg Arg Thr Gly Met Ser Leu Arg His Thr Gly Ser Asn Pro Asn Gln
 660 665 670

Phe Leu Pro Val Pro Phe Ala Lys Gln Gln Asp Val Glu Glu Ser Ser
 675 680 685

Lys Gly Leu Pro Met Lys Asp Gln Arg Tyr Gly Pro Arg Thr Pro Val
 690 695 700

Ser Asp Asp Ala Glu Ser Thr Ser Met Phe Asp Met Arg Cys Glu Glu
 705 710 715 720

Glu Ala Ala Val Gln Pro His Ser Arg Ala Arg Gln Glu Gln Leu Gln
 725 730 735

Leu Ile Asn Asn Gln Leu Arg Glu Glu Asp Asp Lys Trp Gln Asp Asp
 740 745 750

Leu Ala Arg Trp Lys Ser Arg Arg Arg Ser Val Ser Gln Asp Leu Ile
 755 760 765

Lys Lys Glu Glu Glu Arg Lys Lys Met Glu Lys Leu Leu Ala Gly Glu
 770 775 780

Asp Gly Thr Ser Glu Arg Arg Lys Ser Ile Lys Thr Tyr Arg Glu Ile
 785 790 795 800

Val Gln Glu Lys Glu Arg Arg Glu Arg Glu Leu His Glu Ala Tyr Lys
 805 810 815

Asn Ala Arg Ser Gln Glu Glu Ala Glu Gly Ile Leu Gln Gln Tyr Ile
 820 825 830

Glu Arg Phe Thr Ile Ser Glu Ala Val Leu Glu Arg Leu Glu Met Pro
 835 840 845

Lys Ile Leu Glu Arg Ser His Ser Thr Glu Pro Asn Leu Ser Ser Phe
 850 855 860

Leu Asn Asp Pro Asn Pro Met Lys Tyr Leu Arg Gln Gln Ser Leu Pro
 865 870 875 880

Pro Pro Lys Phe Thr Ala Thr Val Glu Thr Thr Ile Ala Arg Ala Ser
 885 890 895

Val Leu Asp Thr Ser Met Ser Ala Gly Ser Gly Ser Pro Ser Lys Thr
 900 905 910

Val Thr Pro Lys Ala Val Pro Met Leu Thr Pro Lys Pro Tyr Ser Gln
 915 920 925

Pro Lys Asn Ser Gln Asp Val Leu Lys Thr Phe Lys Val Asp Gly Lys
 930 935 940

Val Ser Val Asn Gly Glu Thr Val His Arg Glu Glu Glu Lys Glu Arg
 945 950 955 960

Glu Cys Pro Thr Val Ala Pro Ala His Ser Leu Thr Lys Ser Gln Met
 965 970 975

Phe Glu Gly Val Ala Arg Val His Gly Ser Pro Leu Glu Leu Lys Gln
 980 985 990

Asp Asn Gly Ser Ile Glu Ile Asn Ile Lys Lys Pro Asn Ser Val Pro
 995 1000 1005

Gln Glu Leu Ala Ala Thr Thr Glu Lys Thr Glu Pro Asn Ser Gln
 1010 1015 1020

Glu Asp Lys Asn Asp Gly Gly Lys Ser Arg Lys Gly Asn Ile Glu
 1025 1030 1035

Leu Ala Ser Ser Glu Pro Gln His Phe Thr Thr Thr Val Thr Arg
 1040 1045 1050

Cys Ser Pro Thr Val Ala Phe Val Glu Phe Pro Ser Ser Pro Gln
 1055 1060 1065

Leu Lys	Asn Asp Val Ser	Glu	Glu Lys Asp Gln	Lys	Lys Pro Glu
1070		1075		1080	
Asn Glu	Met Ser Gly Lys Val	Glu Leu Val Leu Ser	Gln Lys Val		
1085		1090		1095	
Val Lys	Pro Lys Ser Pro	Glu	Pro Glu Ala Thr	Leu	Thr Phe Pro
1100		1105		1110	
Phe Leu	Asp Lys Met Pro	Glu	Ala Asn Gln Leu His	Leu Pro Asn	
1115		1120		1125	
Leu Asn	Ser Gln Val Asp	Ser	Pro Ser Ser Glu	Lys	Ser Pro Val
1130		1135		1140	
Met Thr	Pro Phe Lys Phe	Trp	Ala Trp Asp Pro	Glu	Glu Glu Arg
1145		1150		1155	
Arg Arg	Gln Glu Lys Trp	Gln	Gln Glu Gln Glu	Arg	Leu Leu Gln
1160		1165		1170	
Glu Arg	Tyr Gln Lys Glu	Gln	Asp Lys Leu Lys	Glu	Glu Trp Glu
1175		1180		1185	
Lys Ala	Gln Lys Glu Val	Glu	Glu Glu Glu Arg	Arg	Tyr Tyr Glu
1190		1195		1200	
Glu Glu	Arg Lys Ile Ile	Glu	Asp Thr Val Val	Pro	Phe Thr Val
1205		1210		1215	
Ser Ser	Ser Ser Ala Asp	Gln	Leu Ser Thr Ser	Ser	Ser Met Thr
1220		1225		1230	
Glu Gly	Ser Gly Thr Met	Asn	Lys Ile Asp Leu	Gly	Asn Cys Gln
1235		1240		1245	
Asp Glu	Lys Gln Asp Arg	Arg	Trp Lys Lys Ser	Phe	Gln Gly Asp
1250		1255		1260	
Asp Ser	Asp Leu Leu Leu	Lys	Thr Arg Glu Ser	Asp	Arg Leu Glu
1265		1270		1275	
Glu Lys	Gly Ser Leu Thr	Glu	Gly Ala Leu Ala	His	Ser Gly Asn
1280		1285		1290	
Pro Val	Ser Lys Gly Val	His	Glu Asp His Gln	Leu	Asp Thr Glu
1295		1300		1305	

Ala Gly Ala Pro His Cys Gly Thr Asn Pro Gln Leu Ala Gln Asp
 1310 1315 1320

Pro Ser Gln Asn Gln Gln Thr Ser Asn Pro Thr His Ser Ser Glu
 1325 1330 1335

Asp Val Lys Pro Lys Thr Leu Pro Leu Asp Lys Ser Ile Asn His
 1340 1345 1350

Gln Ile Glu Ser Pro Ser Glu Arg Arg Lys Lys Ser Pro Arg Glu
 1355 1360 1365

His Phe Gln Ala Gly Pro Phe Ser Pro Cys Ser Pro Thr Pro Pro
 1370 1375 1380

Gly Gln Ser Pro Asn Arg Ser Ile Ser Gly Lys Lys Leu Cys Ser
 1385 1390 1395

Ser Cys Gly Leu Pro Leu Gly Lys Gly Ala Ala Met Ile Ile Glu
 1400 1405 1410

Thr Leu Asn Leu Tyr Phe His Ile Gln Cys Phe Arg Cys Gly Ile
 1415 1420 1425

Cys Lys Gly Gln Leu Gly Asp Ala Val Ser Gly Thr Asp Val Arg
 1430 1435 1440

Ile Arg Asn Gly Leu Leu Asn Cys Asn Asp Cys Tyr Met Arg Ser
 1445 1450 1455

Arg Ser Ala Gly Gln Pro Thr Thr Leu
 1460 1465

<210> 37
 <211> 8456
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <222> (3848)..(3848)
 <223> n is a, c, g, or t

<220>
 <221> misc_feature
 <222> (5680)..(5680)
 <223> n is a, c, g, or t

<220>
 <221> misc_feature
 <222> (5735)..(5735)

<223> n is a, c, g, or t

<220>

<221> misc_feature

<222> (5750)..(5750)

<223> n is a, c, g, or t

<220>

<221> misc_feature

<222> (5858)..(5858)

<223> n is a, c, g, or t

<220>

<221> misc_feature

<222> (8099)..(8100)

<223> n is a, c, g, or t

<220>

<221> misc_feature

<222> (8108)..(8108)

<223> n is a, c, g, or t

<220>

<221> misc_feature

<222> (8130)..(8130)

<223> n is a, c, g, or t

<220>

<221> misc_feature

<222> (8448)..(8448)

<223> n is a, c, g, or t

<400> 37

ctctcgcggc gcgaccgctc agtcacttcg cccagagacc cggacctggg ccgctgggga	60
gcaggcggcc ataaaccccc tctctcccg gttccctgacg ccgcggcagg agctgttaca	120
aacaccctgc ggttggtctc cgatgccctt cagtgcgggtg gggacgcctg gaccctgggtg	180
agcgaacccc aagccacccc ccaccccaac tcagtgtctt cgccggcccc cggcccgtac	240
gcctgtctgg tcgccatggc tgaaaacaca gagggggatc tgaactccaa cctgctccac	300
gccccctacc acaccgggga ccctcagctg gacacggcca tcgggcagtg gctccgctgg	360
gataagaatc ccaaaacaaa agagcagatt gaaaacctgt tacggaatgg gatgaacaag	420
gagctgcgag atcgtctttg ttgccgaatg acttttgga ctgcaggact tcgttctgcc	480
atgggggcag ggttttgcta tattaatgac cttacagtaa tacagtcaac acaggggatg	540
tacaaatacc ttgagagatg tttctcagac ttcaagcaga gaggctttgt ggttgggtat	600
gacactcggg gtcaagtaac tagcagctgc agcagccaga ggcttgctaa actcactgct	660
gcagtcttgc tggccaaaga tgttcctgtg tacctttttt caagatatgt tctacacct	720
tttgtaccat atgcagttca gaagctcaaa gcagttgcag gtgtgatgat tactgcctct	780
cacaaccgca aggaagacaa tggatacaag gtttactggg aaactgggtg tcagatcaca	840
tctcctcatg ataaagaaat tctaaaatgt atagaagaat gtgtggaacc ctggaatggg	900

tccctggaatg ataatttagt ggataccagc ccgctgaaga gagaccctct gcaggacatt	960
tgcaggagat acatggaaga tctgaaaaag atctgttttt acagggagtt aaactcgaag	1020
accaccttga aatttgtgca cacatctttt catggggtcg gacatgacta tgtgcagttg	1080
gcttttaaaag tgtttggttt taagcctcca attccagtac cagaacaaaa agatcctgat	1140
ccagactttt ctaccgttaa atgtccaaat cctgaagaag gagaatctgt gctggaactt	1200
tccttgagac tggcagagaa agaaaatgcc cgggtagtgc tagccacaga tcctgatgca	1260
gacagactgg cagcagcaga acttcaggag aatggttggt ggaaagtttt cacagggaat	1320
gagttggcag ctttgttttg atggtggatg tttgattgct ggaagaaaaa taaatcaaga	1380
aatgctgatg tgaagaacgt ttatatgtta gccaccacag tctcttctaa aattctgaag	1440
gcaattgcac ttaaagaagg atttcatttt gaagaaacat taccaggttt taaatggatt	1500
ggaagtagga taatagacct cctggaaaat gggaaagaag tcctttttgc atttgaagag	1560
tctattgggt ttctctgtgg aacttcagtt ttggataaag atggggtgag tgcagctgtt	1620
gtggttgctg agatggcatc ttacctggaa accatgaata taacattgaa acagcaactg	1680
gttaagggtt atgaaaaata tggttatcat atttcaaaaa ctctctattt cttgtgttat	1740
gaaccaccta ccatcaaaag tatatttgaa aggtctcgta attttgattc tccaaaagaa	1800
tatccaaaat tttgtggaac atttgctata ttgcatgtac gggacgttac cactggatat	1860
gacagtagcc agcctaataa gaaatcagtg ctgcctgtga gtaaaaacag ccaaatgatt	1920
acatttactt ttcaaaatgg ctgtgttgct acccttcgga caagtggaac agaaccaaag	1980
ataaagtatt atgcagagat gtgtgcgtca cctgaccaga gtgacactgc tttactggag	2040
gaagaactga agaaactcat tgatgctctg atagagaatt ttcttcagcc tagtaagaat	2100
ggactgatct ggcgttctgt ttaggggtac accaatatgt catgacactg tgtgggcata	2160
tggaacagag caaccgagaa ctgcatacat tgaaccttgt gttagcattc tctctctatc	2220
tcctctggcc gagtatcttt ttctttttat tttctttctt tttggtcaac tggctagact	2280
aaacaaagta taagtttgaa atgaaatggt ctagaggga atgattcaaa ttttttcata	2340
attcaaacaa aagttattac actaaaagta ttatagtaac gtattgtcct tccgttaaca	2400
gaaacatcac aattcaaaag tgagttttct tattatagtg tgattaagcc tagttctgta	2460
tctcgtaatt gtgtatagca tggtaaaaga aaaaaaaaaa gacaggtaaa tgttatataa	2520
ttgtagggtt agttaaggaa attagtccat aaaattggca agaaaatgggt actttccttg	2580
attgttatga aaaagtgtca ttgtgacatc acaggattaa atttacaagt tagagtattt	2640
gaaataggat atttagcatt tctgtagttt agaacagatg ttacagtgtg tggttttggc	2700
atgtctgttt cagggtgccct tgggtgcagta taaatttact gtatgtattt aaaattctga	2760
tgtttactga acataaaaca atagtggcaa cactcctatt ggctttgttc ttatcaaatt	2820

tctcagtgcc ttctcagggg ctggtatgaa cagaaaaaac agtggttcctg catttaccat	2880
aatttaataca gtcaattttgc agatagctaa ggaaaatctt ccaaaatctg taggtgcttc	2940
ctctatatca agtgagttta ctgattttta aatcttagga atataaataa agatcttaat	3000
aaaagcaagc ttaatgtagt taaggatctt catattagaa ttgtgtcatt tatcaaacct	3060
tttattacag aaaaaaatct gaactgagaa aggagctctg aatttatctg acagtctaac	3120
ttttgaatta cttttttgta agtagattta tcttttattt tatgaagtaa tgttaatttt	3180
actaagaata aaacatccaa attttaagta ttatgaaata ctaaaactac atgttttaaa	3240
acagaggtcc agtttattct aacatcttgc tgacttctgt agaatttctg attgcatgac	3300
tattatttat cttcttttag tgtagaaaat aattttttac aaaaagtcac ttttgggctg	3360
ggcgcagtggt ctcacgctg taatcccagc actttgggag gctgaggcag gtggatcaca	3420
atgtcaggag tttgagacta gcctggccaa tatggtgaaa cctgtctct actaaaaata	3480
caaaaattag cggggtgtgg tgggtgggcac ctgtagtccc agctactcg gaggctgagg	3540
caggagaatt gcttgaacct aggaggctga gattgcagtg agccgaaatc acgccactac	3600
actccagcct gggcaacaga gggagactcc atctcaaaaa aaaaaaagt cacttttggc	3660
taatcatgca gtcttctgaa aaggtttaaat aaaaaaatgc tgtttaggct gggcttggtg	3720
gctcacatct gtaataccag cgctttggga ggctgaggca ggaggattgc ttgaggccag	3780
gagtttaagt tacagaacta tgatcacact gccgcactga agcctggatg acagagtga	3840
accttatntc aaaaaaaaaa aaaaaaagag agagagagag aaaaagaaat gctgtgttta	3900
atcttcttaa aatgtatctt gtcctacata gttctcagaa tacatatggg gtatagttag	3960
gagataacaa taacaagacc taaacaaatt gcttatggga gatagcactg taattagaga	4020
atttatttca tataaaataa taatataaag tttgaactga ttagaatgtt ccttaaatca	4080
tggtttogtt tgttgcagct gtcagctaac atttggttga gctcatataa tccaaacct	4140
attgtccaac tattgtattg ccactattgt taatttgttc ctctcttact ttcttattgt	4200
tggatcttat atgactgtat gactgtagat aatagacagc tagctaaatg tagagcacca	4260
tttttctcaa tgagtttggt ttcaaaggct ttgcactgga gaagatatgt gtagaatttt	4320
tgttttattt taaaatgtg taaaacatct tttgttttat atgcttttaa gaatgtagat	4380
tgtaatntag gaaatggcat atattctgca aagtgggtata atgttgtata tatgaaaaac	4440
aggtatattt ggtttagttc tgtgtttcag tgactgataa acaaaaatga agctaaaatg	4500
aatcaacggc tcattcatag ttactaaagc catgtcatga ctgtcattct attatgaaga	4560
aaacagtttt atttgagtgc tttaatataa tgcaacattt aagtcattct aagggtgaaa	4620
gttattgaag tgttcttctc tcaagttaac aaaattttac aaatgttcct gaagtgtatc	4680
gattgagggg attattttat tctagcctgt catgagtctg tctgagcatg atacttaaca	4740

tgaacttgat	gtagtaggtg	cacatttgcc	taacaggata	aattcagtga	tagcccaagc	4800
tggtcaggag	agacagaagt	gctggctagg	aggtctcagc	ctcggggcact	acaatgttta	4860
tctttttctt	tttgaacaat	attgtgaaaa	tgattgcatt	ctcattgagc	atgaaagtca	4920
aaactggaat	tttctaaagc	tttggagaaa	aggagagaat	tagtagggca	agatcaaagc	4980
atacattgga	aataaacagt	agaaataagc	aaacaatata	aagagacacg	ggaaatgctt	5040
ctcgtgaagg	gcaaattgta	caccataaca	gtagggaata	aactgtgtcc	cacctgttta	5100
cttgtgttta	tttgttggtt	tgatgagtat	gcacctacca	ttttccttta	ttgtgatcac	5160
taaaatgagg	ccaagggtta	ggaggtgaga	aaaagtatcc	cttttacgga	ttcagttcct	5220
gctaggatgt	tgtatcagaa	tcatgtgaga	actcaggaag	aggtagagca	gaaataaaaa	5280
ggaagaaaat	gtagattaga	aaataagact	tctgagaaaa	agttcaagga	ataaagattg	5340
tttgactctg	gagaggagaa	agctgttgag	atagtttgct	aagtttgcta	ataaagcttt	5400
attgagtctt	ttattgaagg	gttcaaagta	agaaatacta	tgccatagtg	ttctccatct	5460
ctagtgaaga	cagaaccaga	gggaataagg	atcttttttt	taaagacagc	atggaaagtt	5520
taagatttag	gaacttcttg	agaaatcttc	ataatatctt	ctattcaaaa	tctttttttt	5580
atttttaagg	gaataaaaaat	tttaaaaaatt	aagagatata	cagatgactt	tttaagggtcc	5640
ttctttccag	cacatttttt	ttaaaaactc	aggttagaan	aaaaatgtca	taaagggcac	5700
taaaagtaac	atttggggtt	cactccatat	ttagnagaat	ttagctcacn	aaacaatttt	5760
tgtgtgtttt	gagtaactga	tgcataagtg	caatgaataa	ttttcctgct	tcactgttat	5820
ttttaattta	acctggcaaa	ttaagtgaga	cttgaacnaa	attttgttca	tcttgaagaa	5880
tgtatagtgg	cttaggttat	ttgaggtttt	cgttttgaaat	atatatgttg	ttattttgat	5940
gtagatacgt	tagcagatgg	agaggttcta	agagattgga	tactttggaa	tgtctttata	6000
taggaaatag	acctcccat	tcttaagcaa	gttgcttctg	gaattgatgc	aacaggggat	6060
attagctgta	gtttaatgga	taagttgttc	tatgagctgc	tgtatcagt	acttttttcc	6120
acattttcca	ttgatatttg	tgatgcttga	aaaatccaaa	cttcacttgg	atacatttaa	6180
ttttatgcct	ttttctctct	acctcccact	tttcctaggc	acccttgcac	attttgttac	6240
tgtgaaggac	tattgccctg	ttacctcact	gataagaaat	ctcaactgta	ttcattcaga	6300
atgctttaag	cttagtcata	aactacattt	gcctatggac	ctttgggttat	tttgttaatg	6360
ttgtcattat	atttaagact	tattgtagga	agtaatttta	aatctatttc	ataacctggt	6420
ttgatctgtc	tgcatactgc	tatatgttaa	tttattacgg	cttttacaat	tcacttgtca	6480
tctgagattg	ttctagaaac	tgatgaggaa	agagtcgtac	taacccaaaat	gaatgggcaa	6540
ttgacataac	attctatttc	aggtttcctt	gtacttgggg	agccaaagca	caagatctaa	6600
ggaaatctta	ttcacatttg	ttcctcagag	catcttacct	taagccacag	aggtataaat	6660

```

agtaagatga gaattgctca ttgtctgact ccttgcacag catgaattta attctagttg 6720
gaaacaaagt tagtggtgac tactgggttac cttaaagtat tcaccatact atatatattat 6780
tagaccatct ttctcttttg tttattcaag actatgtact tataaaagag aacctataag 6840
aatgtgtaga gtgaccaaata agggacaaaag tgacatacct agagtttgct gctcaaattt 6900
ttccactagc gtaaccctaa agcagagggg catgatgaag ggggaagggg ttctgagagc 6960
aaaatcagaa tcagcctacc actgacatga agtgtgttct gtcttttaaa tgtaagtga 7020
ttttgcattc tctataataa aacttttaata ataaaaatga tatgtttaat ctttactgaa 7080
acttatgcca ccacaaagga gcaccatgtt catcacgtgg ctgctcagtt ttttccccc 7140
gtgccctcac cgccagtggg ggcaagtcct tgagctacac aactggctg tagtgcaaga 7200
tatataaaca taggccgggc tcggtggctc atgcctgtaa tcctagcact ttgggaggcc 7260
gaagcaggtg gatcatctga ggtcgggaga ttgagactag cctgaccgac atggagaaac 7320
cctgtgtcca ctaaaaatac aaaattagcc ggggtgtggtg gcacatgcct gtaatcccag 7380
ctactcggga ggctgaggca ggagaatcac ttgaaccag gaggtggagg ttgtggtgag 7440
ccaagatcgc gccgttgac tccagcctgg gcaacaagag tgaaactccg tctcaaaaaa 7500
aaaaaaaaa gatataataa catagaacat atccagacta tttcctgccc ttggaatctg 7560
accttcagaa ggattaacgc caaactaaac tgacctgaaa cagttttgtt actattggca 7620
ttccttgact aaaaaggacc acctgatgct cagtcccag agtaggaatc gtgggtttatt 7680
atttttcata tccataccac ctagttcagt acctgttata gagtgtttaa ttgctgatac 7740
taggaattaa gggatttttt aaaagtagca aatagtttaa ggaacaatca aaaaatgaat 7800
tttctaattc agttttttca atgaagtccc atcactgaac ttataatgt aatgtttatt 7860
aagcagacat ccccttaaata actcagtttc tttggagaga gcaaagcata tctccctgtg 7920
gaaattatat ataatgcctg gtattttaca ctcaaagcaa ggtgttggcc aggaaggtaa 7980
agaaatggtg tctatgtaga cttaatatca ctagtctaag tgtcattttg agcctagtag 8040
cactaccttc caagtgagtc acaacaaatt tgatcctatt ttaggtttt ttccactgnn 8100
atgattcntc atgtatctta caagagccan tcaagcaaga ctctgcttct atgtatggtg 8160
aggccttggt gttctagggt agaataaact ctttgtatgc ctcatgaat atgccaggta 8220
aaatatatgc agtcaagaat gaattatttt tctgactaaa gtgtgtagca gtagttcaaa 8280
attgtgccct tgttttaaca gtttctgtca acatcttctc atttttccct acaaaaacac 8340
cagggtgtat tataagtact gcctgtgaga atttgcactt tatgtatttg tgtgtggatt 8400
tcttgtggtt ttagccaaat gaagtgttat cagtaataaa caggtctntt catagg 8456

```

<210> 38

<211> 622

<212> PRT

<213> Homo sapiens

<400> 38

Met Ala Glu Asn Thr Glu Gly Asp Leu Asn Ser Asn Leu Leu His Ala
 1 5 10 15

Pro Tyr His Thr Gly Asp Pro Gln Leu Asp Thr Ala Ile Gly Gln Trp
 20 25 30

Leu Arg Trp Asp Lys Asn Pro Lys Thr Lys Glu Gln Ile Glu Asn Leu
 35 40 45

Leu Arg Asn Gly Met Asn Lys Glu Leu Arg Asp Arg Leu Cys Cys Arg
 50 55 60

Met Thr Phe Gly Thr Ala Gly Leu Arg Ser Ala Met Gly Ala Gly Phe
 65 70 75 80

Cys Tyr Ile Asn Asp Leu Thr Val Ile Gln Ser Thr Gln Gly Met Tyr
 85 90 95

Lys Tyr Leu Glu Arg Cys Phe Ser Asp Phe Lys Gln Arg Gly Phe Val
 100 105 110

Val Gly Tyr Asp Thr Arg Gly Gln Val Thr Ser Ser Cys Ser Ser Gln
 115 120 125

Arg Leu Ala Lys Leu Thr Ala Ala Val Leu Leu Ala Lys Asp Val Pro
 130 135 140

Val Tyr Leu Phe Ser Arg Tyr Val Pro Thr Pro Phe Val Pro Tyr Ala
 145 150 155 160

Val Gln Lys Leu Lys Ala Val Ala Gly Val Met Ile Thr Ala Ser His
 165 170 175

Asn Arg Lys Glu Asp Asn Gly Tyr Lys Val Tyr Trp Glu Thr Gly Ala
 180 185 190

Gln Ile Thr Ser Pro His Asp Lys Glu Ile Leu Lys Cys Ile Glu Glu
 195 200 205

Cys Val Glu Pro Trp Asn Gly Ser Trp Asn Asp Asn Leu Val Asp Thr
 210 215 220

Ser Pro Leu Lys Arg Asp Pro Leu Gln Asp Ile Cys Arg Arg Tyr Met
 225 230 235 240

Glu Asp Leu Lys Lys Ile Cys Phe Tyr Arg Glu Leu Asn Ser Lys Thr
 245 250 255
 Thr Leu Lys Phe Val His Thr Ser Phe His Gly Val Gly His Asp Tyr
 260 265 270
 Val Gln Leu Ala Phe Lys Val Phe Gly Phe Lys Pro Pro Ile Pro Val
 275 280 285
 Pro Glu Gln Lys Asp Pro Asp Pro Asp Phe Ser Thr Val Lys Cys Pro
 290 295 300
 Asn Pro Glu Glu Gly Glu Ser Val Leu Glu Leu Ser Leu Arg Leu Ala
 305 310 315 320
 Glu Lys Glu Asn Ala Arg Val Val Leu Ala Thr Asp Pro Asp Ala Asp
 325 330 335
 Arg Leu Ala Ala Ala Glu Leu Gln Glu Asn Gly Cys Trp Lys Val Phe
 340 345 350
 Thr Gly Asn Glu Leu Ala Ala Leu Phe Gly Trp Trp Met Phe Asp Cys
 355 360 365
 Trp Lys Lys Asn Lys Ser Arg Asn Ala Asp Val Lys Asn Val Tyr Met
 370 375 380
 Leu Ala Thr Thr Val Ser Ser Lys Ile Leu Lys Ala Ile Ala Leu Lys
 385 390 395 400
 Glu Gly Phe His Phe Glu Glu Thr Leu Pro Gly Phe Lys Trp Ile Gly
 405 410 415
 Ser Arg Ile Ile Asp Leu Leu Glu Asn Gly Lys Glu Val Leu Phe Ala
 420 425 430
 Phe Glu Glu Ser Ile Gly Phe Leu Cys Gly Thr Ser Val Leu Asp Lys
 435 440 445
 Asp Gly Val Ser Ala Ala Val Val Val Ala Glu Met Ala Ser Tyr Leu
 450 455 460
 Glu Thr Met Asn Ile Thr Leu Lys Gln Gln Leu Val Lys Val Tyr Glu
 465 470 475 480
 Lys Tyr Gly Tyr His Ile Ser Lys Thr Ser Tyr Phe Leu Cys Tyr Glu
 485 490 495

Pro Pro Thr Ile Lys Ser Ile Phe Glu Arg Leu Arg Asn Phe Asp Ser
500 505 510

Pro Lys Glu Tyr Pro Lys Phe Cys Gly Thr Phe Ala Ile Leu His Val
515 520 525

Arg Asp Val Thr Thr Gly Tyr Asp Ser Ser Gln Pro Asn Lys Lys Ser
530 535 540

Val Leu Pro Val Ser Lys Asn Ser Gln Met Ile Thr Phe Thr Phe Gln
545 550 555 560

Asn Gly Cys Val Ala Thr Leu Arg Thr Ser Gly Thr Glu Pro Lys Ile
565 570 575

Lys Tyr Tyr Ala Glu Met Cys Ala Ser Pro Asp Gln Ser Asp Thr Ala
580 585 590

Leu Leu Glu Glu Glu Leu Lys Lys Leu Ile Asp Ala Leu Ile Glu Asn
595 600 605

Phe Leu Gln Pro Ser Lys Asn Gly Leu Ile Trp Arg Ser Val
610 615 620

<210> 39

<211> 2015

<212> DNA

<213> Homo sapiens

<400> 39

acaaataata aggaacaagg tgtggttatt gattctctaa aattaagtga ggagatgaag	60
cccaatctag atggtgttga tttattcaac aatggtggtt ctggaaacgg tgaaacgaaa	120
actgggctga gactgaaagc aataaatctg cctttggaaa atgaagtaac tgaaatttca	180
gctttgcagg tgcatttgga tgaattccaa aaaatcttat ggaaggaaag agaaatgcgc	240
acagctttgg aaaaagaaat agagagactg gagtcggctt tgtctctgtg gaagtggaa	300
tatgaagaac tgaaagaatc aaagccaaaa aatgtgaaag agtttgacat tcttcttggt	360
caacataatg atgaaatgca agaactgtca ggcaatataa aggaagaatc caaatctcaa	420
aacagcaaag acagagtgat ttgtgagtta agagcagagc tagagagatt gcaagctgaa	480
aatacctcgg agtgggacaa gagggaaata cttgaaagag aaaagcaggg actggagaga	540
gaaaatagaa ggctgaagat ccaggtgaaa gaaatggaag agcttttgga taagaaaaat	600
agattaagtg caaactctca aagtcctgat ttcaagatgt cacaaattga tctgcaagaa	660
aaaaaccagg aattactgaa ccttcaacat gcctactata aactaaacag acaataccag	720


```

gcaaatattg cagaactgac tcatgcaaac aaccgagtgg atcaaaatga agcagaagta 780
aagaaactaa gattacgagt ggaagaacta aagcagggac tcaaccaaaa agaagatgag 840
cttgatgatt cectgaatca gatccgtaag ctccagaggt ctctggatga agagaaagaa 900
agaaatgaaa acttagagac tgaactcagg cacttgcaaa actggtaatt ttttcacaaa 960
atatgctgaa ttaaagatta gggccttaaa gacatttcca tacccttttc ttaaatatca 1020
gtaaaattgt ttttattaac tagaaatatt aatgaaaaaa acgtagacaa tacacaaatt 1080
aatgggcttc ttcacttctt ctaatttttg cctaacagat actgcatatt ctcaaaaaga 1140
caattttaat gtcattttaa aacaacttta attctaagat gtgtaaatat tttgaaagtc 1200
aaaaagggct ttcagaatac tttttacata aaatctgaaa gagttataat atcggttaaga 1260
aaaagtaagt tgaaaaccat acaagacgct gggtcattaa taagaaaacc attgacttta 1320
agtataaagt actggtttgt ttaaataatt ggtaaacttt tatgtacgtg ttgtctatgt 1380
ggtaggggatg gcaggttgta ttaacaaaaa tgaatcattc tagaggtgta acaatacatt 1440
tcttatataa ttttataagt cattttctaatt ctttgtataa aacagaagtg agcaagatga 1500
atcagaaaaa aggtgttttg tatttttaaag gtaacagata accaggtgat tgaatctaag 1560
acaggctgta agcatcgctg agaaactaaa aggacttttg acttttatct ggatagacat 1620
ttctacagta aaatcatgga aaggcatcag cattgcaaag tagcatctag gtagaaatca 1680
ggccaaaatt aagctgtggt ttccctctga gtagtgggaa tagagaaaat taggaaattg 1740
tggttatgtg aatatttctt taaaactttt atgtacatta tagtttattg cttcatattt 1800
aagtttagtt ttttaaggtaa aatgttattt tgaacaaaaa gacattata attttccata 1860
cctattttca actgaaggca acttgtaaga ttttaactcag tcaataacat actggtttta 1920
ctcatctccc cctccattga ttagccaaaa aaaaatgaaa tcttactaat tcattattga 1980
ataaagacca cttttatcag agtcaaaaaa aaaaaa 2015

```

<210> 40

<211> 297

<212> PRT

<213> Homo sapiens

<400> 40

```

Met Lys Pro Asn Leu Asp Gly Val Asp Leu Phe Asn Asn Gly Gly Ser
1           5           10           15

```

```

Gly Asn Gly Glu Thr Lys Thr Gly Leu Arg Leu Lys Ala Ile Asn Leu
20           25           30

```

```

Pro Leu Glu Asn Glu Val Thr Glu Ile Ser Ala Leu Gln Val His Leu
35           40           45

```

Asp Glu Phe Gln Lys Ile Leu Trp Lys Glu Arg Glu Met Arg Thr Ala
 50 55 60

Leu Glu Lys Glu Ile Glu Arg Leu Glu Ser Ala Leu Ser Leu Trp Lys
 65 70 75 80

Trp Lys Tyr Glu Glu Leu Lys Glu Ser Lys Pro Lys Asn Val Lys Glu
 85 90 95

Phe Asp Ile Leu Leu Gly Gln His Asn Asp Glu Met Gln Glu Leu Ser
 100 105 110

Gly Asn Ile Lys Glu Glu Ser Lys Ser Gln Asn Ser Lys Asp Arg Val
 115 120 125

Ile Cys Glu Leu Arg Ala Glu Leu Glu Arg Leu Gln Ala Glu Asn Thr
 130 135 140

Ser Glu Trp Asp Lys Arg Glu Ile Leu Glu Arg Glu Lys Gln Gly Leu
 145 150 155 160

Glu Arg Glu Asn Arg Arg Leu Lys Ile Gln Val Lys Glu Met Glu Glu
 165 170 175

Leu Leu Asp Lys Lys Asn Arg Leu Ser Ala Asn Ser Gln Ser Pro Asp
 180 185 190

Phe Lys Met Ser Gln Ile Asp Leu Gln Glu Lys Asn Gln Glu Leu Leu
 195 200 205

Asn Leu Gln His Ala Tyr Tyr Lys Leu Asn Arg Gln Tyr Gln Ala Asn
 210 215 220

Ile Ala Glu Leu Thr His Ala Asn Asn Arg Val Asp Gln Asn Glu Ala
 225 230 235 240

Glu Val Lys Lys Leu Arg Leu Arg Val Glu Glu Leu Lys Gln Gly Leu
 245 250 255

Asn Gln Lys Glu Asp Glu Leu Asp Asp Ser Leu Asn Gln Ile Arg Lys
 260 265 270

Leu Gln Arg Ser Leu Asp Glu Glu Lys Glu Arg Asn Glu Asn Leu Glu
 275 280 285

Thr Glu Leu Arg His Leu Gln Asn Trp
 290 295

<210> 41
 <211> 2372
 <212> DNA
 <213> Hmo sapiens

<400> 41
 cgataatgac tttgcacgat tcaactttggg atctcaaagt gcttccaaag cattcagatt 60
 taaaaacaat tcacaagaca ggatcatcttt gtaataccca tacttacaac gaattaacaa 120
 gaggagtgc ttaagattct ccaggaacac agtggcagct attgatgac tgttttctat 180
 ctgtttgata gagcatcatg agaaatcaca aaatacaatg ctatttttct gatgtgtgct 240
 aataaagtca aagaaagcaa atacatcttg acacttttgt ccattttcat taaaaaaaaa 300
 aaaaagttca ggggtgtttg aattttacac ctcagcacac cttactggta tcaatggata 360
 aagcgggtaa ttgacagatc caccctaatg ccactgcagt cagaagcaga tctggacaca 420
 cccttgttta cagtttcata ttgggttgct atagttcccg tgctaaatca ccagctttca 480
 ggaacatgac tgctcctggc agtgggaagg gctgaaacag aaattttaat taaaaacttt 540
 atcaagtact cttcacagtg ctgcttgga ccatagaaaa tcagtacaat atatcgagcc 600
 ctactttgga ggagctggat ttctgagga gctgatccag ttctaagtgt cttctcgaat 660
 taggagatag atgatctttg atggggatct cctcgtcac cacaggccag tcacagaacc 720
 aactagccac gtgctgccag acctcagtgg gcccaagcag gagcaatctc ttctatcccc 780
 catctcccc aggaccatcc ccccatgt caatgtcatc cagggtctt ctggtagtga 840
 gtgacttttc tgcacatgtt tagggcttg gggagctaga acacaggaaa catgaatgca 900
 aaaggcatgg aaaacactgt tttgctttg gttagtataa tgtgggcagg aaaaagatta 960
 ctattggctg gagctttgcc aagtgcagata gaatcaactg tcacccatt cctttcccag 1020
 aaggctctat ggtattaagg atacatccag tattttccca cagattttta ttcaggcgat 1080
 gtttcataaa ttacatatat gaaaacattc attattacat ttccttgtgt gtttcaaca 1140
 gacattggca ccttcctatt gagttaattc tctgcattc ttgcagcagc agcccgcaag 1200
 gagattccca gagatggctc ccctaacaca cagtcctgtg attttacagt tctatgactt 1260
 acagttgatg attcacaaga ttcaggattc tacaagactc aagggggaac taaactttct. 1320
 tacgattgta catgatcagt tatagggtg taatcattaa ttgttggtt caaatgtgga 1380
 cacacacaca cacacatcat gcccaaggagg gaatgggggtg tttcaagtca ggcagcgatg 1440
 attctggaag gttggaaatg taagggtaga agcttggtg gtcttagtaa acttggtccc 1500
 ttgctccac caagaagagg taccaaatgt gagacctgag atctcctcca atatctgtcc 1560
 tctgcagttc cgggaaacta atcatgaagt acacatgcag cagctcctcc acttcctttc 1620
 ctccgaggtc ctcttttcca ttctcccacc tagatactga cacaccgcca cggtttccac 1680
 attggaaggg cagaacactg tgcagtatcg tgcacacttg ctgggttagg aatagagctg 1740

```

ccctagggtc accttcatgc aagtattgac agctacaaat taaagtcctt agagcagttg 1800
acacagatac tacgttctag aagagaatta aatttaaacg tcaagtttaa agggatcata 1860
attctgcagg tatctttctc tgagtgactg aatgtgacta ttgcattagg gtaaataaat 1920
taagacgtgc aagtgggatt tactgtatgt tagaaaggag ttttgcagcc aagactgcct 1980
tgaataaaat gtgtttgcac tgaaaaaaa ttttaaatta cttggtctct ggttgctgta 2040
aagggtcatcc aagatggatg ttctgtttat attgtatagt atttcatatg aaataattac 2100
agttcatgaa atgtcttccc taatgttact gatttataac agcacatttg taacatgggt 2160
tttatcgtgt cagtgtacca tactgtaaat gatgattact tgtcatgctt agtataataa 2220
cttaaaagaa aaaaaaggac agggatTTTT gtaagtctat atttgaaagt ccctccctat 2280
ggtgatactg tgttcatgtt gtttatgtag tgttgtgtga aatatccatt ttggattgtg 2340
ttacttttta agatattaaa taacatttgg tt 2372

```

```

<210> 42
<211> 365
<212> PRT
<213> Homo sapiens

```

```

<400> 42

```

```

Lys Arg Lys Val Ala Gln Glu Ala Ile Glu Ser Leu Ser Ala Ser Lys
1          5          10          15

```

```

Leu Ala Lys Ser Ile Cys Ser Gln Phe Arg Thr Arg Leu Asn Ser Ser
          20          25          30

```

```

His Glu Ala Phe Ala Ala Ser Leu Arg Gln Leu Glu Ala Gly His Ser
          35          40          45

```

```

Gly Arg Leu Glu Lys Thr Glu Asp Leu Trp Leu Arg Val Arg Lys Asp
          50          55          60

```

```

His Ala Pro Arg Leu Ala Arg Leu Ser Leu Glu Ser Arg Ser Leu Gln
65          70          75          80

```

```

Asp Val Leu Leu His Arg Lys Pro Lys Leu Gly Gln Glu Leu Gly Arg
          85          90          95

```

```

Gly Gln Tyr Gly Val Val Tyr Leu Cys Asp Asn Trp Gly Gly His Phe
          100          105          110

```

```

Pro Cys Ala Leu Lys Ser Val Val Pro Pro Asp Glu Lys His Trp Asn
          115          120          125

```

```

Asp Leu Ala Leu Glu Phe His Tyr Met Arg Ser Leu Pro Lys His Glu

```

130		135		140
Arg Leu Val Asp Leu His Gly Ser Val Ile Asp Tyr Asn Tyr Gly Gly				
145		150		155 160
Gly Ser Ser Ile Ala Val Leu Leu Ile Met Glu Arg Leu His Arg Asp				
	165		170	175
Leu Tyr Thr Gly Leu Lys Ala Gly Leu Thr Leu Glu Thr Arg Leu Gln				
	180		185	190
Ile Ala Leu Asp Val Val Glu Gly Ile Arg Phe Leu His Ser Gln Gly				
	195		200	205
Leu Val His Arg Asp Ile Lys Leu Lys Asn Val Leu Leu Asp Lys Gln				
	210		215	220
Asn Arg Ala Lys Ile Thr Asp Leu Gly Phe Cys Lys Pro Glu Ala Met				
	225		230	235 240
Met Ser Gly Ser Ile Val Gly Thr Pro Ile His Met Ala Pro Glu Leu				
	245		250	255
Phe Thr Gly Lys Tyr Asp Asn Ser Val Asp Val Tyr Ala Phe Gly Ile				
	260		265	270
Leu Phe Trp Tyr Ile Cys Ser Gly Ser Val Lys Leu Pro Glu Ala Phe				
	275		280	285
Glu Arg Cys Ala Ser Lys Asp His Leu Trp Asn Asn Val Arg Arg Gly				
	290		295	300
Ala Arg Pro Glu Arg Leu Pro Val Phe Asp Glu Glu Cys Trp Gln Leu				
	305		310	315 320
Met Glu Ala Cys Trp Asp Gly Asp Pro Leu Lys Arg Pro Leu Leu Gly				
	325		330	335
Ile Val Gln Pro Met Leu Gln Gly Ile Met Asn Arg Leu Cys Lys Ser				
	340		345	350
Asn Ser Glu Gln Pro Asn Arg Gly Leu Asp Asp Ser Thr				
	355		360	365

<210> 43
 <211> 1185
 <212> PRT
 <213> Homo sapiens

<400> 43

Lys Lys Ile Arg Glu Lys Phe Asn Arg Tyr Leu Asp Val Val Asn Arg
 1 5 10 15

Asn Lys Gln Val Val Glu Ala Ser Tyr Thr Ala His Leu Thr Ser Pro
 20 25 30

Leu Thr Ala Ile Gln Asp Cys Cys Thr Ile Pro Pro Ser Met Met Glu
 35 40 45

Phe Asp Gly Asn Phe Asn Thr Asn Val Ser Arg Thr Ile Ser Cys Asp
 50 55 60

Arg Leu Ser Thr Thr Val Asn Ser Arg Ala Phe Asn Pro Gly Arg Asp
 65 70 75 80

Leu Asn Ser Val Leu Ala Asp Asn Leu Lys Ser Asn Pro Gly Ile Lys
 85 90 95

Trp Gln Tyr Phe Ser Ser Glu Glu Gly Ile Phe Thr Val Phe Pro Ala
 100 105 110

His Lys Phe Arg Cys Lys Gly Ser Tyr Glu His Arg Ser Arg Pro Ile
 115 120 125

Tyr Val Ser Thr Val Arg Pro Gln Ser Lys His Ile Val Val Ile Leu
 130 135 140

Asp His Gly Ala Ser Val Thr Asp Thr Gln Leu Gln Ile Ala Lys Asp
 145 150 155 160

Ala Ala Gln Val Ile Leu Ser Ala Ile Asp Glu His Asp Lys Ile Ser
 165 170 175

Val Leu Thr Val Ala Asp Thr Val Arg Thr Cys Ser Leu Asp Gln Cys
 180 185 190

Tyr Lys Thr Phe Leu Ser Pro Ala Thr Ser Glu Thr Lys Arg Lys Met
 195 200 205

Ser Thr Phe Val Ser Ser Val Lys Ser Ser Asp Ser Pro Thr Gln His
 210 215 220

Ala Val Gly Phe Gln Lys Ala Phe Gln Leu Ile Arg Ser Thr Asn Asn
 225 230 235 240

Asn Thr Lys Phe Gln Ala Asn Thr Asp Met Val Ile Ile Tyr Leu Ser

	245		250		255
Ala Gly Ile Thr Ser Lys Asp Ser Ser Glu Glu Asp Lys Lys Ala Thr	260		265		270
Leu Gln Val Ile Asn Glu Glu Asn Ser Phe Leu Asn Asn Ser Val Met	275		280		285
Ile Leu Thr Tyr Ala Leu Met Asn Asp Gly Val Thr Gly Leu Lys Glu	290		295		300
Leu Ala Phe Leu Arg Asp Leu Ala Glu Gln Asn Ser Gly Lys Tyr Gly	305		310		315
Val Pro Asp Arg Thr Ala Leu Pro Val Ile Lys Gly Ser Met Met Val	325		330		335
Leu Asn Gln Leu Ser Asn Leu Glu Thr Thr Val Gly Arg Phe Tyr Thr	340		345		350
Asn Leu Pro Asn Arg Met Ile Asp Glu Ala Val Phe Ser Leu Pro Phe	355		360		365
Ser Asp Glu Met Gly Asp Gly Leu Ile Met Thr Val Ser Lys Pro Cys	370		375		380
Tyr Phe Gly Asn Leu Leu Leu Gly Ile Val Gly Val Asp Val Asn Leu	385		390		395
Ala Tyr Ile Leu Glu Asp Val Thr Tyr Tyr Gln Asp Ser Leu Ala Ser	405		410		415
Tyr Thr Phe Leu Ile Asp Asp Lys Gly Tyr Thr Leu Met His Pro Ser	420		425		430
Leu Thr Arg Pro Tyr Leu Leu Ser Glu Pro Pro Leu His Thr Asp Ile	435		440		445
Ile His Tyr Glu Asn Ile Pro Lys Phe Glu Leu Val Arg Gln Asn Ile	450		455		460
Leu Ser Leu Pro Leu Gly Ser Gln Ile Ile Ala Val Pro Val Asn Ser	465		470		475
Ser Leu Ser Trp His Ile Asn Lys Leu Arg Glu Thr Gly Lys Glu Ala	485		490		495

Tyr Asn Val Ser Tyr Ala Trp Lys Met Val Gln Asp Thr Ser Phe Ile
 500 505 510

Leu Cys Ile Val Val Ile Gln Pro Glu Ile Pro Val Lys Gln Leu Lys
 515 520 525

Asn Leu Asn Thr Val Pro Ser Ser Lys Leu Leu Tyr His Arg Leu Asp
 530 535 540

Leu Leu Gly Gln Pro Ser Ala Cys Leu His Phe Lys Gln Leu Ala Thr
 545 550 555 560

Leu Glu Ser Pro Thr Ile Met Leu Ser Ala Gly Ser Phe Ser Ser Pro
 565 570 575

Tyr Glu His Leu Ser Gln Pro Glu Thr Lys Arg Met Val Glu His Tyr
 580 585 590

Thr Ala Tyr Leu Ser Asp Asn Thr Arg Leu Ile Ala Asn Pro Gly Leu
 595 600 605

Lys Phe Ser Val Arg Asn Glu Val Met Ala Thr Ser His Val Thr Asp
 610 615 620

Glu Trp Met Thr Gln Met Glu Met Ser Ser Leu Asn Thr Tyr Ile Val
 625 630 635 640

Arg Arg Tyr Ile Ala Thr Pro Asn Gly Val Leu Arg Ile Tyr Pro Gly
 645 650 655

Ser Leu Met Asp Lys Ala Phe Asp Pro Thr Arg Arg Gln Trp Tyr Leu
 660 665 670

His Ala Val Ala Asn Pro Gly Leu Ile Ser Leu Thr Gly Pro Tyr Leu
 675 680 685

Asp Val Gly Gly Ala Gly Tyr Val Val Thr Ile Ser His Thr Ile His
 690 695 700

Ser Ser Ser Thr Gln Leu Ser Ser Gly His Thr Val Ala Val Met Gly
 705 710 715 720

Ile Asp Phe Thr Leu Arg Tyr Phe Tyr Lys Val Leu Met Asp Leu Leu
 725 730 735

Pro Val Cys Asn Gln Asp Gly Gly Asn Lys Ile Arg Cys Phe Ile Met
 740 745 750

Glu Asp Arg Gly Tyr Leu Val Ala His Pro Thr Leu Ile Asp Pro Lys
 755 760 765

Gly His Ala Pro Val Glu Gln Gln His Ile Thr His Lys Glu Pro Leu
 770 775 780

Val Ala Asn Asp Ile Leu Asn His Pro Asn Phe Val Lys Lys Asn Leu
 785 790 795 800

Cys Asn Ser Phe Ser Asp Arg Thr Val Gln Arg Phe Tyr Lys Phe Asn
 805 810 815

Thr Ser Leu Ala Gly Asp Leu Thr Asn Leu Val His Gly Ser His Cys
 820 825 830

Ser Lys Tyr Arg Leu Ala Arg Ile Pro Gly Thr Asn Ala Phe Val Gly
 835 840 845

Ile Val Asn Glu Thr Cys Asp Ser Leu Ala Phe Cys Ala Cys Ser Met
 850 855 860

Val Asp Arg Leu Cys Leu Asn Cys His Arg Met Glu Gln Asn Glu Cys
 865 870 875 880

Glu Cys Pro Cys Glu Cys Pro Leu Glu Val Asn Glu Cys Thr Gly Asn
 885 890 895

Leu Thr Asn Ala Glu Asn Arg Asn Pro Ser Cys Glu Val His Gln Glu
 900 905 910

Pro Val Thr Tyr Thr Ala Ile Asp Pro Gly Leu Gln Asp Ala Leu His
 915 920 925

Gln Cys Val Asn Ser Arg Cys Ser Gln Arg Leu Glu Ser Gly Asp Cys
 930 935 940

Phe Gly Val Leu Asp Cys Glu Trp Cys Met Val Asp Ser Asp Gly Lys
 945 950 955 960

Thr His Leu Asp Lys Pro Tyr Cys Ala Pro Gln Lys Glu Cys Phe Gly
 965 970 975

Gly Ile Val Gly Ala Lys Ser Pro Tyr Val Asp Asp Met Gly Ala Ile
 980 985 990

Gly Asp Glu Val Ile Thr Leu Asn Met Ile Lys Ser Ala Pro Val Gly
 995 1000 1005

Pro Val Ala Gly Gly Ile Met Gly Cys Ile Met Val Leu Val Leu
1010 1015 1020

Ala Val Tyr Ala Tyr Arg His Gln Ile His Arg Arg Ser His Gln
1025 1030 1035

His Met Ser Pro Leu Ala Ala Gln Glu Met Ser Val Arg Met Ser
1040 1045 1050

Asn Leu Glu Asn Asp Arg Asp Glu Arg Asp Asp Asp Ser His Glu
1055 1060 1065

Asp Arg Gly Ile Ile Ser Asn Thr Arg Phe Ile Ala Ala Val Ile
1070 1075 1080

Glu Arg His Ala His Ser Pro Glu Arg Arg Arg Arg Tyr Trp Gly
1085 1090 1095

Arg Ser Gly Thr Glu Ser Asp His Gly Tyr Ser Thr Met Ser Pro
1100 1105 1110

Gln Glu Asp Ser Glu Asn Pro Pro Cys Asn Asn Asp Pro Leu Ser
1115 1120 1125

Ala Gly Val Asp Val Gly Asn His Asp Glu Asp Leu Asp Leu Asp
1130 1135 1140

Thr Pro Pro Gln Thr Ala Ala Leu Leu Ser His Lys Phe His His
1145 1150 1155

Tyr Arg Ser His His Pro Thr Leu His His Ser His His Leu Gln
1160 1165 1170

Ala Ala Val Thr Val His Thr Val Asp Ala Glu Cys
1175 1180 1185

<210> 44
<211> 410
<212> PRT
<213> Homo sapiens

<400> 44

Met Asp Glu Val Ser Leu Leu Ser Glu Leu Val Glu Ala Ala Ser Phe
1 5 10 15

Leu Gln Val Thr Ser Leu Leu Gln Leu Leu Leu Ser Gln Val Arg Leu
20 25 30

Asn Asn Cys Leu Glu Met Tyr Arg Leu Ala Gln Val Tyr Gly Leu Pro
 35 40 45
 Asp Leu Gln Glu Ala Cys Leu Arg Phe Met Val Val His Phe His Glu
 50 55 60
 Val Leu Cys Lys Pro Gln Phe His Leu Leu Gly Ser Pro Pro Gln Ala
 65 70 75 80
 Pro Gly Asp Val Ser Leu Lys Gln Arg Leu Arg Glu Ala Arg Met Thr
 85 90 95
 Gly Thr Pro Val Leu Val Ala Leu Gly Asp Phe Leu Gly Gly Pro Leu
 100 105 110
 Ala Pro His Pro Tyr Gln Gly Glu Pro Pro Ser Met Leu Arg Tyr Glu
 115 120 125
 Glu Met Thr Glu Arg Trp Phe Pro Leu Ala Asn Asn Leu Pro Pro Asp
 130 135 140
 Leu Val Asn Val Arg Gly Tyr Gly Ser Ala Ile Leu Asp Asn Tyr Leu
 145 150 155 160
 Phe Ile Val Gly Gly Tyr Arg Ile Thr Ser Gln Glu Ile Ser Ala Ala
 165 170 175
 His Ser Tyr Asn Pro Ser Thr Asn Glu Trp Leu Gln Val Ala Ser Met
 180 185 190
 Asn Gln Lys Arg Ser Asn Phe Lys Leu Val Ala Val Asn Ser Lys Leu
 195 200 205
 Tyr Ala Ile Gly Gly Gln Ala Val Ser Asn Val Glu Cys Tyr Asn Pro
 210 215 220
 Glu Gln Asp Ala Trp Asn Phe Val Ala Pro Leu Pro Asn Pro Leu Ala
 225 230 235 240
 Glu Phe Ser Ala Cys Glu Cys Lys Gly Lys Ile Tyr Val Ile Gly Gly
 245 250 255
 Tyr Thr Thr Arg Asp Arg Asn Met Asn Ile Leu Gln Tyr Cys Pro Ser
 260 265 270
 Ser Asp Met Trp Thr Leu Phe Glu Thr Cys Asp Val His Ile Arg Lys
 275 280 285

Gln Gln Met Val Ser Val Glu Glu Thr Ile Tyr Ile Val Gly Gly Cys
 290 295 300

Leu His Glu Leu Gly Pro Asn Arg Arg Ser Ser Gln Ser Glu Asp Met
 305 310 315 320

Leu Thr Val Gln Ser Tyr Asn Thr Val Thr Arg Gln Trp Leu Tyr Leu
 325 330 335

Lys Glu Asn Thr Ser Lys Ser Gly Leu Asn Leu Thr Cys Ala Leu His
 340 345 350

Asn Asp Gly Ile Tyr Ile Met Ser Arg Asp Val Thr Leu Ser Thr Ser
 355 360 365

Leu Glu His Arg Val Phe Leu Lys Tyr Asn Ile Phe Ser Asp Ser Trp
 370 375 380

Glu Ala Phe Arg Arg Phe Pro Ala Phe Gly His Asn Leu Leu Val Ser
 385 390 395 400

Ser Leu Tyr Leu Pro Asn Lys Ala Glu Thr
 405 410

<210> 45
 <211> 1101
 <212> PRT
 <213> Homo sapiens

<400> 45

Arg Arg Arg Arg Pro Ser Ser Ser Arg Arg Leu Arg Gly Arg Gly Ala
 1 5 10 15

Ala Gln Met Ala Cys Pro Ala Leu Gly Leu Glu Ala Leu Gln Pro Leu
 20 25 30

Gln Pro Glu Pro Pro Pro Glu Pro Ala Phe Ser Glu Ala Gln Lys Trp
 35 40 45

Ile Glu Gln Val Thr Gly Arg Ser Phe Gly Asp Lys Asp Phe Arg Thr
 50 55 60

Gly Leu Glu Asn Gly Ile Leu Leu Cys Glu Leu Leu Asn Ala Ile Lys
 65 70 75 80

Pro Gly Leu Val Lys Lys Ile Asn Arg Leu Pro Thr Pro Ile Ala Gly
 85 90 95

Leu Asp Asn Ile Ile Leu Phe Leu Arg Gly Cys Lys Glu Leu Gly Leu
 100 105 110

Lys Glu Ser Gln Leu Phe Asp Pro Ser Asp Leu Gln Asp Thr Ser Asn
 115 120 125

Arg Val Thr Val Lys Ser Leu Asp Tyr Ser Arg Lys Leu Lys Asn Val
 130 135 140

Leu Val Thr Ile Tyr Trp Leu Gly Lys Ala Ala Asn Ser Cys Thr Ser
 145 150 155 160

Tyr Ser Gly Thr Thr Leu Asn Leu Lys Glu Phe Glu Gly Leu Leu Ala
 165 170 175

Gln Met Arg Lys Asp Thr Asp Asp Ile Glu Ser Pro Lys Arg Ser Ile
 180 185 190

Arg Asp Ser Gly Tyr Ile Asp Cys Trp Asp Ser Glu Arg Ser Asp Ser
 195 200 205

Leu Ser Pro Pro Arg His Gly Arg Asp Asp Ser Phe Asp Ser Leu Asp
 210 215 220

Ser Phe Gly Ser Arg Ser Arg Gln Thr Pro Ser Pro Asp Val Val Leu
 225 230 235 240

Arg Gly Ser Ser Asp Gly Arg Gly Ser Asp Ser Glu Ser Asp Leu Pro
 245 250 255

His Arg Lys Leu Pro Asp Val Lys Lys Asp Asp Met Ser Ala Arg Arg
 260 265 270

Thr Ser His Gly Glu Pro Lys Ser Ala Val Pro Phe Asn Gln Tyr Leu
 275 280 285

Pro Asn Lys Ser Asn Gln Thr Ala Tyr Val Pro Ala Pro Leu Arg Lys
 290 295 300

Lys Lys Ala Glu Arg Glu Glu Tyr Arg Lys Ser Trp Ser Thr Ala Thr
 305 310 315 320

Ser Pro Leu Gly Gly Glu Arg Pro Phe Arg Tyr Gly Pro Arg Thr Pro
 325 330 335

Val Ser Asp Asp Ala Glu Ser Thr Ser Met Phe Asp Met Arg Cys Glu
 340 345 350

Glu Glu Ala Ala Val Gln Pro His Ser Arg Ala Arg Gln Glu Gln Leu
 355 360 365

Gln Leu Ile Asn Asn Gln Leu Arg Glu Glu Asp Asp Lys Trp Gln Asp
 370 375 380

Asp Leu Ala Arg Trp Lys Ser Arg Arg Arg Ser Val Ser Gln Asp Leu
 385 390 395 400

Ile Lys Lys Glu Glu Glu Arg Lys Lys Met Glu Lys Leu Leu Ala Gly
 405 410 415

Glu Asp Gly Thr Ser Glu Arg Arg Lys Ser Ile Lys Thr Tyr Arg Glu
 420 425 430

Ile Val Gln Glu Lys Glu Arg Arg Glu Arg Glu Leu His Glu Ala Tyr
 435 440 445

Lys Asn Ala Arg Ser Gln Glu Glu Ala Glu Gly Ile Leu Gln Gln Tyr
 450 455 460

Ile Glu Arg Phe Thr Ile Ser Glu Ala Val Leu Glu Arg Leu Glu Met
 465 470 475 480

Pro Lys Ile Leu Glu Arg Ser His Ser Thr Glu Pro Asn Leu Ser Ser
 485 490 495

Phe Leu Asn Asp Pro Asn Pro Met Lys Tyr Leu Arg Gln Gln Ser Leu
 500 505 510

Pro Pro Pro Lys Phe Thr Ala Thr Val Glu Thr Thr Ile Ala Arg Ala
 515 520 525

Ser Val Leu Asp Thr Ser Met Ser Ala Gly Ser Gly Ser Pro Ser Lys
 530 535 540

Thr Val Thr Pro Lys Ala Val Pro Met Leu Thr Pro Lys Pro Tyr Ser
 545 550 555 560

Gln Pro Lys Asn Ser Gln Asp Val Leu Lys Thr Phe Lys Val Asp Gly
 565 570 575

Lys Val Ser Val Asn Gly Glu Thr Val His Arg Glu Glu Glu Lys Glu
 580 585 590

Arg Glu Cys Pro Thr Val Ala Pro Ala His Ser Leu Thr Lys Ser Gln
 595 600 605

Met Phe Glu Gly Val Ala Arg Val His Gly Ser Pro Leu Glu Leu Lys
 610 615 620

Gln Asp Asn Gly Ser Ile Glu Ile Asn Ile Lys Lys Pro Asn Ser Val
 625 630 635 640

Pro Gln Glu Leu Ala Ala Thr Thr Glu Lys Thr Glu Pro Asn Ser Gln
 645 650 655

Glu Asp Lys Asn Asp Gly Gly Lys Ser Arg Lys Gly Asn Ile Glu Leu
 660 665 670

Ala Ser Ser Glu Pro Gln His Phe Thr Thr Thr Val Thr Arg Cys Ser
 675 680 685

Pro Thr Val Ala Phe Val Glu Phe Pro Ser Ser Pro Gln Leu Lys Asn
 690 695 700

Asp Val Ser Glu Glu Lys Asp Gln Lys Lys Pro Glu Asn Glu Met Ser
 705 710 715 720

Gly Lys Val Glu Leu Val Leu Ser Gln Lys Val Val Lys Pro Lys Ser
 725 730 735

Pro Glu Pro Glu Ala Thr Leu Thr Phe Pro Phe Leu Asp Lys Met Pro
 740 745 750

Glu Ala Asn Gln Leu His Leu Pro Asn Leu Asn Ser Gln Val Asp Ser
 755 760 765

Pro Ser Ser Glu Lys Ser Pro Val Thr Thr Pro Gln Phe Lys Phe Trp
 770 775 780

Ala Trp Asp Pro Glu Glu Glu Arg Arg Arg Gln Glu Lys Trp Gln Gln
 785 790 795 800

Glu Gln Glu Arg Leu Leu Gln Glu Arg Tyr Gln Lys Glu Gln Asp Lys
 805 810 815

Leu Lys Glu Glu Trp Glu Lys Ala Gln Lys Glu Val Glu Glu Glu Glu
 820 825 830

Arg Arg Tyr Tyr Glu Glu Glu Arg Lys Ile Ile Glu Asp Thr Val Val
 835 840 845

Pro Phe Thr Val Ser Ser Ser Ser Ala Asp Gln Leu Ser Thr Ser Ser
 850 855 860

Ser Met Thr Glu Gly Ser Gly Thr Met Asn Lys Ile Asp Leu Gly Asn
865 870 875 880

Cys Gln Asp Glu Lys Gln Asp Arg Arg Trp Lys Lys Ser Phe Gln Gly
885 890 895

Asp Asp Ser Asp Leu Leu Leu Lys Thr Arg Glu Ser Asp Arg Leu Glu
900 905 910

Glu Lys Gly Ser Leu Thr Glu Gly Ala Leu Ala His Ser Gly Asn Pro
915 920 925

Val Ser Lys Gly Val His Glu Asp His Gln Leu Asp Thr Glu Ala Gly
930 935 940

Ala Pro His Cys Gly Thr Asn Pro Gln Leu Ala Gln Asp Pro Ser Gln
945 950 955 960

Asn Gln Gln Thr Ser Asn Pro Thr His Ser Ser Glu Asp Val Lys Pro
965 970 975

Lys Thr Leu Pro Leu Asp Lys Ser Ile Asn His Gln Ile Glu Ser Pro
980 985 990

Ser Glu Arg Arg Lys Lys Ser Pro Arg Glu His Phe Gln Ala Gly Pro
995 1000 1005

Phe Ser Pro Cys Ser Pro Thr Pro Pro Gly Gln Ser Pro Asn Arg
1010 1015 1020

Ser Ile Ser Gly Lys Lys Leu Cys Ser Ser Cys Gly Leu Pro Leu
1025 1030 1035

Gly Lys Gly Ala Ala Met Ile Ile Glu Thr Leu Asn Leu Tyr Phe
1040 1045 1050

His Ile Gln Cys Phe Arg Cys Gly Ile Cys Lys Gly Gln Leu Gly
1055 1060 1065

Asp Ala Val Ser Gly Thr Asp Val Arg Ile Arg Asn Gly Leu Leu
1070 1075 1080

Asn Cys Asn Asp Cys Tyr Met Arg Ser Arg Ser Ala Gly Gln Pro
1085 1090 1095

Thr Thr Leu
1100

<210> 46
 <211> 1161
 <212> PRT
 <213> Homo sapiens

<400> 46

Met Lys Lys Phe Ser Arg Met Pro Lys Ser Glu Gly Gly Ser Gly Gly
 1 5 10 15

Gly Ala Ala Gly Gly Gly Ala Gly Gly Ala Gly Ala Gly Ala Gly Cys
 20 25 30

Gly Ser Gly Gly Ser Ser Val Gly Val Arg Val Phe Ala Val Gly Arg
 35 40 45

His Gln Val Thr Leu Glu Glu Ser Leu Ala Glu Gly Gly Phe Ser Thr
 50 55 60

Val Phe Leu Val Arg Thr His Gly Gly Ile Arg Cys Ala Leu Lys Arg
 65 70 75 80

Met Tyr Val Asn Asn Met Pro Asp Leu Asn Val Cys Lys Arg Glu Ile
 85 90 95

Thr Ile Met Lys Glu Leu Ser Gly His Lys Asn Ile Val Gly Tyr Leu
 100 105 110

Asp Cys Ala Val Asn Ser Ile Ser Asp Asn Val Trp Glu Val Leu Ile
 115 120 125

Leu Met Glu Tyr Cys Arg Ala Gly Gln Val Val Asn Gln Met Asn Lys
 130 135 140

Lys Leu Gln Thr Gly Phe Thr Glu Pro Glu Val Leu Gln Ile Phe Cys
 145 150 155 160

Asp Thr Cys Glu Ala Val Ala Arg Leu His Gln Cys Lys Thr Pro Ile
 165 170 175

Ile His Arg Asp Leu Lys Val Glu Asn Ile Leu Leu Asn Asp Gly Gly
 180 185 190

Asn Tyr Val Leu Cys Asp Phe Gly Ser Ala Thr Asn Lys Phe Leu Asn
 195 200 205

Pro Gln Lys Asp Gly Val Asn Val Val Glu Glu Glu Ile Lys Lys Tyr
 210 215 220

Thr Thr Leu Ser Tyr Arg Ala Pro Glu Met Ile Asn Leu Tyr Gly Gly
 225 230 235 240
 Lys Pro Ile Thr Thr Lys Ala Asp Ile Trp Ala Leu Gly Cys Leu Leu
 245 250 255
 Tyr Lys Leu Cys Phe Phe Thr Leu Pro Phe Gly Glu Ser Gln Val Ala
 260 265 270
 Ile Cys Asp Gly Asn Phe Thr Ile Pro Asp Asn Ser Arg Tyr Ser Arg
 275 280 285
 Asn Ile His Cys Leu Ile Arg Phe Met Leu Glu Pro Asp Pro Glu His
 290 295 300
 Arg Pro Asp Ile Phe Gln Val Ser Tyr Phe Ala Phe Lys Phe Ala Lys
 305 310 315 320
 Lys Asp Cys Pro Val Ser Asn Ile Asn Asn Ser Ser Ile Pro Ser Ala
 325 330 335
 Leu Pro Glu Pro Met Thr Ala Ser Glu Ala Ala Ala Arg Lys Ser Gln
 340 345 350
 Ile Lys Ala Arg Ile Thr Asp Thr Ile Gly Pro Thr Glu Thr Ser Ile
 355 360 365
 Ala Pro Arg Gln Arg Pro Lys Ala Asn Ser Ala Thr Thr Ala Thr Pro
 370 375 380
 Ser Val Leu Thr Ile Gln Ser Ser Ala Thr Pro Val Lys Val Leu Ala
 385 390 395 400
 Pro Gly Glu Phe Gly Asn His Arg Pro Lys Gly Ala Leu Arg Pro Gly
 405 410 415
 Asn Gly Pro Glu Ile Leu Leu Gly Gln Gly Pro Pro Gln Gln Pro Pro
 420 425 430
 Gln Gln His Arg Val Leu Gln Gln Leu Gln Gln Gly Asp Trp Arg Leu
 435 440 445
 Gln Gln Leu His Leu Gln His Arg His Pro His Gln Gln Gln Gln Gln
 450 455 460
 Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln
 465 470 475 480

Gln Gln Gln Gln Gln Gln His His His His His His His His Leu Leu
 485 490 495

Gln Asp Ala Tyr Met Gln Gln Tyr Gln His Ala Thr Gln Gln Gln Gln
 500 505 510

Met Leu Gln Gln Gln Phe Leu Met His Ser Val Tyr Gln Pro Gln Pro
 515 520 525

Ser Ala Ser Gln Tyr Pro Thr Met Met Pro Gln Tyr Gln Gln Ala Phe
 530 535 540

Phe Gln Gln Gln Met Leu Ala Gln His Gln Pro Ser Gln Gln Gln Ala
 545 550 555 560

Ser Pro Glu Tyr Leu Thr Ser Pro Gln Glu Phe Ser Pro Ala Leu Val
 565 570 575

Ser Tyr Thr Ser Ser Leu Pro Ala Gln Val Gly Thr Ile Met Asp Ser
 580 585 590

Ser Tyr Ser Ala Asn Arg Ser Val Ala Asp Lys Glu Ala Ile Ala Asn
 595 600 605

Phe Thr Asn Gln Lys Asn Ile Ser Asn Pro Pro Asp Met Ser Gly Trp
 610 615 620

Asn Pro Phe Gly Glu Asp Asn Phe Ser Lys Leu Thr Glu Glu Glu Leu
 625 630 635 640

Leu Asp Arg Glu Phe Asp Leu Leu Arg Ser Asn Arg Leu Glu Glu Arg
 645 650 655

Ala Ser Ser Asp Lys Asn Val Asp Ser Leu Ser Ala Pro His Asn His
 660 665 670

Pro Pro Glu Asp Pro Phe Gly Ser Val Pro Phe Ile Ser His Ser Gly
 675 680 685

Ser Pro Glu Lys Lys Ala Glu His Ser Ser Ile Asn Gln Glu Asn Gly
 690 695 700

Thr Ala Asn Pro Ile Lys Asn Gly Lys Thr Ser Pro Ala Ser Lys Asp
 705 710 715 720

Gln Arg Thr Gly Lys Lys Thr Ser Val Gln Gly Gln Val Gln Lys Gly
 725 730 735

Asn Asp Glu Ser Glu Ser Asp Phe Glu Ser Asp Pro Pro Ser Pro Lys
 740 745 750

Ser Ser Glu Glu Glu Glu Gln Asp Asp Glu Glu Val Leu Gln Gly Glu
 755 760 765

Gln Gly Asp Phe Asn Asp Asp Asp Thr Glu Pro Glu Asn Leu Gly His
 770 775 780

Arg Pro Leu Leu Met Asp Ser Glu Asp Glu Glu Glu Glu Lys His
 785 790 795 800

Ser Ser Asp Ser Asp Tyr Glu Gln Ala Lys Ala Lys Tyr Ser Asp Met
 805 810 815

Ser Ser Val Tyr Arg Asp Arg Ser Gly Ser Gly Pro Thr Gln Asp Leu
 820 825 830

Asn Thr Ile Leu Leu Thr Ser Ala Gln Leu Ser Ser Asp Val Ala Val
 835 840 845

Glu Thr Pro Lys Gln Glu Phe Asp Val Phe Gly Ala Val Pro Phe Phe
 850 855 860

Ala Val Arg Ala Gln Gln Pro Gln Gln Glu Lys Asn Glu Lys Asn Leu
 865 870 875 880

Pro Gln His Arg Phe Pro Ala Ala Gly Leu Glu Gln Glu Glu Phe Asp
 885 890 895

Val Phe Thr Lys Ala Pro Phe Ser Lys Lys Val Asn Val Gln Glu Cys
 900 905 910

His Ala Val Gly Pro Glu Ala His Thr Ile Pro Gly Tyr Pro Lys Ser
 915 920 925

Val Asp Val Phe Gly Ser Thr Pro Phe Gln Pro Phe Leu Thr Ser Thr
 930 935 940

Ser Lys Ser Glu Ser Asn Glu Asp Leu Phe Gly Leu Val Pro Phe Asp
 945 950 955 960

Glu Ile Thr Gly Ser Gln Gln Gln Lys Val Lys Gln Arg Ser Leu Gln
 965 970 975

Lys Leu Ser Ser Arg Gln Arg Arg Thr Lys Gln Asp Met Ser Lys Ser
 980 985 990

Asn Gly Lys Arg His His Gly Thr Pro Thr Ser Thr Lys Lys Thr Leu
 995 1000 1005

Lys Pro Thr Tyr Arg Thr Pro Glu Arg Ala Arg Arg His Lys Lys
 1010 1015 1020

Val Gly Arg Arg Asp Ser Gln Ser Ser Asn Glu Phe Leu Thr Ile
 1025 1030 1035

Ser Asp Ser Lys Glu Asn Ile Ser Val Ala Leu Thr Asp Gly Lys
 1040 1045 1050

Asp Arg Gly Asn Val Leu Gln Pro Glu Glu Ser Leu Leu Asp Pro
 1055 1060 1065

Phe Gly Ala Lys Pro Phe His Ser Pro Asp Leu Ser Trp His Pro
 1070 1075 1080

Pro His Gln Gly Leu Ser Asp Ile Arg Ala Asp His Asn Thr Val
 1085 1090 1095

Leu Pro Gly Arg Pro Arg Gln Asn Ser Leu His Gly Ser Phe His
 1100 1105 1110

Ser Ala Asp Val Leu Lys Met Asp Asp Phe Gly Ala Val Pro Phe
 1115 1120 1125

Thr Glu Leu Val Val Gln Ser Ile Thr Pro His Gln Ser Gln Gln
 1130 1135 1140

Ser Gln Pro Val Glu Leu Asp Pro Phe Gly Ala Ala Pro Phe Pro
 1145 1150 1155

Ser Lys Gln
 1160